

<u>MA (EU) number</u>	<u>(Invented) name</u>	<u>Strength</u>	<u>Pharmaceutical form</u>	<u>Route of administration</u>	<u>Immediate Packaging</u>	<u>Content (concentration)</u>	<u>Pack size</u>
EU/1/21/1612/001	Prevenar 20	-- ¹	Suspension for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	1 pre-filled syringe
EU/1/21/1612/002	Prevenar 20	-- ¹	Suspension for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	1 pre-filled syringe + 1 needle
EU/1/21/1612/003	Prevenar 20	-- ¹	Suspension for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	10 pre-filled syringes
EU/1/21/1612/004	Prevenar 20	-- ¹	Suspension for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	10 pre-filled syringes + 10 needles
EU/1/21/1612/005	Prevenar 20	-- ¹	Suspension for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	50 pre-filled syringes
EU/1/21/1612/006	Prevenar 20	-- ¹	Suspension for injection	Intramuscular use	pre-filled syringe (glass)	0.5 ml	50 pre-filled syringes + 50 needles

--¹ --¹

1 dose (0.5 mL) contains:

Pneumococcal polysaccharide serotype 1 ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 3 ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 4 ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 5 ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 6A ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 6B ^{1 2}	4.4 µg
Pneumococcal polysaccharide serotype 7F ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 8 ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 9V ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 10A ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 11A ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 12F ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 14 ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 15B ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 18C ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 19A ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 19F ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 22F ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 23F ^{1 2}	2.2 µg
Pneumococcal polysaccharide serotype 33F ^{1 2}	2.2 µg

¹ Conjugated to CRM₁₉₇ carrier protein (approximately 51 µg per dose)

² Adsorbed on aluminium phosphate (0.125 mg aluminium per dose)