



London, 7 September 2007  
Doc. Ref: EMEA/HMPC/359388/2006

**COMMITTEE ON HERBAL MEDICINAL PRODUCTS (HMPC)**

**LIST OF REFERENCES FOR ASSESSMENT OF:**

***PIMPINELLA ANISUM L., fructus and aetheroleum (aniseed and anise oil)***

*The EMEA acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.*

1. Abraham SK. Anti-genotoxicity of trans-anethole and eugenol in mice. *Food Chem Toxicol* 2001;39:493-8.
2. Aburjai T, Darwish RM, Al-Khalil S, Mahafzah A, Al-Abbadi A. Screening of antibiotic resistant inhibitors from local plant materials against two different strains of *Pseudomonas aeruginosa*. *J Ethnopharmacol* 2001; 76:39-44.
3. Al-Bataina BA, Maslat AO, Al.KOfahi MM. Element analysis and biological studies on ten oriental spices using XRF and Ames test, *J.Trace Elem. Med. Biol.*2003; 17(2); 85-90.
4. Albert-Puleo MJ. Fennel and anise as estrogenic agents. *J.Ethnopharmacol.* 1980;2(4);337-344.
5. Al-Harbi MM, Qureshi S, Raza M, Ahmed MM, Giangreco AB, Shah AH. Influence of anethole treatment on the tumour induced by Ehrlich ascites carcinoma cells in paw of Swiss albino mice. *Eur J Cancer Prev* 1995;4:307-18.
6. Andersen KE, Contact allergy to toothpaste flavours. *Contact Dermatitis.* 1978;4;195-198.
7. Andersen KE. Contact allergy to toothpaste flavours. *Contact Dermatitis* 1978;4:195-8.
8. Aniseed - Anisi fructus. *European Pharmacopoeia*, Council of Europe.
9. Arslan N, Gurbuz B, Sarihan EO, Bayrak A, Gumuscu A. Variation in essential oil content and composition in Turkish anise (*Pimpinella anisum L.*) populations. *Turkish J. Agricul. Forestry.*2004;28(3);173-177.
10. Baerheim Svendsen A. Uber das Vorkommen der Chlorogen- and Kaffeesaure in der Pflanzenfamilie der Umbelliferen. *Pharm Acta Helv* 1951;26:253-8
11. Balachandran B, Sivaswamy SN, Sivaranakrishan VM Genotoxic effects of some foods & food components in Swiss mice. *Indian J Med Res* 1991; 94:378-383.

12. Barnes J, Anderson LA, Phillipson JD. Aniseed. In: Herbal Medicines - A guide for healthcare professionals, 2nd ed. London-Chicago: Pharmaceutical Press, 2002:51-4.
13. Bellakhdar J, Claisse R, Fleurentin J, Younos C. Repertory of standard herbal drugs in the Moroccan Pharmacopoeia. J. Ethnopharmacol. 1991;35(2); 123-143.
14. Blumenthal M, Busse WR, Goldberg A, et al (eds). The Complete German Commission E Monographs: Therapeutic Guide to Herbal Medicines. Austin: American Botanical Council and Boston: Integrative Medicine Communications, 1998, 82-83
15. Blumenthal M, Goldberg A, Brikmann J. Herbal medicine expanded, Commission E Monograph. American Botanical Council, Austin, Texas. 2000.
16. Boskabady MH, Ramazani-Assari M. Relaxant effect of *Pimpinella anisum* on isolated guinea pig tracheal chains and its possible mechanism(s). J Ethnopharmacol 2001;74:83-8.
17. Bounds SVJ and Caldwell J. Pathways of Metabolism of (1'-14C)trans-anethole in the rat and mouse. Drug Metabolism and Disposition. 1996;
18. Boyd EM, Pearson GL. Am. J. Med. Sci. 1946;211:602-610.
19. Boyd EM, Pearson GL. On the expectorant action of volatile oils. Am J Med Sci 1946;211:602-10.
20. Boyd EM, Sheppard EP. The effect of steam inhalation of volatile oils on the output and composition of respiratory tract fluid. J Pharmacol Exp Therap 1968;163:250-6.
21. Boyd EM. Expectorants and respiratory tract fluid. Pharmacol Rev 1954;6:521-42.
22. British Herbal Pharmacopoeia Pimpinella. 1983. Bournemouth: British Herbal Medicine Association, 1983:160-1.
23. British Pharmacopoeia. The Stationary Office. 1999;260-261.
24. Buechi S, Vogel R, von Eiff MM, Ramos M, Melzer J. Open trial to assess aspects of safety and efficacy of a combined herbal cough syrup with ivy and thyme. Forsch Komplementarmed Klass Naturheilkd. 2005 Dec;12(6):328-32.
25. Burkhardt G, Reichling J, Martin R, Becker H. Terpene hydrocarbons in *Pimpinella anisum* L. Pharm Weekbl Sci. 1986; 8: 190-3.
26. Caldwell J. Perspective on the usefulness of the mouse lymphoma assay as an indicator of a genotoxic carcinogen : ten compounds which are positive in the test but are not genotoxic carcinogens. Teratogenesis, Carcinogenesis and Mutagenesis. 1993;13:185-190.
27. Caldwell J, Sutton JD. Influence of dose size on the disposition of trans-[methoxy-<sup>14</sup>C] anethole in human volunteers. Food Chem Toxicol 1988;26:87-91.
28. Caldwell J, Chan VSW, Marshall AD, Hasheminejad G , Bounds SVJ. 1'-hydroxylation is the only metabolic pathway of simple alkylbenzenes involved in their genotoxicity. Toxicologist. 1992; 12;56.
29. Cambar PJ, Aviado DM. Arch.Env,Health .1970; 20(4); 488.
30. Casciano DA, Talaska G, Clive D. The potent hepatocarcinogen methapyriline induces mutation in L5178Y mouse lymphoma cells in the absence of DNA adduct formation. Mutation Res. 1992;263:127-132.

31. Ceska O, Chaudhary SK, Warrington PJ, Ashwood-Smith MJ. Photoactive furocoumans in fruits of some umbellifers. *Phytochemistry* 1987;26:165-9.
32. Chainy GBN, Manna SK, Chaturvedi MM, Aggarwal BB. Anethole blocks both early and late cellular responses transduced by tumor necrosis factor: effect on NF-KB, AP-1, JNK, MAPKK and apoptosis. *Oncogene* 2000; 19: 2943-2950.
33. Chandler RF, Hawkes D. Aniseed - a spice, a flavor, a drug. *Can Pharm J* 1984; 117: 28-29.
34. Chopra RN, Nayer SC, Chopra IC. *Phytochemistry.2003;62 Glossary of Indian Medical Plants*. Council of Scientific and Industrial Research:New Delhi.1956.
35. Czygan F-C, Hiller K. Anisi fructus - Anis. In: Wichtl M, editor. *Teedrogen and Phytopharmaka. Ein Handbuch für- die Praxis auf wissenschaftlicher Grundlage*, 4th ed. Stuttgart: W issenschaftl iche Verlagsgesellschaft, 2002: 42-4.
36. Czygan F-C. Anis (Anisi fructus DAB 10) - *Pimpinella anisum L.* *Z Phytotherapie* 1992; 13:101-6.
37. De Smet PAGM. Legislative outlook on the safety of herbal remedies: *Pimpinella anisum*. In: De Smet PAGM, Keller K, Hansel R, Chandler RF, editors. *Adverse effects of herbal drugs*, Volume 2. Berlin: Springer, 1993:62.
38. De Vincenzi M, Silano M, Maialetti F, Scazzocchio B. Constituents of aromatic plants: II. Estragole. *Fitoterapia* 2000;71:725-9.
39. Dhar SK. Anti-fertility activity and hormonal profile of trans-anethole in rats. *Indian J. Physiol Pharmacol* 1995;39:63-7.
40. Dirks U, Herrmann K. 4-(13-D-glucoopyranosyloxy)-benzoic acid, a characteristic phenolic constituent of the Apiaceae. *Phytochemistry* 1984;23:1811-2.
41. Dirks U, Herrmann K. Hochleistungsfl (üssigkeitschromatographie der Hydroxycinnamoylchinasäuren und der 4-O-D-glucoopyranosyloxy)-benzoesäure in Gewürzen. *Z Lebensm Unters Forsch* 1984;179:12-6.
42. Dorsch W, Loew D, Meyer-Buchtela E, Schilcher H. In: *Kooperation Phytopharmaka*, editor. *Kinderdosierung von Phytopharmaka*, 3rd ed. Teil I. Empfehlungen zur Anwendung and Dosierung von Phytopharmaka, monograph ierteArzneidrogen and ihren Zubereitungen in der Padiatrie: *Foeniculi fructus (Fenchelfruchte)*. Bonn: Kooperation Phytopharmaka, 2002:70-1.
43. Drinkwater NR, Miller EC, Miller JA, Pitot HC. Hepatocarcinogenicity of estragole (1-allyl-4-methoxybenzene) and 1'-hydroxyestragole in the mouse and mutagenicity of 1'-acetoxyestragole in bacteria. *J Natl Cancer Inst.* 1976; 57: 1323-31.
44. EMEA, The European Agency for the Evaluation of Medicinal Products. Committee for Veterinary Medicinal Products: Anisi aetheroleum, summary report. .London: EMEA/MRL/413/98-FINAL; Mai 1998: 1-3 <http://www.eudra.org/emea.html>
45. El-Moghazi AM, All AA, Ross SA, Mottaleb MA. Flavonoids of *Pimpinella anisum L.* growing in Egypt. *Fitoterapia* 1979;50:267-8.
46. El-Wakeil F, Khairy M, Morsi S, Farag RS, Shihata AA, Badel AZMA. Biochemical studies on the essential oils of somefruits of Umbelliferaefamily. *Seifen-Ole-Fette-Wachse* 1986;112:77-80.

47. ESCOP (European Scientific Cooperative on Phytotherapy): Aniseed. In: Monographs on the medicinal uses of plant drugs. Argyle House, ESCOP, Exeter, UK; 1996-99.
48. European Commission: Scientific Committee on Food. Opinion of the Scientific Committee on food on estragole (1-allyl-4-methoxybenzene). 26 September 2001.
49. European Pharmacopoeia 5<sup>th</sup> ed., Aniseed - *Anisi fructus*. Council of Europe. 1/2005:0262; 1006
50. European Pharmacopoeia 5<sup>th</sup> ed., Anise oil - *Anisi aetheroleum*. Council of Europe. 1/2005:0804; 1004
51. Farag R, Daw Z, Abo-Raya S. Influence of some spice essential oils on *A. parasiticus* growth and production of aflatoxins in a synthetic medium. *J. Food Sci.*1989;54(1);74-76.
52. Fraj J, Lezaun A, Colas C, Duce F, Dominguez MA, Alonso MD. Occupational asthma induced by aniseed. *Allergy.*1996;51(5);337-339.
53. Franks A. Contact allergy to anethole in toothpaste associated with loss of taste. *Contact Dermatitis.*1998;38;354.
54. Friedman M, Henika PR, Mandrell RR. Bactericidal activities of plant essential oils and some of their isolated constituents against *Campylobacter jejuni*, *Escherichia coli*, *Listeria monocytogenes* and Fujimatu E, Ishikawa T, Kitajima J. Aromatic glucosides, alkyl glucoside and glucide from the fruit of anise. *Phytochemistry.*2003;63;609-616.
55. Fujimatu E, Ishikawa T, Kitajima J. Aromatic compound glucosides, alkyl glucoside and glucide from the fruit of anise. *Phytochemistry.*2003;63;609-616.
56. Fyfe L, Armstrong F, Stewart J. Inhibition of *Listeria monocytogenes* and *Salmonella enteritidis* by combinations of plant oils and derivatives of benzoic acid: the development of synergistic antimicrobial combinations. *Int J Antimicrobial Agents* 1998; 9:195-199.
57. Garcia-Bravo B et al. Occupational contact dermatitis from anethole in food handlers. *Contact Dermatitis* 1997; 37: 38.
58. Garcia Gonzalez JJ, Bartolomè Zavala B, Fernandez Meléndez S, Barcelò-Munoz JM, Pàez AM, Carmona-Bueno MJH, Vega-Chicote JM, Negro Carrasco M, Godoy AA, Espinosa RP.
59. Occupational rinoconjunctivitis and food allergy because of aniseed sensitization. *Ann. A llergy, Asthma Immunol.* 2002; 88; 518-522.
60. Gershbein LL. Regeneration of rat liver in the presence of essential oils and their components. *Food Cosmet Toxicol* 1977;15:173-81.
61. Ghelardini C, Galeotti N, Mazzanti G. Local anaesthetic activity of monoterpenes and phenylpropanes of essential oils. *Planta Med* 2001;67:564-6.
62. Gorelik NJ. Genotoxicity of trans-anethole *in vitro*. *Mutation Res.* 1995;326; 199-209.
63. Gosselin RE, Smith RP, Hodge HC., *Clinical Toxicology of commercial products* 5<sup>th</sup> Ed . Baltimore: Williams and Wilkins.1984; II-230.
64. Gulcin I, Oktay M, Kirecci E, Irfan Kufrevioglu O, Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extracts. *Food Chemistry.*2002;83;371-382.

65. Guneyli A, Karacali I. Effects of storage conditions on the quality of aniseed. *Ege Universitesi Ziraat Fakultesi Dergisi*. 2002;39(3);17-24.
66. Hagan EC, Hansen WH, Fitzhugh OG, Jenner PM, Jones WI, Taylor JM et al. Food flavourings and compounds of related structure. II. Subacute and chronic toxicity. *Food Cosmet Toxicol* 1967;5:141-57.
67. Haggag EG, Abou-Moustafa MA, Boucher W, Theoharides TC. The effect of a herbal water-extract on histamine release from mast cells and on allergic asthma. *J Herb Pharmacother*. 2003;3(4):41-54.
68. Hammer KA, Carson CF, Riley TV. Antimicrobial activity of essential oils and other plant extracts. *J Applied Microbiol* 1999;86:985-90.
69. Hansel R, Keller K, Rimpler H, Schneider G, editors. Pimpinella. In: *Hagers Handbuch der Pharmazeutischen Praxis*, 5th ed. Volume 6: Drogen P-Z. Berlin-Heidelberg-New York-London: Springer-Verlag, 1994:135-56.
70. Hansel R, Sticher O, Steinegger E. Anis and Anisil. In: *Pharmakognosie- Phytopharmazie*, 6th ed. Berlin-Heidelberg-New York-London: Springer-Verlag, 1999:692-5.
71. Hasan H.A.H. Inhibition of mycoflora and zearalenone on rice by selected essential oils. *Pakistan J. Sci. Ind. Res.* 1994;37(11);471-473.
72. Hausen BM. The sensitizing capacity of sulfuretin. In: *Allergiepflanzen-Pflanzenallergene*. Part]. Landsberg/ Munchen: ecomed, 1988:300.
73. Heck AM, DeWitt BA, Lukes AL: Potential interactions between alternative therapies and warfarin. *Am J Health Syst Pharm* 2000; 57: 1221-1227.
74. Heck JD, Vollmuth TA, Cifone MA, Jagannath DR, Mhyr B. and Current RD. Evaluation of food flavouring ingredients in a genetic toxicity screening battery. *Toxicologist*. 1989;9;257.
75. Howes AJ, Chan VSW, Caldwell J. Structure-specificity of the genotoxicity of some naturally occurring alkenylbenzenes determined by the unscheduled DNA synthesis assay in rat hepatocytes. *Food Chemical Toxicol*. 1990; 28;537-542.
76. Howes MJ, Houghton PJ, Barlow DJ, Pocock VJ, Milligan SR. Assessment of estrogenic activity in some common essential oil constituents. *J Pharm Pharmacol*. 2002 Nov;54(11):1521-8.
77. Hsia MTS, Adamovics JA, Kreamer BL. Microbial studies of insect growth regulators and other potential insecticidal compounds in *Salmonella typhimurium*. *Chemosphere*. 1979; 8; 521-5529.
78. Ibrahim YKE, Ogunmodede MS. Growth and survival of *Pseudomonas aeruginosa* in some aromatic waters. *Pharm Acta Helv* 1991;66:286-8.
79. Imaseki I, Katabake. *Yakugaku Zasshi*. 1962;82;1326-1328.

80. Ishidate M, Sofuni T, Yoshikawa K, Hayashi M, Nohmi T, Sawada M, Matsuoka A. Primary mutagenicity screening of food additives currently used in Japan. *Food Chem Toxicol* 1984;22:623-36.
81. Ishikawa T, Fujimatu E, Kitajima J. Water-soluble components of anise: new glucosides of anethol glycol and its related compounds. *Chem. Pharm.Bull.* 2002;50(11):1460-1466.
82. Jensen-Jarolim E, Leitner A, Hirschwehr R, Kraft D, Wuthrich B, Scheiner O, Graf J, Ebner C. Characterization of allergens in Apiaceae spices: anise, fennel, coriander and cumin. *Clin Exp Allergy.* 1997; 27:1299-306.
83. Joint FAO/WHO Expert Committee on Food Additives (JECFA) safety evaluation of certain food additives. Trans-anethole. World Health Organization, Geneva, 1999.
84. Kartnig T, Scholz G. Über einige Lipoid-Inhaltsstoffe aus den Früchten von *Pimpinella anisum* L. and *Carum carvi* L. *Fette Seifen Anstrichmittel* 1969;71:276-80.
85. Kassi E, Papoutsis Z, Fokialakis N, Messari J, Mitakou S, Moutsatsou P: Greek plant extracts exhibit selective estrogen receptor modulator (SERM)-like properties. *J.Agric. Food Chem.*2004;52:6956-6961.
86. Keller K (1992) In: De Smet PAGN, Keller K, Hänsel R, Chandler F (Hrsg.) *Adverse Effects of Herbal Drugs*, Springer, Berlin Heidelberg New York, Bd. 1, S. 135–142.
87. Kitajima J, Ishikawa T, Fujimatu E, Kondho K, Takayanagi T. Glycosides of 2-C-methyl-D-erythritol from the fruits of anise, coriander and cumin. *Phytochemistry.*2003;62:115-120.
88. Kreydiyyeh SI, Usta J. Copti,R. Effects of cinnamon, clove and some of their constituents on the Na<sup>+</sup>-K<sup>+</sup>- ATPase activity and alanine absorption in the rat jejunum. *Food Cjem.Toxicol.* 2000; 38; 755-762.
89. Kreydiyyeh SI, Usta J, Knio K., Markossian S, Dagher S. Aniseed oil increases glucose absorption and reduces urine output in the rat. *Life Sci.*2003;74:663-673.
90. Kubeczka K-H. Grundlagen der Qualitätsbeurteilung arzneilich verwendeter ätherischer Öle. *Acta Horticulturae* 1978;73:85-93.
91. Kunzemann J, Herrmann K. Isolierung and Identifizierung der Flavon(ol)-O-glykoside in Kümmel (*Carum carvi* L.), Fenchel (*Foeniculum vulgata* Mill.), Anis (*Pimpinella anisum* L.) and Koriander (*Coriandrum sativum* L.) and von Flavon-C-Glykosiden im Anis. *Z Lebensm Unters-Forsch* 1977;164:194-200.
92. Le Bourhis B. Proprietes biologiques du trans-anethole. Essai de determination de la dose journal iere acceptable. *Parfums Cosmet Sav Fr* 1973;3:450-6.
93. Leclerc Henri: *Précis de Phytothérapie*: 5<sup>e</sup> edition, Masson 1983
94. Lin FSD. Trans-anethole. In: Joint FAO/WHO Expert Committee on Food Additives. *Toxicological evaluation of certain food additives and contaminants. WHO Food Additives Series 28.* Geneva: World Health Organization 1991:135-52.
95. Linares E, Bye RA. *J.Ethnopharmacol.* 1987;19:153-183.
96. Mahady GB, Pendland SL, Stoia A, Hamill FA. *In vitro* susceptibility of *Helicobacter pylori* to botanicals used traditionally for the treatment of gastro-intestinal disorders. *Phytomedicine Suppl.* 2000;7(2); 79.

97. Marcus C, Lichtenstein EP. Interactions of naturally occurring food plant components with insecticides and pentobarbital in rats and mice. *J Agric Food Chem* 1982;30:563-8.
98. Maruzzella JC, Freundlich M. Antimicrobial substances from seeds. *J Am Pharm Assoc* 1959;48:356-8.
99. Marzin D. Recherche d'une action mutagene par le test du micronucleus chez le souris ( unpublished). Department Recherche et Essais Biologiques Stallergnes.1979.
100. Miller EC, Swanson AB, Phillips DH, Fletcher L, Liem A, Miller JA. Structure-activity studies of the carcinogenicities in the mouse and rat of some naturally occurring and synthetic alkenylbenzene derivatives related to safrole and estragole. *Cancer Res* 1983 ; 43: 1124-34.
101. Mortelmans K, Haworth S, Lawlor T, Speck W., Tainer B, and Zeiger E. Salmonella mutagenicity tests: II. Results from the testing of 270 chemicals. *Environ. Mutagenesis*. 1986; 6(suppl.7); 1-119.
102. Moyler DA.Spices-Recent Advances. In *Spices, Herbs and Edible Fungi*; Charalambous G. (Ed). Elsevier Science;London,UK.1994;1-70.
103. Muller L, Kasper P, Muller-Tegethoff K., Petr T. The genotoxic potential in vitro and in vivo of the alkylbenzene etheric oils estragole ,basil oil and trans-anethole. *Mutation Res*. 1994; 325; 129-136.
104. Mueller-Riebau F, Berger B, Yeken O. Chemical composition and fungitoxic properties to phytopathogenic fungi of essential oils of selected aromatic plants growing wild in Turkey, *J.Agric. Food Chem*. 1955; 43(8); 2262-2266
105. Muller-Limmroth W, Frohlich H-H. Wirkungsnachweis einiger phytotherapeutischer Expektorantien auf den mukoziliaren Transport. *Fortschr Med* 1980;98:95-101.
106. Murray RD. The naturally occurring coumarins. *Fortschr Chem Org Naturst*. 2002;83:1-619.
107. Nestmann ER, Lee EGH. Mutagenicity of constituents of pulp and paper mill effluent in growing cells of *Saccharomyces cerevisiae* *Mutation Res*. 1983; 119; 273-280.
108. Nestmann ER, Lee EGH, Matula TI, Douglas GR, Meuller JC. Mutagenicity of constituents of pulp and paper mill effluent using the salmonella/mammalian-microsome assay. *Mutation Res*. 1980; 79; 203—212.
109. Newall CA, Anderson LA, Phillipson JD: *Herbal Medicines. A Guide for Health Care Professionals*. The Pharmaceutical Press, London, England; 1996.
110. Newberne PM, Carlton WW, Brown WR. Histopathological evaluation of proliferative lesions in rats fed trans-anethole in chronic studies. *Food Chem. Toxicol* . 1989; 27; 21-26.
111. Newberne P, Smith RL, Doull J, Goodman JI, Munro IC, Portoghese PS, Wagner BM, Weil CS, Woods LA, Adams TB, Lucas CD, Ford RA. The FEMA GRAS assessment of trans-anethole used as a flavouring substance. *Flavour and Extract Manufacturer's Association. Food Chem Toxicol*. 1999 Jul;37(7):789-811.
112. Okazaki K, Nakayama S, Kawazoe K,Takaishi Y. Antiaggregant effect of human platelets of culinary herbs. *Phytother.Res*.1998;12(8);603-605.

113. Okuyama T, Matsuda, Masuda Y, Baba M, Masubuchi H, Adachi M, Okada Y, Hashimoto T, Zou LB, Nishino H. Studies on cancer bio-chemioprevention onf natural resources.X. Inhibitory effect of spices on TPA-enhanced 3H-choline incorporation in phospholipid of C3H10T1/2 cells and on TPA-induced ear edema. *Zhonghua Yaoxue Zashi*, 1995; 47(5),421-430.
114. Omidbaigi R, HadjiakhoondiA, Saharkhiz M. Changes in the content and chemical composition of *Pimpinella anisum* oil at various harvest time. *J. essential oil-bearing plants*.2003;6(1);46-50.
115. Ondarza M, Sanchez A. Steam distillation and supercritical fluid extraction of some Mexican spices. *Chromatographia*.1990;30;16-18.
116. Phillips DH, Reddy MV, Randerath K. 32-P-post labelling analysis of DNA adducts formed in the livers of animals treated with safrole, estragole and other naturally-occurring alkenylbenzenes.II. Newborn male B6C3F1 mice. *Carcinogenesis*. 1984; 5; 1623- 1628.
117. Pourgholami MH, Majizoob S, Javadi M, Kamalinejad M., Fanae GHR, Sayyah M. The fruit essential oil of *Pimpinella anisum* exerts anticonvulsivant effects in mice. *J. Ethnopharmacol*. 1999;66;211-215.
118. Ramadan FM, El-Zanfaly RT, El-Waked FA, Alian AM. On the antibacterial effects of some essential oils. I. Use of agar diffusion method. *Chem Mikrobiol Technol Lebensm* 1972;2:51-5.
119. Reed PM, Caldwell J. Induction of hepatic cytochrome P450 and related activities following dietary adminiustrartion of trans-anethole in SD-CD rats. *Hum. Exprimen. Toxicol*. 1992; 11; 580-581.
120. Reiter M, Brandt W. Relaxant effects on tracheal and ileal smooth muscles of the guinea pig. *Arzneim-Forsch/Drug Res* 1985;35:408-14.
121. Rodrigues VM, Paulo T, Rosa V, Marques MMO, Petenate A, Meireles MAA. Spercritical extraction of essential oil from aniseed (*Pimpinella anisum* L.) using CO<sub>2</sub>: solubility, kinetics and composition data. *J.Agric. Food Chem*. 2003;51;1518-1523.
122. Rempelberg CJM et al. Effects of the naturally occurring alkenylbenzenes eugenol and trans- anethole on drug-metabolising enzymes in the rat liver. *Food Chem Toxicol* 1993; 31: 637-645.
123. Sahraei H, Ghoshooni H, Hossein SS, Mohseni AA, Shafaghi B, Falahi M, Kamalnegad M. The effects of fruit essential oil of the *Pimpinella anisum* on the acquisition and expresswion of morphine induced conditioned place preference in mice. *J. Ethnopharmacol*.2002;80(1);43-47.
124. *Salmonella enterica*. *J. Food Prot*. 2002;65(10);1545-1560.
125. Sangster SA, Caldwell J, Hutt AJ, Anthony A, Smith RL. The metabolic disposition of [methoxy-<sup>14</sup>C]-labelled trans-anethole, estragole and p-propylanisole in human volunteers. *Xenobiotica* 1987;17:1223-32.
126. Sangster SA, Caldwell J, Smith RL, Farmer PB. Metabolism of anethole. I. Pathways of metabolism in the rat and mouse. *Food Chem Toxicol* 1984b; 22: 695-706.
127. Sangster SA, Caldwell J, Smith RL. Metabolism of anethole. II. influence of dose size on the route of metabolism of trans-anethole in the rat and mouse. *Food Chem Toxicol* 1984a; 22: 707-13.



128. Sathiyamoorthy P, Lugasi Evgil H, Schlesinger P, Kedar I, Gopas J, Pollack Y, Golan Goldhirsh A. Screening for cytotoxic and antimalarial activities in desert plants of the Negev and Bedouin market products. *Pharmaceutical Biol.* 1999;37(3);188-195.
129. Schultze W, Lange G, Kubeczka K-H. Direkte massenspektrometrische Analyse von *Pimpinella anisum* L. - Anispulver and Anisi 1. *Dtsch Apoth Ztg* 1987;127:372-8.
130. Schulz JM, Herrmann K. Analysis of hydroxybenzoic and hydroxycinnamic acids in plant material. II. Determination by gas-liquid chromatography. *J Chromatogr* 1980;195:95-104.
131. Schulz V, Hansel R, Tyler VE. *Rational Phytotherapy. A Physician's Guide to Herbal Medicine.* Springer, New York, NY; 1998: 159-160.
132. Sekizawa J, Shibamoto T. Genotoxicity of safrolerelated chemicals in microbial test systems. *Mutation Res.* 1982; 101: 127-40.
133. Shashikanth KN, Hosono A. *In vitro* mutagenicity of tropical spices to streptomycin-dependent strains of *Salmonella typhimurium* TA 98. *Agric Biol Chem* 1986;50:2947-8.
134. Shukla HS, Tripathi SC. Antifungal substance in the essential oil of anise (*Pimpinella anisum* L.). *Agric Biol Chem* 1987;51:1991-3.
135. Singh G, Kapoor IPS, Pandey SK, Singh UK, Singh RK. Studies on essential oil: Part 10; Antibacterial activity of volatile oils of some spices. *Phytother. Res.* 2002;16:680-682.
136. Soher EA. Prevention of the growth and aflatoxin production of *Apergillus flavus* by some spice essential oils. *Minufiya J. Agric. Res.* 1999;24(2);563-576.
137. Sokmen A, Jones BM, Ertkurk M. The *in vitro* antibacterial activity of Turkish Medicinal Plants. *J. Ethnopharmacol.* 1999;67(1);79-86.
138. Soliman KM, Badeaa RI. Effect of oil extracted from some medicinal plants on different mycotoxigenic fungi. *Food Chem. Toxicol.* 2002;40;1669-1675.
139. Sousa MP, Matos EO, Matos JJA, Machado MIL, Craveiro AA: *Constituentes Quimicos Ativos de Plantas edicinais Brasileiras.* Ed. UFC/laboratorio de Produtos Naturais; Fortaleza, Brazil, 1991.
140. Swanson AB, Chambliss DD, Blomquist JC, Miller EC, Miller JA. The mutagenicities of safrole, estragole, eugenol, trans-anethole and some of their known or possible metabolites for *Salmonella typhimurium* mutants. *Mutation Res.* 1979; 60; 143-145.
141. Sweetmann SE, editor. Aniseed, Anise oil. In: *Martindale - The complete drug reference*, 33rd ed. London: Pharmaceutical Press, 2002:1580.
142. Teissedre PL, Waterhouse AL. Inhibition of oxidation of human Low-Density Lipoproteins by phenolic substances in different essential oil varieties. *J. Agric. Food Chem.* 2000; 48(9);3801-3805.
143. Tisserand R, Balacs T. Anise. In: *Essential Oil Safety - A Guide for Health Care Professionals.* Edinburgh: Churchill Livingstone, 1995:117.
144. To LP, Hunt TP, Andersen ME. Mutagenicity of trans-anethole, estragole, eugenol and safrole in the Ames *Salmonella typhimurium* assay. *Bull. Environ. Contam. Toxicol.* 1982; 28; 647-654.

145. Truhaut R, LeBourhis B, Attia M, Glomot R, Newman J, Calwell J, Chronic toxicity/carcinogenicity of trans-anethole in rats. *Food Chem. Toxicol.* 1989; 27; 11-19.
146. Tuckler V, Peck C, Nesbit C., Coleman M, Weimer S, Martinez J, Ryan M, Arnold T. Seizure in an infant from aniseed oil toxicity. *J. Toxicol. Clin. Toxicol.* 2002; 40(5); 689
147. Tunc I, Berger B.M., Erler F., Dagli F. Ovicidal activity of essential oil from five plants against two stored-product insects. *J. Stored Prod. Res.* 2000;36;161-168.
148. Valnet, Jeanç *Aromathérapie*, 11<sup>e</sup> édition, Maloine 1990
149. Van Dongen K, Leusink H. The action of opium-alkaloids and expectorantson the ciliary movements in the air passages. *Arch Int Pharmacodyn* 1953;93:261-76.
150. Van Loon J. Z. *Lebensm..Unters.Forsch.* 1973;153;289-293.
151. von Skramlik E. Ober die Giftigkeit and Vertraglichkeit von atherischen Olen. *Pharmazie* 1959;14:435-45.
152. Weiss RF. *Herbal Medicine*. Gothenberg, Sweden: Ab Arcanum and Beaconsfield: Beaconsfield Publishers Ltd, 1985:203-4.
153. Weiss RF. In: *Lehrbuch der Phytotherapie*, 8th ed. Stuttgart: Hippokrates, 1997:83 and 443.
154. Wuthrich B, Dietsch R. *Schweiz. Med. Wochenschr* . 1985;115;358-364.

Other consulted bibliographic documentation:

1. Hagan EC, Hansen WH, Fitzhugh OG, Jenner PM, Jones WI, Taylor JM, Long EL, Nelson AA, Brouwer JB (1967) *Food Cosmet Toxicol* 5:141–157
2. Koller WD (1976) *Z Lebensm Unters Forsch* 160:143–147 [MEDLINE 2001]
3. Schulte KE, Rücker G, Backe W (1970) *Arch Pharm* 303:912–919
4. Boyd EM, Sheppard EP (1971) *Pharmacology* 6:65–80
5. Boyd EM (1970) *Int J Clin Pharmacol* 3:55–60
6. Boyd EM, Sheppard EP (1970) *Pharmacol* 3:345–352
7. Reiter M, Brandt W (1985) *Arzneim Forsch* 35:408–414
8. Sharaf A, Goma N (1964) *J Endocrinol* 310:289–290
9. Leclerc A, Brugere J, Luce D, Point D, Guenel P (1987) *Eur J Cancer Clin Oncol* 23:529–534 [MEDLINE 2001]
10. Abdul-Ghani A-S et al. Anticonvulsant effects of some Arab medicinal plants. *Int J Crude Drug Res* 1987; 25: 39-43