

London, 14 January 2009
Doc. Ref.: EMEA/HMPC/212894/2008

COMMITTEE ON HERBAL MEDICINAL PRODUCT (HMPC)

DRAFT

LIST OF REFERENCES FOR ASSESSMENT OF:

Taraxaci radix cum herba
***Taraxacum officinale* Weber ex Wigg., radix cum herba**
(dandelion root and herb)

The EMEA acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Ahuja TS, Freeman D, Mahnken JD, Agraharkar M, Siddiqui M, Memon A.
Predictors of the development of hyperkalemia in patients using angiotensin-converting enzyme inhibitors.
Am J Nephrol. 2000; 20: 268-272.

Akashi T, Furuno T, Takahashi T, Ayabe SI.
Biosynthesis of triterpenoids in cultured cells, and regenerated and wild plant organs of *Taraxacum officinale*.
Phytochemistry 1994; 36: 303–308.

Akhtar MS, Khan QM, Khaliq T.
Effects of *Portulaca oleracea* (kulfa) and *Taraxacum officinale* (dhudhal) in normoglycaemic and alloxan-treated hyperglycaemic rabbits.
J Pakistan Med Assoc. 1985; 35: 207–210.

Bisset NG, Phillipson JD, Czygan FC, Frohne D, Höltzel D, Nagell A., Pfander HJ, Willuhn G, Buff W. (eds)
Herbal Drugs and Phytopharmaceuticals: A Handbook for Practice on a Scientific Basis.
CRC Press, Boca Raton, Ann Arbor, London Tokyo, 1994; 486–489.

Blumenthal M, Busse WR, Goldberg A, Gruenwald J, Hall T, Riggins CW, Rister RS. (eds)
“Dandelion herb” and “Dandelion root with herb”
In: The Complete German Commission E Monographs. Therapeutic Guide to Herbal Medicines.
American Botanical Council, Austin, Texas, 1998; 118–120.

Böhm K.
Untersuchungen über choleretische Wirkungen einiger Arzneipflanzen.
Arzneim.Forsch./Drug Res. 1959; 9: 376-378.

Bradley PR ed. British Herbal Compendium Volume I, British Herbal Medicine Association,
Bournemouth 1992, 73-75.

7 Westferry Circus, Canary Wharf, London, E14 4HB, UK
Tel. (44-20) 74 18 84 00 Fax (44-20) 75 23 70 51

E-mail: mail@emea.europa.eu <http://www.emea.europa.eu>

© European Medicines Agency, 2009. Reproduction is authorised provided the source is acknowledged

British Herbal Pharmacopoeia (BHP). Fourth Edition, British Herbal Medicine Association, Exeter UK, 1996.

Budzianowski J.

Coumarins, caffeoyleltartaric acids and their artifactual methyl esters from Taraxacum officinale leaves. *Planta Medica* 1997; 63: 288- 288.

Burrows S, Simpson J.

The triterpene alcohols of Taraxacum root. The triterpene group. Part IV. *J Chem Soc. (Part II)*, 1938; 2042–2047.

Chabrol E, Charonnat R, Maximin M, Walzt R, Porin J.

L'action choléretique des Compositées.

CR Soc.Biol. 1931; 108: 1100-1102.

Choi HR, Choi JS, Han YN, Bae SJ, Chung HY.

Peroxynitrite Scavenging Activity of Herb Extracts.

Phytother Res. 2002; 16: 364–367.

Choi JH, Shin KM, Kim NY, Hong JP, Lee YS, Kim HJ, Park HJ, Lee KT.

Taraxinic Acid, a Hydrolysate of Sesquiterpene Lactone Glycoside from the Taraxacum coreanum NAKAI, Induces the Differentiation of Human Acute Promyelocytic Leukemia HL-60 Cells.

Biol Pharm Bull. 2002; 25: 1446–1450.

Clifford MN, Shutler S, Thomas GA, Ohiokpehai O.

The chlorogenic acids content of coffee substitutes.

Food Chem. 1987; 24: 99–107.

DAC 2004. Löwenzahn. Taraxaci herba cum radice.

Day JRS, Chaudhry AN, Hunt I, Taylor KM.

Heparin-induced hyperkalemia after cardiac surgery.

Ann Thorac Surg. 2002; 74: 1698–1700.

Dietary Guidelines for Americans 2005. Chapter 8: Sodium and Potassium [article online]. U.S. Department of Agriculture.

Available from <http://www.health.gov/Dietary-Guidelines/dga2005/document/html/chapter8.htm>. Accessed 7 November 2008.

Ertas ÖS, Aktaş HF, Hazedaroğlu MZ.

Analysis of Sodium and Potassium Levels in Taraxacum officinale by Flame Emission Photometry.

Acta Pharm Turcica 2005; 47: 127-130.

Evans KJ and Greenberg A.

Hyperkalemia: A Review

J Intensive Care Med 2005; 20: 272-290.

Faber K.

Dandelion—Taraxacum officinale Weber.

Pharmazie 1958; 13: 423–436.

Fernandez C, Martin-Esteban M, Fiandor A, Pascual C, Lopez Serrano C, Martinez Alzamora F, Diaz Pena JM, Ojeda Casas JA.

Analysis of cross-reactivity between sunflower pollen and other pollens of the Compositae family.

J Allergy Clin Immunol. 1993; 92: 660-667.

Funke I and Melzig MF.
Phytotherapie bei Typ-2 Diabetes mellitus.
Z Phytother. 2005; 26: 271-274.

Gehrmann B, Koch WG, Tschirch CO, Brinkmann H.
Medicinal Herbs. A Compendium.
The Haworth Medicinal Press, New York London Oxford, 2005; 64.

Gennari FJ.
Hypokalemia.
N Engl J Med. 1998; 339: 451-458.

Goldberg ED, Amosova EN, Zueva EP, Razina TG, Krylova SG, Reikhart DV.
Effects of Extracts from Medicinal Plants on the Development of Metastatic Process.
Bull Exp Biol Med. 2004; 138(3): 288-294.

Grases F, Melero G, Costa-Bauzá A, Prieto R, March JG.
Urolithiasis and phytotherapy.
Int Urol Nephrol. 1994; 26: 507–511.

Gross JL, de Azevedo MJ, Silveiro SP, Canani LH, Caramori ML, Zelmanovitz T.
Diabetic nephropathy: diagnosis, prevention, and treatment.
Diabetes Care 2005; 28: 164–176.

HagerROM 2006. Hagers Handbuch der Drogen und Arzneistoffe..) Version 5.0, Blaschek W, Ebel S, Hackenthal E, Holzgrabe U, Keller K, Reichling J, Schulz V. (eds). Deutscher Apotheker Verlag, Stuttgart, Germany.

Hagymási K, Blázovics A, Fehér J, Lugasi A, Kristó ST, Kéry A.
The in vitro effect of dandelion antioxidants on microsomal lipid peroxidation.
Phytother Res. 2000a; 14: 43–44.

Hagymási K, Blázovics A, Lugasi A, Kristó ST, Fehér J, Kéry A.
In vitro antioxidant evaluation of dandelion (*Taraxacum officinale* WEB.) water extracts.
Acta Alimentaria 2000b; 29: 1–7.

Hagymási K, Blázovics A, Lugasi A, Kristó ST , Fehér J, Kéry A.
In vitro antioxidant evaluation of dandelion (*Taraxacum officinale* WEB.) water extracts.
Accredit Quality Assurance 2000c; 5: 1-7.

Hance P, Martin Y, Vasseur J, Hilbert JL, Trotin F.
Quantification of cichory root bitterness by an ELISA for 11 β ,13-dihydrolactucin.
Food Chem. 2007; 105: 742-748.

Harkness R, Bratman S.
Mosby's Handbook of Drug-Herb and Drug-Supplement Interactions. Mosby, St. Louis London Philadelphia Sydney Toronto, 2003; 163-163.

Hausen BM.
Taraxinsäure-1'-O- β -D-glucopyranosid, das Kontaktallergen des Löwenzahns (*Taraxacum officinale* Wiggers).
Dermatosen 1982; 30: 51-53.

Hausen BM and Vieluf IK.
Allergiepflanzen/Pflanzenallergene: Handbuch und Atlas der Allergie-induzierenden Wild- und Kulturpflanzen, 2nd edition.
Landsberg München: Ecomed Verlagsgesellschaft mbH, 1997.

Hänsel R, Kartarabhardja M, Huang JT, Bohlmann F.
Sesquiterpenlacton- β -D-glucopyranoside sowie ein neues Eudesmanolid aus Taraxacum officinale.
Phytochemistry 1980; 19: 857–861.

Hänsel R.
Phytopharmaka. Grundlagen und Praxis. 2., völlig überarbeitete Auflage, unter Mitarbeit von Wiltrud Juretzek.
Springer Verlag, Berlin Heidelberg New York London, 1991; 237.

Hänsel R, Sticher O.
Pharmakognosie-Phytopharmazie. 8., überarbeitete und aktualisierte Auflage.
Springer Medizin Verlag, Heidelberg; 2007, 866-868.

He FJ and MacGregor GA.
Beneficial effects of potassium on human health.
Physiol Plant. 2008; 133: 725-735

Hemmerle H, Burger HJ, Below P, Schubert G, Rippel R, Schindler PW, Paulus E, Herling AW.
Chlorogenic acid and synthetic chlorogenic acid derivatives : novel inhibitors of hepatic glucose-6-phosphate translocase.
J Med Chem. 1997; 40: 137-145.

Hook I, McGee A, Henman M.
Evaluation of dandelion for diuretic activity and variation in potassium content.
Int J Pharmacogn. 1993; 31: 29–34.

Hussain Z, Waheed A, Qureshi RA, Burdi DK, Verspohl EJ, Khan N, Hasan M.
The Effect of Medicinal Plants of Islamabad and Murree Region of Pakistan on Insulin Secretion from INS-1 Cells.
Phytother Res. 2004; 18: 73–77.

Hyung M, -Hwan C, Chung CK.
Activation of inducible nitric oxide synthase by Taraxacum officinale in mouse peritoneal macrophages.
Gen Pharmacol. 1999; 32: 683–688.

Jarman PR, Kehely AM, Mather HM.
Hyperkalemia in diabetes: prevalence and associations.
Postgrad Med J. 1995; 71: 551-552.

Jeon HJ, Kang HJ, Jung HJ, Kang YS, Lim CJ, Kim YM, Park EH.
Anti-inflammatory activity of Taraxacum officinale.
J Ethnopharmacol. 2008; 115: 82–88.

Johnston KL, Clifford MN, Morgan LM.
Coffee acutely modifies gastrointestinal hormone secretion and glucose tolerance in humans: glycemics effects of chlorogenic acid and caffeine.
Am J Clin Nutr. 2003; 78: 728-733.

- Jovanović M, Mimica-Dukić N, Poljackić M, Boža P.
Erythema multiforme due to contact with weeds: A recurrence after patch testing.
Contact Dermatitis 2003; 48: 17-25.
- Kashiwada Y, Takanaka K, Tsukada H, Miwa Y, Taga T, Tanaka S, Ikeshiro Y.
Sesquiterpene glucosides from anti-leukotriene B4 release fraction of Taraxacum officinale.
J Asian Nat Prod Res. 2001; 3: 191–197.
- Kaurinovic B, Popovic M, Cebovic T, Mimica-Dukic N.
Effects of Calendula officinalis L. and Taraxacum officinale WEBER (Asteraceae) extracts on the production of OH radicals.
Fresenius Environ Bull. 2003; 12: 250–253.
- Keane B, Collier MH, Shann JR, Rogstad SH.
Metal content of dandelion Taraxacum officinale leaves in relation to soil contamination and airborne particulate matter.
Sci Total Environ. 2001; 281: 63-78
- Khan MG.
Cardiac Drug Therapy, 6th edition.
Philadelphia: W.B. Saunders, 2003.
- Kim HM, Lee EH, Shin TY, Lee KN, Lee JS.
Taraxacum officinale restores inhibition of nitric oxide production by cadmium in mouse peritoneal macrophages.
Immunopharmacol Immunotoxicol. 1998; 20: 283–297.
- Kim HM, Oh CH, Chung CK.
Activation of inducible nitric oxide synthase by Taraxacum officinale in mouse peritoneal macrophages.
Gen Pharmacol. 1999; 32: 683–688.
- Kim MK, Hyun SH, Choung SY.
Effect of Herbal Extract Mixtures on Serum and Liver Lipid Metabolism in Chronic Ethanol — Administered Rats.
J Health Sci. 2006; 52: 344–351.
- Kim J, Noh K, Cho M, Jang J, Song Y.
Anti-oxidative, anti-inflammatory and anti-atherogenic effects of Dandelion (Taraxacum officinale) extracts in C57BL/6 mice fed atherogenic diet.
The FASEB Journal. 2007; 21: 862.7
- Kisiel W, Barszcz B.
Further sesquiterpenoids and phenolics from Taraxacum officinale.
Fitoterapia 2000; 71: 269–273.
- Kober L, Torp-Pedersen C, Carlsen JE, Bagger H, Eliasen P, Lyngborg K, Videbaek J, Cole DS, Auclert L, Pauly NC.
A clinical trial of the angiotensin-converting-enzyme inhibitor trandolapril in patients with left ventricular dysfunction after myocardial infarction. Trandolapril Cardiac Evaluation. (TRACE) Study Group.
N Engl J Med. 1995; 333: 1670–1676.
- Koo HN, Hong SH, Song BK, Kim CH, Yoo YH, Kim HM.
Taraxacum officinale induces cytotoxicity through TNF-and IL-1secretion in Hep G2 cells.
Life Sci. 2004; 74: 1149–1157.

Kostis JB, Shelton B, Gosselin G, Goulet C, Hood WB Jr, Kohn RM, Kubo SH, Schron E, Weiss MB, Willis PW III, Young JB, Probstfield J.
Adverse effects of enalapril in the Studies of Left Ventricular Dysfunction (SOLVD).
Am Heart J. 1996; 131: 350–355.

Kristó ST, Ganzler K, Apáti P, Szöke E, Kéry A.
Analysis of antioxidant flavonoids from Asteraceae and Moraceae plants by capillary electrophoresis.
Chromatographia 2002; 56: 121–126.

Kroeber L.
Zur Pharmakologie der Inulindrogen und ihre therapeutische Verwendung.
Pharmazie 1950; 5: 122–127.

Królak E.
Accumulation of Zn, Cu, Pb and Cd by dandelion (*Taraxacum officinale* WEB.) in environments with various degrees of metallic contamination.
Pol J Environ. Studies 2003; 12: 713-721.

Kubelka W, Länger R.
Phytokodex. Pflanzliche Arzneispezialitäten in Österreich.
Österreichischer Apotheker-Verlag, Wien, 1996; 332-334.

Kuusi T, Pyysalo H, Autio K.
The bitterness properties of dandelion. II. Chemical investigations.
Lebensmittel-Wiss Technol. 1985; 18: 347–349.

Lopatina KA, Razina TG, Zueva EP, Krylova SG, Amosova EN, Guryev AM.
Plant Polysaccharides in Combined Therapy of Transplanted Tumors.
Bull Exp Biol Med. 2007; 143 Suppl 1: 24-27.

Lovell CR, Rowan M.
Dandelion dermatitis.
Contact Dermatitis 1991; 25: 185–188.

Macdonald JE and Struthers AD.
What is the optimal serum potassium level in cardiovascular patients?
J Am Coll Cardiol. 2004; 43: 155–161.

Mark KA, Brancaccio RR, Soter NA, Cohen DE.
Allergic contact and photoallergic contact dermatitis to plant and pesticide allergens.
Arch Dermatol. 1999; 135: 67–70.

Mascolo N, Autore G, Capasso F, Menghini A, Fasolo MP.
Biological screening of Italian medicinal plants for anti-inflammatory activity.
Phytother Res. 1987; 1: 28–31.

Neef H, Cilli F, Declerck PJ, Laekeman G.
Platelet antiaggregating activity of *Taraxacum officinale* Weber.
Phytother Res. 1996; 10: 138–140.

Önal S, Timur S, Okutucu B, Zihnioğlu F.
Inhibition of α -Glucosidase by Aqueous Extracts of Some Potent Antidiabetic Medicinal Plants.
Prep Biochem Biotechnol. 2005; 35: 29-36.

- Palmer BF.
Managing hyperkalemia caused by inhibitors of the renin-angiotensin-aldosterone system.
N Engl J Med. 2004; 351: 585–592.
- Park C, Zhou Y, Song YS.
Hepatoprotective effect of dandelion (*Taraxacum officinale*) against acute liver injury induced by Carbon tetrachloride in Sprague-Dawley rats.
The FASEB Journal 2007; 21: 862.8
- Park JY, Park CM, Kim JJ, Song YS.
Hepatoprotective Activity of Dandelion (*Taraxacum officinale*) Water Extract against D-Galactosamine-Induced Hepatitis in Rats.
J Korean Soc Food Sci Nutr. 2008; 37: 177-183.
- Pharmacopoeia Bohemoslovaca. Vol.2 , Ed.4, Avicenum-zdravotnické nakladatelství, Praha; 1987, 816-817.
- Popovic M, Kaurinovic B, Mimica-Dukic N, Vojinovic-Miloradov M, Dordevic A.
Combined effects of plant extracts and xenobiotics on liposomal lipid peroxidation. Part 3. Dandelion extract—CCl₄/fullnerol.
Oxidation Commun. 2001; 24: 335–343.
- Rauwald HW, Huang JT.
Taraxacoside, a type of acylated-butyrolactone glucoside from *Taraxacum officinale*.
Phytochemistry 1985; 24: 1557–1559.
- Rácz-Kotilla E, Rácz G, Solomon A.
The action of *Taraxacum officinale* extracts on the body weight and diuresis of laboratory animals.
Planta Med. 1974; 26: 212–217.
- Rote Liste 1975. Arzneimittelverzeichnis. Bundesverband der Pharmazeutischen Industrie (ed.), Aulendorf/Württ., Editio Cantor.
- Rote Liste 2006. Arzneimittelverzeichnis für Deutschland (einschliesslich EU-Zulassungen und bestimmter Medizinprodukte). Rote Liste® Service GmbH, Frankfurt/Main, Germany.
- Queralt I, Ovejero M, Carvalho ML, Marques AF, Llabrés JM.
Quantitative determination of essential and trace element content of medicinal plants and their infusions by XRF and ICP techniques.
X-Ray Spectrom. 2005; 34: 213-217.
- Saulnier P, Crosbie L, Duttaroy AK.
Inhibitory effect of aqueous extracts of some herbs on human platelet aggregation in vitro.
Platelets 2005; 8: 469-473.
- Schütz K, Kammerer DR, Carle R, Schieber A.
Characterization of phenolic acids and flavonoids in dandelion (*Taraxacum officinale* WEB. ex WIGG.) root and herb by high-performance liquid chromatography/electrospray ionization mass spectrometry.
Rapid Commun Mass Spectrom. 2005; 19: 179–186.
- Schütz K, Muks E, Carle R, Schieber A.
Separation and quantification of inulin from Compositae plants.
Biomed Chromatogr. 2006; 20: 1295–1303.

Seo SW, Koo HN, An HJ, Kwon KB, Lim BC, Seo EA, Ryu DG, Moon G, Kim HY, Kim HM, Hong SH.

Taraxacum officinale protects against cholecystokinin-induced acute pancreatitis in rats
World J Gastroenterol. 2005; 11: 597-599.

Sigstedt SC, Hooten CJ, Callewaert MC, Jenkins AR, Romero AE, Pullin MJ, Kornienko A, Lowrey TK, Slambrouck SV, Steelant WF.

Evaluation of aqueous extracts of Taraxacum officinale on growth and invasion of breast and prostate cancer cells.

Int J Oncology 2008; 32: 1085-1090.

Simándi B, Kristó ST, Kéry Á, Selmeczi LK, Kmecz I, Kemény S.
Supercritical fluid extraction of dandelion leaves.

J Supercrit Fluids 2002; 23: 135-142.

Sumanth M, Rana AC.

In vivo antioxidant activity of hydroalcoholic extract of Taraxacum officinale roots in rats.
Indian J Pharmacol. 2006; 38: 54-55.

Tita B, Bello U, Faccendini P, Bartolini R, Bolle P.

Taraxacum officinale W.: pharmacological effect of ethanol extract.

Pharmacol Res. 1993; 27: 23-24.

Trojanová I, Rada V, Kokoska L, Vlková E.

The bifidogenic effect of Taraxacum officinale root.

Fitoterapia 2004; 75: 760-763.

Tsialtas JT, Kassioumi M, Veresoglou DS.

Evaluating Leaf Ash Content and Potassium Concentration as Surrogates of Carbon Isotope Discrimination in Grassland Species.

J. Agronomy Crop Sci. 2002; 188: 168-175.

Wagner H, Wiesenauer M.

Phytotherapie. Phytopharmaka und pflanzliche Homöopatika.

Gustav Fischer Verlag. Stuttgart Jena New York, 1995, 132-134.

Westermann I, Roddick K.

Annual variation in sterol levels in leaves of Taraxacum officinale Weber.

Plant Physiol. 1981; 68: 872-875.

Welsch CA, Lachance PA, Wasserman BP.

Dietary phenolic compounds: inhibition of Na^+ dependent D-glucose uptake in rat intestinal brush border membrane vesicles.

J Nutr. 1989; 119: 1698-1704.

WHO monographs on selected medicinal plants. vol.3, WHO Press, Geneva, Switzerland, 2007; 328-337.

Wichtl M.

Teedrogen. Ein Handbuch für Apotheker und Ärzte

Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, 1984; 217-219.

Williams CA, Goldstone F, Greenham J.

Flavonoids, cinnamic acids and coumarins from the different tissues and medicinal preparations of Taraxacum officinale.

Phytochemistry 1996; 42: 121-127.

Wilman D, Riley JA.
Potential nutritive value of a wide range of grassland species.
J Agric Sci. 1993; 120: 43–49.

Wilman, D., Derrick, R.W.
Concentration and availability to sheep of N, P, K, Ca, Mg, and Na in chickweed, dandelion, dock, ribwort and spurrey, compared with perennial ryegrass.
J Agric Sci. 1994; 122: 217–223.

Wojdylo A, Oszmianski J, Czeremys R.
Antioxidant activity and phenolic compounds in 32 selected herbs.
Food Chem. 2007; 105: 940-949.

Wolbis M and Krolikowska M.
Polyphenolic compounds of dandelion (*Taraxacum officinale*).
Acta Pol Pharm. 1985; 42: 215.

Wolbis M, Królikowska M, Bednarek P.
Polyphenolic compounds in *Taraxacum officinale*.
Acta Pol Pharm. 1993; 50: 153–158.

Wu SH, Luo XD, Ma YB, Hao XJ, Wu DG.
Anti-gastric ulcer sesquiterpene lactone glycosides from *Crepis napifera*.
Yaoxue Xuebao 2002; 37: 33-36.

Yasukawa K, Akihisa T, Inoue Y, Tamura T, Yamanouchi S, Takido M.
Inhibitory effect of the methanol extracts from Compositae plants on 12-O-tetradecanoylphorbol-13-acetate-induced ear oedema in mice.
Phytother Res. 1998; 12: 484–487.

Zeller W, de Gols M, Hausen BM.
The sensitizing capacity of Compositae plants VI. Guinea pig sensitization experiments with ornamental plants and weeds using different methods.
Arch Dermatol Res. 1985; 277: 28–35.