

24 November 2014
EMA/HMPC/261891/2014
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Echinacea purpurea* (L.) Moench, herba recens

Final

The European Medicine Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Abdul MI, Jiang X, Williams KM, Day RO, Roufogalis BD, Liauw WS, et al. Pharmacokinetic and pharmacodynamic interactions of *Echinacea* and policosanol with warfarin in healthy subjects. *Br J Clin Pharmacol* 2010, 69(5):508-15

Ardjomand-Woelkart K, Kollroser M, Magnes C, Sinner F, Frye RF, Derendorf H, et al. Absolute/relative bioavailability and metabolism of dodeca-2E,4E,8Z,10E/Z-tetraenoic acid isobutylamides (tetraenes) after intravenous and oral single doses to rats. *Planta Med* 2011, 77(16):1794-1799

Ardjomand-Woelkart K, Kollroser M, Derendorf H, Bauer R, Butterweck V. Herb-Drug Interactions: Effects of *Echinacea* preparations on cytochrome P450 activities in rats. *Planta Med* 2012, 78(11):PH10

Ayrancı U, Son N, Son O. Prevalence of nonvitamin, nonmineral supplement usage among students in a Turkish university. *BMC Public Health* 2005, 5(47)

Baetgen D. Erfolge in der Keuchhusten – Behandlung mit Echinacin. *Therapiewoche* 1984, 34:5115-5119

Baetgen D. Behandlung der akuten Bronchitis im Kindesalter – Praxisstudie mit einem Immunstimulans aus *Echinacea purpurea*. *Pädiatrie* 1988, 1:66-70

Barnes J, Anderson L, Gibbons S, Phillipson JD. *Echinacea* species (*Echinacea angustifolia* (DC.) Hell., *Echinacea pallida* (Nutt.) Nutt., *Echinacea purpurea* (L.) Moench): a review of their chemistry, pharmacology and clinical properties. *J Pharm Pharmacol* 2005, 57(8):929-954

Barnes J, Anderson LA, Phillipson JD. *Herbal Medicines*. 3rd ed. Pharmaceutical Press, London 2007

Barrett B. Medicinal properties of *Echinacea*: a critical review. *Phytomedicine* 2003, 10(1):66-86

Barrett B. Efficacy and safety of *Echinacea* in treating upper respiratory tract infections in children: a randomized controlled trial. *J Pediatr* 2004, 145(1):135-136

Barrett B, Vohmann M, Calabrese C. *Echinacea* for upper respiratory infection. *J Fam Practice* 1999, 48(8):628-635



Barrett B, Brown RL, Locken K, Maberry R, Bobula JA, D'Alessio D. Treatment of the common cold with unrefined *Echinacea*. A randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 2002, 137(12):939-946

Bauer R. [New knowledge regarding the effect and effectiveness of *Echinacea purpurea* extracts.] *Wien Med Wochenschr* 2002, 152(15-16):407-411 [German]

Bauer R, Jurcic K, Puhlmann J, Wagner H. [Immunologic *in vivo* and *in vitro* studies on *Echinacea* extracts.] *Arzneimittelforschung* 1988a, 38(2):276-281 [German]

Bauer R, Remiger P, Wagner H. *Echinacea*. Comparative TLC and HPLC analysis of herbal drugs from *Echinacea purpurea*, *E. pallida* and *E. angustifolia*. Part 3. *Dtsch Apoth Ztg* 1988b, 128:174-180

Bauer R, Remiger P, Jurcic K, Wagner H. Beeinflussung der Phagozytose-Aktivität durch *Echinacea*-Extrakte. *Z Phytother* 1989, 10:43-48

Beuscher N, Bodinet C, Willigmann I, Egert D. Immonomodulierende Eigenschaften von Wurzelextrakten verschiedener *Echinacea*-Arten. *Z Phytother* 1995, 16:157-166

Binns SE, Purgina B, Bergeron C, Smith ML, Ball L, Baum BR, et al. Light-mediated antifungal activity of *Echinacea* extracts. *Planta Med* 2000, 66(3):241-244

Binns SE, Livesey JF, Arnason JT, Baum BR. Phytochemical variation in *Echinacea* from roots and flowerheads of wild and cultivated populations. *J Agric Food Chem* 2002a, 50:3673-3687

Binns SE, Hudson J, Merali S, Arnason JT. Antiviral activity of characterized extracts from *Echinacea* spp. (Heliantheae: Asteraceae) against *Herpes simplex* virus (HSV-I). *Planta Med* 2002b, 68(9):780-783

Blaschek W, Ebel S, Hackenthal E, Holzgrabe U, Keller K, Reichling J, et al., editors. HagerROM 2004: Hagers Handbuch der Drogen und Arzneistoffe. CD-ROM 5. 1st ed. Springer-Verlag, Heidelberg 2004

Blaschek W, Ebel S, Hilgenfeldt U, Holzgrabe U, Reichling J, Schulz, V, editors. HagerROM 2008: Hagers Enzyklopädie der Arzneistoffe und Drogen. 6th ed. Springer-Verlag, Stuttgart 2008

Blumenthal M. *Echinacea* trial. *Altern Ther Health Med* 2004, 10(2):12-14

Blumenthal M, Goldberg A, Brinckmann J, editors. Herbal medicine. Expanded Commission E Monographs. The American Botanical Council, Austin Texas, 2000

Bossaer JB, Odle BL. Probable etoposide interaction with *Echinacea*. *J Diet Suppl* 2012, 9(2):90-95

Bourne LC, Rice-Evans C. Bioavailability of ferulic acid. *Biochem Biophys Res Commun* 1998, 253:222-227

Brinkeborn RM, Shah DV, Degenring FH. Echinaforce and other *Echinacea* fresh plant preparations in the treatment of the common cold. A randomized, placebo-controlled, double-blind clinical trial. *Phytomedicine* 1999, 6(1):1-6

Burger RA, Torres AR, Warren RP, Caldwell VD, Hughes BG. *Echinacea*-induced cytokine production by human macrophages. *Int J Immunopharmacol* 1997, 19(7):371-378

Cheminat A, Zawatzky R, Becker H, Bruillard R. Caffeoyl conjugates from *Echinacea* species: Structures and biological activity. *Phytochemistry* 1988, 27(9):2787-2794

Chow G, Johns T, Miller SC. Dietary *Echinacea purpurea* during murine pregnancy: effect on maternal hemopoiesis and fetal growth. *Biol Neonate* 2006, 89(2):133-138

Clifford LJ, Nair MG, Rana J, Dewitt DL. Bioactivity of alkamides isolated from *Echinacea purpurea* (L.) Moench. *Phytomedicine* 2002, 9(3):249-253

Currier NL, Miller SC. Natural killer cells from aging mice treated with extracts from *Echinacea purpurea* are quantitatively and functionally rejuvenated. *Exp Geron* 2000, 35(5):627-639

Cuzzolin L, Francini-Pesenti F, Verlato G, Joppi M, Baldelli P, Benoni G. Use of herbal products among 392 Italian pregnant women: focus on pregnancy outcome. *Pharmacopidemiol Drug Saf* 2010, 19(11):1151-1158

Dietz B, Heilmann J, Bauer R. Absorption of dodeca-2E,4E,8Z,10E/Z-tetraenoic acid isobutylamides after oral application of *Echinacea purpurea* tincture. *Planta Med* 2001, 67(9):863-864

Dorsch W. Klinische Anwendung von Extrakten aus *Echinacea purpurea* oder *Echinacea pallida*. *Zeitschrift für ärztliche Fortbildung* 1996, 90:117-122

Dorsch W, Loew D, Meyer-Buchtela E, Schilcher H. Kinderdosierungen von Phytopharmaka. Vol 1. 3rd ed. Kooperation Phytopharmaka, Bonn 2002

ESCOP Monographs 2nd ed. *Echinaceae purpureae herba - Purple coneflower herb*. European Scientific Cooperative on Phytotherapy, editor. Thieme, Stuttgart 2003, 129-136

Facino RM, Carini M, Aldini G, Saibene L, Pietta P, Mauri P. Echinacoside and caffeoyl conjugates protect collagen from free radica induced degradation: a potential use of *Echinacea* extracts in the prevention of skin photocarcinogenesis. *Planta Med* 1995, 61(6):510-514

Firenzuoli F, Gori L. *Echinacea* for treating colds in children. *JAMA* 2004, 291(11):1323-1324

Foster S. The Purple Coneflowers. American Botanical Council, Austin 1996

Gallo M, Sarkar M, Au W, Pietrzak K, Comas B, Smith M, et al. Pregnancy outcome following gestational exposure to *Echinacea*: a prospective controlled study. *Arch Intern Med* 2000, 160(20):3141-3143

Gan XH, Zhang L, Heber D, Bonavida B. Mechanism of activation of human peripheral blood NK cells at the single cell level by *Echinacea* water soluble extracts: recruitment of lymphocyte-target conjugates and killer cells and activation of programming for lysis. *Int Immunopharmacol* 2003, 3(6):811-824

German homeopathic pharmacopoeia 8th ed. Suppl. *Echinacea purpurea*. Deutscher Apotheker Verlag, Stuttgart 2011, 1-3

Gertsch J, Schoop R, Kuenzle U, Suter A. *Echinacea* alkylamides modulate TNF-a gene expression via cannabinoid receptor CB2 and multiple signal transduction pathways. *FEBS Letters* 2004, 577(3):563-569

Giancaspro G. *Echinacea purpurea* Aerial Parts. *Pharmacopeial Forum* 2004, 30(2):557-560

Giles JT, Palat CT, Chien SH, Chang ZG, Kennedy DT. Evaluation of *Echinacea* for treatment of the common cold. *Pharmacotherapy* 2000, 20(6):690-697

Glavač NK, Košir IJ, Rode J, Kreft S. Optimization and use of a spectrophotometric method for determining polysaccharides in *Echinaceae purpurea*. *Cent Eur J Biol* 2012, 7(1):126-131

Godwin M, Crellin J, Mathews M, Chowdhury NL, Newhook LA, Pike A, et al. Use of natural health products in children: Survey of parents in waiting rooms. *Can Fam Physician* 2013, 59(8):e364-371

Goel V, Chang C, Slama JV, Barton R, Bauer R, Gahler R, et al. Alkylamides of *Echinacea purpurea* stimulate alveolar macrophage function in normal rats. *Int Immunopharmacol* 2002, 2(2-3):381-387

Goel V, Lovlin R, Barton R, Lyon MR, Bauer R, Lee TD, et al. Efficacy of a standardized *Echinacea* preparation (Echinilin) for the treatment of the common cold: a randomized, double-blind, placebo-controlled trial. *J Clin Pharm Ther* 2004, 29(1):75-83

Goel V, Lovlin R, Chang C, Slama JV, Barton R, Gahler R, et al. A proprietary extract from the *Echinacea* plant (*Echinacea purpurea*) enhances systemic immune response during a common cold. *Phytother Res* 2005, 19(8):689-694

Goey AK, Rosing H, Meijerman I, Sparidans RW, Schellens JH, Beijnen JH. The bioanalysis of the major *Echinacea purpurea* constituents dodeca-2E,4E,8Z,10E/Z-tetraenoic acid isobutylamides in human plasma using LC-MS/MS. *J Chromatogr B* 2012, 902:151-156

Gorski JC, Huang SM, Pinto A, Hamman MA, Hilligoss JK, Zaheer NA, et al. The effect of *Echinacea* (*Echinacea purpurea* root) on cytochrome P450 activity *in vivo*. *Clin Pharmacol Ther* 2004, 75:89-100

Götte K, Roschke I. A supportive measure in the treatment of acute infections of the respiratory tract in children with recurring infections of the upper respiratory tract. *PÄD Praktische Pädiatrie* 2001, 7:1-7

Gotti R, Pomponio R, Bertucci C, Cavrini V. Simultaneous analysis of the lipophilic and hydrophilic markers of *Echinacea* plant extracts by capillary electrophoresis. *J Sep Sci* 2002, 25(15-17):1079-1086

Grimm W, Müller HH. A randomized controlled trial of the effect of fluid extract of *Echinacea purpurea* on the incidence and severity of colds and respiratory infections. *Am J Med* 1999, 106(2):138-143

Grünwald J, Büttel K. Der europäische Markt für Phytotherapeutika. Zahlen, Trends und Analysen. *Pharm Ind* 1996, 3:209-214

Gurley BJ, Gardner SF, Hubbard MA, Williams DK, Gentry WB, Carrier J, et al. *In vivo* assessment of botanical supplementation on human cytochrome P450 phenotypes: *Citrus aurantium*, *Echinacea purpurea*, milk thistle, and saw palmetto. *Clin Pharmacol Ther* 2004, 76(5):428-440

Gurley BJ, Swain A, Williams DK, Barone G, Battu SK. Gauging the clinical significance of P-glycoprotein-mediated herb-drug interactions: Comparative effects of St. John's wort, *Echinacea*, clarithromycin, and rifampin on digoxin pharmacokinetics. *Mol Nutr Food Res* 2008, 52(7):772-779

Hänsel R, Keller K, Rimpler H, Schneider G, editors. Hagers Handbuch der Pharmazeutischen Praxis. Drogen E-O. Vol 5. 5th ed. Springer-Verlag, New York 1993

Hansen TS, Nielsen OG. *In vitro* CYP3A4 metabolism: Inhibition by *Echinacea purpurea* and choice of substrate for the evaluation of herbal inhibition. *Basic Clin Pharmacol Toxicol* 2008, 103(5):445-449

Hansen TS, Nielsen OG. *Echinacea purpurea* and P-glycoprotein drug transport in Caco-2 cells. *Phytother Res* 2009, 23(1):86-91

Heinen-Kammerer T, Holtmannspotter C, Schnabel S, Motzkat K, Kiencke P, Rychlik R. Effectiveness of echinacin in therapy of chronic recurrent respiratory disease. *Gesundheitswesen* 2005, 67(4):296-301

Hermann R, Richter O. Clinical evidence of herbal drugs as perpetrators of pharmacokinetic drug interactions. *Planta Med* 2012, 78(13):1458-1477

Hoheisel O, Sandberg M, Bertram S, Bulitta M, Schafer M. Echinagard treatment shortens the course of the common cold: a double-blind, placebo-controlled clinical trial. *Eur J Clin Res* 1997, 9:261-268

Holst L, Wright D, Haavik S, Nordeng H. Safety and efficacy of herbal remedies in obstetrics-review and clinical implications. *Midwifery* 2011, 27(1):80-86

Huntley AL, Coon JT, Ernst E. The safety of herbal medicinal products derived from *Echinacea* species. A systematic review. *Drug Safety* 2005, 28(5):387-400

Izzo AA, Ernst E. Interactions between herbal medicines and prescribed drugs: an updated systematic review. *Drugs* 2009, 69(13):1777-1798

Izzo AA, Ernst E. Interactions between herbal medicines and prescribed drugs. A systematic review. *Drugs* 2001, 61(15):2163-2175

Jacobsson I, Jönsson AK, Gerdén B, Hägg S. Spontaneously reported adverse reactions in association with complementary and alternative medicine substances in Sweden. *Pharmacoepidemiol Drug Saf* 2009, 18(11):1039-1047

Jager H, Meinel L, Dietz B, Lapke C, Bauer R, Merkle HP, et al. Transport of Alkamides from *Echinacea* Species through Caco-2 Monolayers. *Planta Med* 2002, 68(5):469-471

Jawad M, Schoop R, Suter A, Klein P, Eccles R. Safety and Efficacy Profile of *Echinacea purpurea* to Prevent Common Cold Episodes: A Randomized, Double-Blind, Placebo-Controlled Trial. *Evid Based Complement Alternat Med* 2012, Article ID 841315

Jeschke E, Ostermann T, Lüke C, Tabali M, Kröz M, Bockelbrink A, et al. Remedies containing Asteraceae extracts: a prospective observational study of prescribing patterns and adverse drug reactions in German primary care. *Drug Safety* 2009, 32(8):691-706

Karsch-Völk M, Barrett B, Kiefer D, Bauer R, Ardjomand-Woelkart K, Linde K. *Echinacea* for preventing and treating the common cold. *The Cochrane Database of Systematic Reviews* 2014, 2:CD000530

Kemp DE, Franco KN. Possible leukopenia associated with long-term use of *Echinacea*. *J Am Board Farm Pract* 2002, 15(5):417-419

Kim L, Wollner D, Anderson P, Brammer D. *Echinacea* for treating colds in children. *JAMA* 2004, 291(11):1323-1324

Kinkel HJ, Plate M, Tullner HU. Objektivierbare wirkung von Echinacin®-Salbe auf die Wunderheilung. [Effect of Echinacin ointment in healing of skin lesions.] *Med Klin* 1984, 79(21):580-583

Kommission E. Aufbereitungsmonographie *Echinaceae purpurea*. *Bundesanzeiger* vom 02/03/1989

Kreft S. Cichoric acid content and biomass production of *Echinaceae purpurea* plants cultivated in Slovenia. *Pharm Biol* 2005, 43(8):662-665

Lenk W. Akute toxicität von verschiedenen polysacchariden aus *Echinaceae purpurea* an der Maus. [Acute toxicity of various polysaccharides from *Echinaceae purpurea* in the mouse.] *Z Phytother* 1989, 10:49-51

Le Tourneau M. *Echinacea* fails to show efficacy in treating colds in a pediatric population. *Altern Ther Health Med* 2004, 10(1):16

Liasos G, Elefsiniotis R, Todorova R, Moulakakis A. Severe thrombotic thrombocytopenic purpura (TTP) induced or exacerbated by the immunostimulatory herb *Echinacea*. *Am J Hematol* 2006, 81(3):224

Linde K, Barrett B, Wölkart K, Bauer R, Melchart D. *Echinacea* for preventing and treating the common cold. *The Cochrane Database of Systematic Reviews* 2006, 1: CD000530

Lindenmuth GF, Lindenmuth EB. The efficacy of *Echinacea* compound herbal tea preparation on the severity and duration of upper respiratory and flu symptoms: a randomized, double-blind placebo-controlled study. *J Altern Complement Med* 2000, 6(4):327-334

Liu R, Tam TW, Mao J, Saleem A, Krantis A, Arnason JT, et al. The effect of natural health products and traditional medicines on the activity of human hepatic microsomal-mediated metabolism of oseltamivir. *J Pharm Pharmaceut Sci* 2010, 13(1):43-55

Logan JL, Ahmed J. Critical hypokalemic renal tubular acidosis due to Sjogren's syndrome: association with the purported immune stimulant *Echinacea*. *Clin Rheumatol* 2003, 22(2):158-159

Luettig B, Steinmuller C, Gifford GE, Wagner H, Lohmann-Matthes ML. Macrophage activation by the polysaccharide arabinogalactan isolated from plant cell cultures of *Echinaceae purpurea*. *J Natl Cancer Inst* 1989, 81(9):669-675

Manček B, Kretf S. Determination of cichoric acid content in dried press juice of purple coneflower (*Echinaceae purpurea*) with capillary electrophoresis. *Talanta* 2005, 66(5):1094-1097

Martindale: The Complete Drug Reference. 36th ed. Pharmaceutical Press, London 2009

Maskatia ZK, Baker K. Hypereosinophilia associated with *Echinacea* use. *South Med J* 2010, 103(11):1173-1174

Matthias A, Addison RS, Penman KG, Dickinson RG, Bone KM, Lehmann RP. *Echinacea* alkamide disposition and pharmacokinetics in humans after tablet ingestion. *Life Sci* 2005a, 77(16):2018-2029

Matthias A, Blanchfield JT, Penman KG, Toth I, Lang CS, De Voss JJ, et al. Permeability studies of alkylamides and caffeic acid conjugates from *Echinacea* using a Caco-2 cell monolayer model. *J Clin Pharm Ther* 2004, 29(1):7-13

Matthias A, Gillam EMJ, Penman KG, Matovic NJ, Bone KM, De Voss JJ, et al. Cytochrome P450 enzyme-mediated degradation of *Echinacea* alkylamides in human liver microsomes. *Chem Biol Interact* 2005b, 155(1-2):62-70

Melchart D, Linde K, Worku F, Bauer R, Wagner H. Immunomodulation with *Echinacea* – a systematic review of controlled clinical trials. *Phytomedicine* 1994, 1:245-254

Melchart D, Clemm C, Weber B, Draczynski T, Worku F, Linde K, et al. Polysaccharides isolated from *Echinaceae purpurea* herba cell cultures to counteract undesired effects of chemotherapy - a pilot study. *Phytother Res* 2002, 16(2):138-142

Mengs U, Clare CB, Poiley JA. Toxicity of *Echinaceae purpurea*. Acute, subacute and genotoxicity studies. *Arzneimittel forschung* 1991, 41(10):1076-1081

Merali S, Binns S, Paulin-Levasseur M, Ficker C, Smith M, Baum B, et al. Antifungal and Anti-inflammatory Activity of the Genus *Echinacea*. *Pharm Biol* 2003, 41(6):412-420

Mishima S, Saito K, Maruyama H, Inoue M, Yamashita T, Ishida T, et al. Antioxidant and immuno-enhancing effects of *Echinaceae purpurea*. *Biol Pharm Bull* 2004, 27(7):1004-1009

Modarai M, Yang M, Suter A, Kortenkamp A, Heinrich M. Metabolomic profiling of liquid *Echinacea* medicinal products with *in vitro* inhibitory effects on cytochrome P450 3A4 (CYP3A4). *Planta Med* 2010, 76(4):378-385

Moltó J, Valle M, Miranda C, Cedeño S, Negredo E, Barbanoj MJ, et al. Herb-drug interaction between *Echinaceae purpurea* and darunavir-ritonavir in HIV-infected patients. *Antimicrob Agents Chemother* 2011, 55(1):326-330

Moltó J, Valle M, Miranda C, Cedeño S, Negredo E, Clotet B. Herb-drug interaction between *Echinaceae purpurea* and etravirine in HIV-infected patients. *Antimicrob Agents Chemother* 2012, 56(10):5328-5331

Mrozikiewicz PM, Bogacz A, Karasiewicz M, Mikolajczak PL, Ozarowski M, Seremak-Mrozikiewicz A, et al. The effect of standardized *Echinaceae purpurea* extract on rat cytochrome P450 expression level. *Phytomedicine* 2010, 17(10):830-833

Mullins RJ. *Echinacea*-associated anaphylaxis. *Med J Aust* 1998, 168(4):170-171

Mullins RJ, Heddle R. Adverse reactions associated with *Echinacea*: the Australian experience. *Ann Allergy Asthma Immunol* 2002, 88(1):42-51

Nordeng H, Bayne K, Havnen GC, Paulsen BS. Use of herbal drugs during pregnancy among 600 Norwegian women in relation to concurrent use of conventional drugs and pregnancy outcome. *Complement Ther Clin Pract* 2011, 17(3):147-151

Nordeng H, Havnen GC. Use of herbal drugs in pregnancy: a survey among 400 Norwegian women. *Pharmacoepidemiology Drug Safety* 2004, 13(6):371-380

Ondrzek RR, Chan PJ, Patton WC, King A. Inhibition of human sperm motility by specific herbs used in alternative medicine. *J Assist Reprod Genet* 1999, 16(2):87-91

Orinda VD, Diederich J, Wacker A. Antiviral activity of compounds of *Echinaceae purpurea*. *Arzneimittelforschung* 1973, 23:1119-1120

Parnham MJ. Benefit-risk assessment of the squeezed sap of the purple coneflower (*Echinaceae purpurea*) for long term oral immunostimulation. *Phytomedicine* 1996, 3(1):95-102

Pasco D, Pugh ND, Khan I, Moraes R. Immunostimulatory agents in botanical. US Patent 20050002962 A1, 2005

Pellati F, Benvenuti S, Magro L, Melegari M, Soragni F. Analysis of phenolic compounds and radical scavenging activity of *Echinacea* spp. *J Pharm Biomed Anal* 2004, 35(2):289-301

Penzak SR, Robertson SM, Hunt JD, Chairez C, Malati CY, Alfaro RM, et al. *Echinaceae purpurea* significantly induces cytochrome P450 3A activity but does not alter lopinavir-ritonavir exposure in healthy subjects. *Pharmacotherapy* 2010, 30(8):797-805

Perri D, Dugoua JJ, Mills E, Koren G. Safety and efficacy of *Echinacea* (*Echinacea angustifolia*, *E. purpurea* and *E. pallida*) during pregnancy and lactation. *Can J Clin Pharmacol* 2006, 13(3):e262-267

Pharmeuropa 26.3. European Directorate for the Quality of Medicines & HealthCare, Strasbourg 2014, 303-312

Pietta P, Mauri P, Bauer R. MEKC Analysis of different *Echinacea* species. *Planta Med* 1998, 64(7):649-652

Pleschka S, Stein M, Schoop R, Hudson JB. Anti-viral properties and mode of action of standardized *Echinaceae purpurea* extract against highly pathogenic avian influenza virus (H5N1, H7N7) and swine-origin H1N1 (S-OIV). *Virol J* 2009, 6:197

Proksch A, Wagner H. Structural analysis of a 4-O-methylglucuronoarabinoxylan with immunostimulating activity from *Echinaceae purpurea*. *Phytochemistry* 1987, 26(7):1989-1993

Pugh ND, Balachandran P, Lata H, Dayan FE, Joshi V, Bedir E, et al. Melanin: dietary mucosal immune modulator from *Echinacea* and other botanical supplements. *Int Immunopharmacol* 2005, 5(4):637-647

Pugh ND, Jackson CR, Pasco DS. Total bacterial load within *Echinaceae purpurea*, determined using a new PCR-based quantification method, is correlated with LPS levels and *in vitro* macrophage activity. *Planta Med* 2013, 79(1):9-14

Randolph RK, Gellenbeck K, Stonebrook K, Brovelli E, Qian Y, Bankaitis-Davis D, et al. Regulation of human immune gene expression as influenced by a commercial blended *Echinacea* product: preliminary studies. *Exp Biol Med* 2003, 228(9):1051-1056

Raner GM, Cornelious S, Moulick K, Wang Y, Mortenson A, Cech NB. Regulation of human immune gene expression as influenced by a commercial blended *Echinacea* product: preliminary studies. *Food Chem Toxicol* 2007, 45(12):2359-2365

Razinger-Mihovec B. Vpliv socialno-demografskih značilnosti in zdravstvenega stanja populacije na uporabo zdravil rastlinskega izvora v sloveniji. [The influence of socio-demographic characteristics and health status of the population on the use of herbal medicines in Slovenia.] Master thesis, University of Ljubljana 2007

Reisch J, Spitzner W, Schulte KE. The problem of the microbiological activity of simple acetylene compounds. *Arzneimittelforschung* 1967, 17:816-824

Rininger JA, Kickner S, Chigurupati P, McLean A, Franck Z. Immunopharmacological activity of *Echinacea* preparations following simulated digestion on murine macrophages and human peripheral blood mononuclear cells. *J Leukoc Biol* 2000, 68(4):503-510

Roesler J, Steinmuller C, Kiderlen A, Emmendorffer A, Wagner H, Lohmann-Matthes ML. Application of purified polysaccharides from cell cultures of the plant *Echinaceae purpurea* to mice mediates protection against systemic infections with *Listeria monocytogenes* and *Candida albicans*. *Int J Immunopharmacol* 1991, 13(1):27-37

Ross SM. A standardized *Echinacea* extract demonstrates efficacy in the prevention and treatment of colds in athletes. *Holist Nurs Pract* 2010, 24(2):107-109

Rote liste. Preparativerzeichnis. Bundesverband der Pharmazeutischen Industrie e. V. Editio Cantor, Aulendorf 1961, 295

Rote liste. Wundbehandlungsmittel. Bundesverband der Pharmazeutischen Industrie e. V. Editio Cantor, Aulendorf 1984, 7

Schaffer DM, Gordon NP, Jensen CD, Avins AL. Nonvitamin, nonmineral supplement use over a 12-month period by adult members of a large health maintenance organization. *J Am Diet Assoc* 2003, 103(11):1500-1505

Schimmer O, Abel G, Behninger C. Untersuchungen zur gentoxischen potenzeines neutralen Polysaccharids aus *Echinacea*-Gewebe-kulturen in menschlichen lymphozytenkulturen. [Investigations on the genotoxic potency of a neutral polysaccharide from *Echinacea* cell cultures in cultured human lymphocytes.] *Z Phytother* 1989, 10:39-42

Schoop R, Büechi S, Suter A. Open, multicenter study to evaluate the tolerability and efficacy of Echinaforce Forte tablets in athletes. *Adv Ther* 2006, 23(5):823-833

Schulte KE, Rücker G, Boehme R. Polyacetylenes as compounds of the roots of burs. *Arzneimittelforschung* 1967a, 17:829-833

Schulte KE, Rücker G, Perlick J. The presence of polyacetylene compounds in *Echinaceae purpurea* and *Echinacea angustifolia*. *Arzneimittelforschung* 1967b, 17(7):825-829

Schulzen B, Bulitta M, Ballering-Bruhl B, Koster U, Schafer M. Efficacy of *Echinaceae purpurea* in patients with a common cold. A placebo-controlled, randomised, double-blind clinical trial. *Arzneimittelforschung* 2001, 51(11):563-568

Schwarz S, Knauth M, Schwab S, Walter-Sack I, Bonmann E, Storch-Hagenlocher B. Acute disseminated encephalomyelitis after parenteral therapy with herbal extracts: a report of two cases. *J Neurol Neurosurg Psychiatry* 2000, 69(4):516-518

Schwarz E, Metzler J, Diedrich JP, Freudenstein J, Bode C, Bode JC. Oral administration of freshly expressed juice of *Echinaceae purpurea* herbs fail to stimulate the nonspecific immune response in healthy young men: results of a double-blind, placebo-controlled crossover study. *J Immunother* 2002, 25(5):413-420

Schwarz E, Parlesak A, Henneicke-von Zepelin HH, Bode JC, Bode C. Effect of oral administration of freshly pressed juice of *Echinaceae purpurea* on the number of various subpopulations of B- and T-lymphocytes in healthy volunteers: results of a double-blind, placebo-controlled cross-over study. *Phytomedicine* 2005, 12(9):625-631

Shah SA, Sander S, White CM, Rinaldi M, Coleman CI. Evaluation of *Echinacea* for the prevention and treatment of the common cold: a meta-analysis. *Lancet Infect Dis* 2007, 7(7):473-480

Sharma M, Schoop R, Hudson JB. The efficacy of *Echinacea* in a 3-D tissue model of human airway epithelium. *Phytother Res* 2010, 24(6):900-904

Skwarek T, Tynecka Z, Glowniak K, Lutostanska E. *Echinacea L.*—Inducer of interferons. *Herba Pol* 1996, 42:110-117

Sperber SJ, Shah LP, Gilbert RD, Ritchey TW, Monto AS. *Echinaceae purpurea* for prevention of experimental rhinovirus colds. *Clin Infect Dis* 2004, 38(10):1367-1371

Soon SL, Crawford RI. Recurrent erythema nodosum associated with *Echinacea* herbal therapy. *J Am Acad Dermatol* 2001, 44(2):298-299

Steinmüller C, Roesler J, Gröttrup E, Franke G, Wagner H, Lohmann-Matthes ML. Polysaccharides isolated from plant cell cultures of *Echinaceae purpurea* enhance the resistance of immunosuppressed mice against systemic infections with *Candida albicans* and *Listeria monocytogenes*. *Int J Immunopharmacol* 1993, 15(5):605-614

Stimpel M, Proksch A, Wagner H, Lohmann-Matthes ML. Macrophage activation and induction of macrophage cytotoxicity by purified polysaccharide fractions from the plant *Echinaceae purpurea*. *Infect Immun* 1984, 46(3):845-849

Stotzem CD, Hungerland U, Mengs U. Influence of *Echinaceae purpurea* on the phagocytosis of human granulocytes. *Med Sci Res* 1992, 20:719-720

Tamta H, Pugh ND, Balachandran P, Moraes R, Sumiyanto J, Pasco DS. Variability in *in vitro* macrophage activation by commercially diverse bulk *Echinacea* plant material is predominantly due to bacterial lipoproteins and lipopolysaccharides. *J Agric Food Chem* 2008, 56(22):10552-10556

Taylor JA, Weber W, Calabrese C. *Echinacea* for treating colds in children. *JAMA* 2004, 291(11):1324

Taylor JA, Weber W, Standish L, Quinn H, Goesling J, McGann M, et al. Efficacy and safety of *Echinacea* in treating upper respiratory tract infections in children: a randomized controlled trial. *JAMA* 2003, 290(21):2824-2830

Tsai YL, Chiou SY, Chan KC, Sung JM, Lin SD. Caffeic acid derivatives, total phenols, antioxidant and antimutagenic activities of *Echinaceae purpurea* flower extracts. *LWT - Food Sci Technol* 2012, 46(1):169-176

Tünnerhoff FK, Schwabe HK. Studies in human beings and animals on the influence of *Echinacea* extracts on the formation of connective tissue following the implantation of fibrin.

Arzneimittelforschung 1956, 6:330-334

Turner RB, Riker DK, Gangemi JD. Ineffectiveness of *Echinacea* for prevention of experimental rhinovirus colds. *Antimicrob Agents Ch* 2000, 44(6):1708-1709

Uang YS, Hsu KY. A dose-dependent pharmacokinetic study on caffeic acid in rabbits after intravenous administration. *Biopharm Drug Dispos* 1997, 18(8):727-736

Vimalanathan S, Kang L, Amiguet VT, Livesey J, Arnason JT, Hudson J. *Echinaceae purpurea*. Aerial Parts Contain Multiple Antiviral Compounds. *Pharm Biol* 2005, 43(9):740-745

Vonau B, Chard S, Mandalia S, Wilkinson D, Barton SE. Does the extract of the plant *Echinaceae purpurea* influence the clinical course of recurrent genital herpes? *Int J STD AIDS* 2001, 12(3):154-158

Wacker A, Hilbig W. Virusemmung mit *Echinaceae purpurea*. [Virus-inhibition by *Echinaceae purpurea*.] *Planta Med* 1978, 33:89-102

Wagner H, Proksch A, Riess-Maurer I, Vollmar A, Odenthal S, Stuppner H, et al. Immunstimulierend wirkende polysaccharide (heteroglycane) aus höheren pflanzen. [Immunostimulating action of polysaccharides (heteroglycans) from higher plants.] *Arzneimittelforschung* 1985, 35(7):1069-1075

Wagner H, Stuppner H, Schafer W, Zenk M. Immunologically active polysaccharides of *Echinacea purpurea* cell cultures. *Phytochemistry* 1988, 27(1):119-126

Washam C. *Echinacea* no cure-all for kids. *Environ Health Perspect* 2004, 112(8):A466

Weber W, Taylor JA, Stoep AV, Weiss NS, Standish LJ, Calabrese C. *Echinacea purpurea* for prevention of upper respiratory tract infections in children. *J Altern Comp Med* 2005, 11(6):1021-1026

Westendorf J, Czok G. Untersuchungen der pharmacokinetik von 14C-Zimtsäure-Derivaten an Ratten. [Pharmacokinetic of ¹⁴C labelled derivatives of cinnamic acid on rats.] *Z Ernährungswiss* 1978, 17:26-36

WHO monographs on selected medicinal plants. Vol 1. Herba Echinaceae Purpureae. World Health Organization. Geneva 1999, 136-144

Woelkart K, Koidl C, Grisold A, Gangemi JD, Turner RB, Marth E, et al. Bioavailability and pharmacokinetics of alkamides from the roots of *Echinacea angustifolia* in humans. *J Clin Pharmacol* 2005, 45(6):683-689

Yale SH, Liu K. *Echinaceae purpurea* therapy for the treatment of the common cold: a randomized, double-blind, placebo-controlled clinical trial. *Arch Intern Med* 2004, 164(11):1237-1241

Yale SH, Glurich I. Analysis of the inhibitory potential of *Ginkgo biloba*, *Echinaceae purpurea*, and *Serenoa repens* on the metabolic activity of cytochrome P450 3A4, 2D6, and 2C9. *J Altern Complement Med* 2006, 11(3):433-439

Zoutewelle G, Van Wijk R. Effects of *Echinaceae purpurea* extracts on fibroblast populated collagen lattice contraction. *Phytother Res* 1990, 4(2):77-81