

30 May 2017
EMA/HMPC/424585/2016
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Echinacea purpurea* (L.) Moench, radix

Final

The European Medicines Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Agnew LL, Guffogg SP, Matthias A, Lehmann RP, Bone KM, Watson K. Echinacea intake induces an immune response through altered expression of leucocyte hsp70, increased white cell counts and improved erythrocyte antioxidant defences. *J Clin Pharm Ther* 2005, 30: 363-369

Ardjomand-Woelkart K, Kollroser M, Magnes C, Sinner F, Frye RF, Derendorf H et al. Absolute/relative bioavailability and metabolism of dodeca-2E, 4E, 8Z, 10E/Z-tetraenoic acid isobutylamides (tetraenes) after intravenous and oral single doses to rats. *Planta Med* 2011, 77(16): 1794-1799

Balciunaite G, Juodsnukyte J, Savickas A, Ragazinskiene O, Siatkute L, Zvirblyte G, Mistiniene E, Savickiene N. Fractionation and evaluation of proteins in roots of *Echinacea purpurea* (L.) Moench. *Acta Pharm* 2015, 65(4):473-479.

Barbour EK, Abou Assi CA, Shaib H, Hamadeh S, Murtada M, Mahmoud G et al. Evaluation of a *Salmonella Enteritidis* vaccine and related ELISA for respective induction and assessment of acquired immunity to the vaccine and/or *Echinacea purpurea* in Awassi Ewes. *Vaccine* 2015, 33(19): 2228-2231

Barnes J, Anderson LA, Phillipson JD. *Echinacea*. In: *Herbal Medicines*. 3rd ed. Pharmaceutical Press, London 2007, UK: 217-236

Barnes J, Anderson, LA, Gibbons S, Phillipson JD. *Echinacea* species (*Echinacea angustifolia* (DC.) Hell., *Echinacea pallida* (Nutt.) Nutt., *Echinacea purpurea* (L.) Moench): A review of their chemistry, pharmacology and clinical properties. *J Pharm Pharmacol* 2005, 57: 929-954

Barrett B. Medicinal properties of *Echinacea*: a critical review. *Phytomedicine* 2003, 10: 66-86

Barrett B, Brown R, Rakel D, Mundt M, Bone K, Barlow S et al. *Echinacea* for treating the common cold: a randomized trial. *Ann Intern Med* 2010, 153(12): 769-777



Barrett B, Brown RL, Locken K, Maberry R, Bobula JA, D'Alessio D. Treatment of the common cold with unrefined Echinacea. A randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 2002, 137: 939-946

Barrett B, Vohmann M, Calabrese C. Echinacea for upper respiratory tract infection. *J Fam Pract* 1999, 48: 628-635

Barth A, Hovhannisyan A, Jamalyan K, Naramyan M. Antitussive effect of a fixed combination of *Justicia adhatoda*, *Echinacea purpurea* and *Eleutherococcus senticosus* extracts in patients with acute upper respiratory tract infection: A comparative, randomized, double-blind, placebo-controlled study. *Phytomedicine* 2015, 22(13): 1195-1200

Bauer R, Liersch R. Echinacea. In: Hagers Buch der Pharmazeutischen Praxis. Drogen E-O. Vol 5. 5th ed. Springer-Verlag, Berlin 1993, Germany: 1-34

Bauer R, Remiger P. TLC and HPLC Analysis of Alkaloids in Echinacea Drugs. *Planta Med* 1989a, 55: 367-371

Bauer R, Remiger P, Jurcic K, Wagner H. Beeinflussung der Phagozytose-Aktivität durch Echinacea-Extrakte. *Z Phytother* 1989b, 10: 43-48

Benson JM, Pokorny AJ, Rhule A, Wenner CA, Kandhi V, Cech NB et al. Echinacea purpurea extracts modulate murine dendritic cell fate and function. *Food Chem Toxicol* 2010, 48(5): 1170-1177

Beuscher N, Bodinet I, Willigmann D, Egert. Immunomodulating Properties of Root Extracts from Various Echinacea Species. *Z Phytother* 1995, 16: 157-166

Beuscher N, Schet KH, Bodinet C, Egert D. Modulation of host resistance by polymer substances from *Baptisia tinctoria* and *Echinacea purpurea*. In: Immunotherapeutic prospects of infectious diseases (Masih KN, Lange W, eds.). Springer-Verlag, Berlin-Heidelberg 1990, Germany: 59-63

Binns SE, Hudson J, Merali S, Arnason JT. Antiviral activity of characterized extracts from Echinacea spp. (Heliantheae: Asteraceae) against herpes simplex virus (HSV-1). *Planta Med* 2002, 68: 780-783

Binns SE, Purgina B, Bergeron C, Smith ML, Ball L, Baum BR, et al. (2000). Light-mediated antifungal activity of Echinacea extracts. *Planta Med* 66: 241-244

Blumenthal M, Goldberg A, Brinckmann J. *Echinacea purpurea* root. In: Herbal Medicine - Expanded Commission E monographs, American Botanical Council, Austin 2000, TX, USA: 98-102

Bodinet C, Beuscher N. Antiviral and Immunological Activity of Glycoproteins from *Echinacea purpurea* Root. *Planta Med* 1991, 57(2): 33-34

Bodinet C, Freudenstein J. Effect of an orally applied aqueous-ethanolic extract of a mixture of *Thujae occidentalis* herba, *Baptisiae tinctoriae* radix, *Echinaceae purpureae* radix and *Echinaceae pallidae* radix on antibody response against sheep red blood cells in mice. *Planta Med* 1999, 65: 695-699

Bodinet K. Immunopharmacological Investigations of a Plant Immunomodulator [Dissertation]. Ernst-Moritz-Arndt-Universität, Greifswald 1999, Germany

Bossaer JB, Odle BL. Probable etoposide interaction with *Echinacea*. *J Diet Suppl* 2012, 9(2): 90-95

Bradley P. Echinacea. In: British Herbal Compendium. Vol. 2. British Herbal Medicine Association, Bournemouth 2006, UK: 129-141

Bräunig B, Dorn M, Limburg E, Knick E, Bausendorf. *Echinacea purpurea* radix: zur Stärkung der körpereigenen Abwehr bei grippalen Infekten. *Z Phytother* 1992, 13: 7-13

Brinkeborn RM, Shah DV, Degenring FH. Echinaforce® and other Echinacea fresh plant preparations in the treatment of the common cold. A randomized, placebo-controlled, double-blind clinical trial.

Phytomedicine 1999, 6: 1-6

Brousseau M, Miller SC. Enhancement of natural killer cells and increased survival of aging mice fed daily Echinacea root extract from youth. *Biogerontology* 2005, 6: 157-163

Budzinski JW, Foster BC, Vandenoever S, Arnason JT. An in vitro evaluation of human cytochrome P450 3A4 inhibition by selected commercial herbal extracts and tinctures. *Phytomedicine* 2000, 7: 273-282

Cech NB, Kandhi V, Davis JM, Hamilton A, Eads D, Laster SM. *Echinacea* and its alkylamides: effects on the influenza A-induced secretion of cytokines, chemokines, and PGE₂ from RAW 264.7 macrophage-like cells. *Int Immunopharmacol* 2010, 10(10): 1268-1278

Chen Y, Fu T, Tao T, Yang J, Chang Y, Wang M, et al. Macrophage Activating Effects of New Alkamides from the Roots of Echinacea Species. *J Nat Prod* 2005, 68: 773-776

Chicca A, Adinolfi B, Martinotti E, Fogli S, Breschi MC, Pellati F, et al. Cytotoxic effects of Echinacea root hexanic extracts on human cancer cell lines. *J Ethnopharmacol* 2007, 110: 148-153

Chicca A, Pellati F, Adinolfi B, Matthias A, Massarelli I, Benvenuti S, et al. Cytotoxic activity of polyacetylenes and polyenes isolated from roots of *Echinacea pallida*. *Br J Pharmacol* 2008, 153: 879-885

Clifford LJ, Nair MG, Rana J, Dewitt DL. Bioactivity of alkamides isolated from *Echinacea purpurea* (L.) Moench. *Phytomedicine* 2002, 9: 249-253

Currier NL, Miller SC. Natural killer cells from aging mice treated with extracts from *Echinacea purpurea* are quantitatively and functionally rejuvenated. *Exp Gerontol* 2000, 35: 627-639

Currier NL, Miller SC. *Echinacea purpurea* and melatonin augment natural-killer cells in leukemic mice and prolong life span. *J Altern Complement Med* 2001, 7: 241-251

Currier NL, Miller SC. The effect of immunization with killed tumor cells, with/without feeding of *Echinacea purpurea* in an erythroleukemic mouse model. *J Altern Complement Med* 2002, 8: 49-58

Cuzzolin L, Francini-Pesenti F, Verlato G, Joppi M, Baldelli P, Benoni G. Use of herbal products among 392 Italian pregnant women: focus on pregnancy outcome. *Pharmacoepidemiol Drug Saf.* 2010, 19(11):1151-1158

Dalby-Brown L, Barsett H, Landbo AK, Meyer AS, Mølgaard P. Synergistic antioxidative effects of alkamides, caffeic acid derivatives, and polysaccharide fractions from *Echinacea purpurea* on *in vitro* oxidation of human low-density lipoproteins. *J Agric Food Chem* 2005, 53: 9413-9423

Di Carlo G, Nuzzo I, Capasso R, Sanges MR, Galdiero E, Capasso F, et al. Modulation of apoptosis in mice treated with *Echinacea* and St. John's wort. *Pharmacol Res* 2003, 48: 273-277

El-Ashmawy NE, El-Zamarany EA, Salem ML, El-Bahrawy HA, Al-Ashmawy GM. In vitro and in vivo studies of the immunomodulatory effect of *Echinacea purpurea* on dendritic cells. *J Gen Eng Biotech* 2015, 13(2): 185-192

European Pharmacopoeia. Purple Coneflower Root. In: European Pharmacopoeia, 6.0., Vol. 2, Council of Europe. 01/2008: 1824

European Scientific Cooperative On Phytotherapy (ESCOP). *Echinaceae purpureae radix*. In: ESCOP Monographs, 2nd ed., Thieme, New York 2003, NY, USA: 137-141

European Scientific Cooperative On Phytotherapy (ESCOP). Echinaceae purpurea radix. In: ESCOP Monographs, 2nd ed., Suppl. 2009, Thieme, New York 2009, NY, USA: 102-109

Fast DJ, Balles JA, Scholten JD, Mulder T, Rana J. Echinacea purpurea root extract inhibits TNF release in response to Pam3Csk4 in a phosphatidylinositol-3-kinase dependent manner. *Cell Immunol* 2015, 297(2): 94-99

Freeman C, Spelman K. A critical evaluation of drug interactions with Echinacea spp. *Mol Nutr Food Res* 2008, 52: 789-798

Gallo M, Sarkar M, Au W, Pietrzak K, Comas B, Smith M, et al. Pregnancy outcome following gestational exposure to Echinacea: a prospective controlled study. *Arch Intern Med* 2000, 160(20): 3141-3143

Gertsch J, Schoop R, Kuenzle U, Suter A. Echinacea alkylamides modulate TNF-alpha gene expression via cannabinoid receptor CB2 and multiple signal transduction pathways. *FEBS Lett* 2004, 577: 563-569

Godwin M, Crellin J, Mathews M, Chowdhury NL, Newhook LA, Pike A, et al. Use of natural health products in children: survey of parents in waiting rooms. *Can Fam Physician*. 2013, 59(8):364-371

Goel V, Chang C, Slama J, Barton R, Bauer R, Gahler R, et al. Echinacea stimulates macrophage function in the lung and spleen of normal rats. *J Nutr Biochem* 2002a, 13: 487-492

Goel V, Chang C, Slama JV, Barton R, Bauer R, Gahler R, et al. Alkylamides of *Echinacea purpurea* stimulate alveolar macrophage function in normal rats. *Int Immunopharmacol* 2002b, 2: 381-387

Goel V, Lovlin R, Barton R, Lyon MR, Bauer R, Lee TD, et al. Efficacy of a standardized Echinacea preparation (Echinilin) for the treatment of the common cold: a randomized, double-blind, placebo-controlled trial. *J Clin Pharm Ther* 2004, 29: 75-83

Goel V, Lovlin R, Chang C, Slama JV, Barton R, Gahler R, et al. A proprietary extract from the Echinacea plant (*Echinacea purpurea*) enhances systemic immune response during a common cold. *Phytother Res* 2005, 19: 689-694

Gorski JC, Huang SM, Pinto A, Hamman MA, Hilligoss JK, Zaheer NA, et al. The effect of Echinacea (*Echinacea purpurea* root) on cytochrome P450 activity in vivo. *Clin Pharmacol Ther* 2004, 75: 89-100

Gryzlak BM, Wallace RB, Zimmerman MB, Nisly NL. National surveillance of herbal dietary supplement exposures: the poison control center experience. *Pharmacoepidemiol Drug Saf* 2007, 16: 947-957

Haller CA, Anderson IB, Kim SY, Blanc PD. An evaluation of selected herbal reference texts and comparison to published reports of adverse herbal events. *Adverse Drug React Toxicol* 2002, Rev 21: 143-150

Hellum BH, Nilsen OG. The *in vitro* inhibitory potential of trade herbal products on human CYP2D6-mediated metabolism and the influence of ethanol. *Basic Clin Pharmacol Toxicol* 2007, 101: 350-358

Henneicke-von Zepelin H, Hentschel C, Schnitker J, Kohnen R, Köhler G, Wüstenberg P. Efficacy and safety of a fixed combination phytomedicine in the treatment of the common cold (acute viral respiratory tract infection): results of a randomised, double blind, placebo controlled, multicentre study. *Curr Med Res Opin* 1999, 15: 214-227

Hohmann J, Rédei D, Forgo P, Szabó P, Freund TF, Haller J, et al. Alkamides and a neolignan from *Echinacea purpurea* roots and the interaction of alkamides with G-protein-coupled cannabinoid receptors. *Phytochemistry* 2011, 72(14-15): 1848-1853

Holst L, Wright D, Haavik S, Nordeng H. Safety and efficacy of herbal remedies in obstetrics-review and clinical implications. *Midwifery* 2011, 27(1):80-86.

Hou CC, Chen CH, Yang NS, Chen YP, Lo CP, Wang SY et al. Comparative metabolomics approach coupled with cell- and gene-based assays for species classification and anti-inflammatory bioactivity validation of Echinacea plants. *J Nutr Biochem* 2010, 21(11): 1045-1059

Hou CC, Huang CC, Shyur LF. Echinacea alkamides prevent lipopolysaccharide/D-galactosamine-induced acute hepatic injury through JNK pathway-mediated HO-1 expression. *J Agric Food Chem* 2011, 59(22): 11966-11974

Hu C, Kitts DD. Studies on the antioxidant activity of Echinacea root extract. *J Agric Food Chem* 2000, 48: 1466-72

Huntley AL, Thompson Coon J, Ernst E. The safety of herbal medicinal products derived from Echinacea species: a systematic review. *Drug Saf* 2005, 28: 387-400

Jager H, Meinel L, Dietz B, Lapke C, Bauer R, Merkle HP, et al. Transport of Alkamides from Echinacea Species through Caco-2 Monolayers. *Planta Med* 2002, 68: 469-471

Jawad M, Schoop R, Suter A, Klein P, Eccles R. Safety and Efficacy Profile of Echinacea purpurea to Prevent Common Cold Episodes: A Randomized, Double-Blind, Placebo-Controlled Trial. *Evid Based Compl Altern Med* 2012, 841315

Jurcic K, Melchart D, Holzmann M, Martin P, Bauer R, Doenecke A, et al. Zwei Probandenstudien zur Stimulierung der Granulozytenphagozytose durch Echinacea-Extrakt-haltige Präparate. *Z Phytother* 1989, 10: 67-70

Karsch-Völk M, Barrett B, Kiefer D, Bauer R, Arjomand-Woelkart K, Linde K. Echinacea for preventing and treating the common cold. *Cochrane Database of Systematic Reviews* 2014, 2: 1-91

Kotowska D, El-Houri RB, Borkowski K, Petersen RK, Fretté XC, Wolber G et al. Isomeric C12-alkamides from the roots of Echinacea purpurea improve basal and insulin-dependent glucose uptake in 3T3-L1 adipocytes. *Planta Med* 2014, 80(18): 1712-1720

LaLone CA, Hammer KD, Wu L, Bae J, Leyva N, Liu Y, et al. Echinacea species and alkamides inhibit prostaglandin E(2) production in RAW264.7 mouse macrophage cells. *J Agric Food Chem* 2007, 55: 7314-7322

Linde K, Barret B, Bauer R, Melchart D, Woelkart K. Echinacea for preventing and treating the common cold. *Cochrane Database of Systematic Reviews*, Issue 1, In: The Cochrane Library 2008, Issue 3, John Wiley & Sons, Ltd., Chichester 2006, UK: 1-88

Lindenmuth GF, Lindenmuth EB. The efficacy of Echinacea compound herbal tea preparation on the severity and duration of upper respiratory and flu symptoms: a randomized, double-blind placebo-controlled study. *J Alt Complement Med* 2000, 6: 327-334

Maffei Facino R, Carini M, Aldini G, Saibene L, Pietta P, Mauri P. Echinacoside and caffeoyl conjugates protect collagen from free radical-induced degradation: a potential use of Echinacea extracts in the prevention of skin photodamage. *Planta Med* 1995, 61: 510-514

Matthias A, Addison RS, Agnew LL, Bone KM, Watson K, Lehmann RP. Comparison of Echinacea alkylamide pharmacokinetics between liquid and tablet preparations. *Phytomedicine* 2007, 14: 587-590

Matthias A, Addison RS, Penman KG, Dickinson RG, Bone KM, Lehmann RP. Echinacea alkamide disposition and pharmacokinetics in humans after tablet ingestion. *Life Sci* 2005b, 77: 2018-2029

Matthias A, Banbury L, Bone KM, Leach DN, Lehmann RP. Echinacea alkylamides modulate induced immune responses in T-cells. *Fitoterapia* 2008, 79: 53-58

Matthias A, Blanchfield JT, Penman KG, Toth I, Lang CS, De Voss JJ, et al. Permeability studies of alkylamides and caffeic acid conjugates from Echinacea using a Caco-2 cell monolayer model. *J Clin Pharm Ther* 2004, 29: 7-13

Matthias A, Gillam EM, Penman KG, Matovic NJ, Bone KM, De Voss JJ, et al. Cytochrome P450 enzyme-mediated degradation of *Echinacea* alkylamides in human liver microsomes. *Chem Biol Interact* 2005a, 155: 62-70

Mazza G, Cottrell T. Volatile components of roots, stems, leaves, and flowers of *Echinacea* species. *J Agric Food Chem* 1999, 47: 3081-3085

McCann DA, Solco A, Liu Y, Macaluso F, Murphy PA, Kohut ML, et al. Cytokine- and interferon-modulating properties of *Echinacea* spp. root tinctures stored at -20°C for 2 years. *J Interferon Cytokine Res* 2007, 27: 425-436

Melchart D, Linde K, Fischer P, Kaesmayr J. Echinacea for preventing and treating the common cold (Cochrane review). In: The Cochrane Library, Issue 2, John Wiley & Sons, Chichester 2004, UK

Melchart D, Linde K, Worku F, Bauer R, Wagner H. Immunomodulation with *Echinacea*. A systematic review of controlled clinical trials. *Phytomed* 1994, 1: 245-254

Melchart D, Linde K, Worku F, Sarkady L, Holzmann M, Jurcic K, et al. Results of five randomized studies on the immunomodulatory activity of preparations of *Echinacea*. *J Altern Complem Med* 1995, 1: 145-160

Melchart D, Walther E, Linde K, Brandmaier R, Lersch C. Echinacea root extracts for the prevention of upper respiratory tract infections: a double-blind, placebo-controlled randomized trial. *Arch Fam Med* 1998, 7: 541-545

Merali S, Binns S, Paulin-Levasseur M, Ficker C, Smith M, Baum B, et al. Antifungal and Anti-inflammatory Activity of the Genus *Echinacea*. *Pharm Biol* 2003, 41: 412-420

Moltó J, Valle M, Miranda C, Cedeño S, Negredo E, Barbanjo MJ et al. Herb-drug interaction between *Echinacea purpurea* and darunavir-ritonavir in HIV-infected patients. *Antimicrob Agents Chemother* 2011, 55(1): 326-330

Moltó J, Valle M, Miranda C, Cedeño S, Negredo E, Clotet B. Herb-drug interaction between *Echinacea purpurea* and etravirine in HIV-infected patients. *Antimicrob Agents Chemother* 2012, 56(10): 5328-5331

Müller-Jakic B, Breu W, Pröbstle A, Redl K, Greger H, Bauer R. In vitro inhibition of cyclooxygenase and 5-lipoxygenase by alkamides from *Echinacea* and *Achillea* species. *Planta Med* 1994, 60: 37-40

Mullins RJ. Echinacea-associated anaphylaxis. *Med J Austr* 1998, 168: 170-171

Mullins RJ, Heddle R. Adverse reactions associated with *Echinacea*: the Australian experience. *Ann Allergy Asthma Immunol* 2002, 88: 42-51

Nordeng H, Bayne K, Havnen GC, Paulsen BS. Use of herbal drugs during pregnancy among 600 Norwegian women in relation to concurrent use of conventional drugs and pregnancy outcome. *Complement Ther Clin Pract*. 2011, 17(3):147-151.

PDR. *Echinacea*. In: Gruenwald J, Brendler T, Jaenicke C (editors). PDR for herbal medicines, 4th Edition, Thomson Healthcare Inc., Montvale 2007, NJ, USA: 266-276

- Pellati F, Benvenuti S, Magro L, Melegari M, Soragni F. Analysis of phenolic compounds and radical scavenging activity of *Echinacea* spp. *J Pharm Biomed Anal* 2004, 35: 289-301
- Perri D, Dugoua JJ, Mills E, Koren G. Safety and efficacy of *Echinacea* (*Echinacea angustifolia*, *E. purpurea* and *E. pallida*) during pregnancy and lactation. *Can J Clin Pharmacol* 2006, 13(3): 262-267
- Pugh ND, Jackson CR, Pasco DS. Total bacterial load within *Echinacea purpurea*, determined using a new PCR-based quantification method, is correlated with LPS levels and in vitro macrophage activity. *Planta Med* 2013, 79(1): 9-14
- Raduner S, Majewska A, Chen JZ, Xie XQ, Hamon J, Faller B, et al. Alkylamides from *Echinacea* are a new class of cannabinomimetics. Cannabinoid type 2 receptor-dependent and -independent immunomodulatory effects. *J Biol Chem* 2006, 281: 14192-14206
- Raso GM, Pacilio M, Di Carlo G, Esposito E, Pinto L, Meli R. In-vivo and in-vitro anti-inflammatory effect of *Echinacea purpurea* and *Hypericum perforatum*. *J Pharm Pharmacol* 2002, 54: 1379-1383
- Rauš K, Pleschka S, Klein P, Schoop R, Fisher P. Effect of an *Echinacea*-Based Hot Drink Versus Oseltamivir in Influenza Treatment: A Randomized, Double-Blind, Double-Dummy, Multicenter, Noninferiority Clinical Trial. *Curr Therap Res* 2015, 77: 66-72
- Rininger JA, Kickner S, Chigurupati P, McLean A, Franck Z. Immunopharmacological activity of *Echinacea* preparations following simulated digestion on murine macrophages and human peripheral blood mononuclear cells. *J Leukoc Biol* 2000, 68: 503-10
- Rizzello CG, Coda R, Macías DS, Pinto D, Marzani B, Filannino P et al. Lactic acid fermentation as a tool to enhance the functional features of *Echinacea* spp. *Microb Cell Fact* 2013, 12: 44
- Romiti N, Pellati F, Nieri P, Benvenuti S, Adinolfi B, Chieli E. P-Glycoprotein inhibitory activity of lipophilic constituents of *Echinacea pallida* roots in a human proximal tubular cell line. *Planta Med* 2008, 74: 264-266
- Schoop R, Klein P, Suter A, Johnston SL. Echinacea in the prevention of induced rhinovirus colds: a meta-analysis. *Clin Ther* 2006, 28: 174-183
- Senchina DS, Flagel LE, Wendel JF, Kohut ML. Phenetic comparison of seven *Echinacea* species based on immunomodulatory characteristics. *Econ Bot* 2008, 60: 205-211
- Senchina DS, McCann DA, Asp JM, Johnson JA, Cunnick JE, Kaiser MS, et al. Changes in immunomodulatory properties of *Echinacea* spp. root infusions and tinctures stored at 4 degrees C for four days. *Clin Chim Acta* 2005, 355: 67-82
- Senchina DS, Wu L, Flinn GN, Konopka del N, McCoy JA, Widrlechner MP, et al. Year-and-a-half old, dried *Echinacea* roots retain cytokine-modulating capabilities in an in vitro human older adult model of influenza vaccination. *Planta Med* 2006, 72: 1207-1215
- Shah SA, Sander S, White CM, Rinaldi M, Coleman CI. Evaluation of *Echinacea* for the prevention and treatment of the common cold: a meta-analysis. *Lancet Infect Dis* 2007, 7: 473-480
- Sharma M, Arnason JT, Burt A, Hudson JB. *Echinacea* extracts modulate the pattern of chemokine and cytokine secretion in rhinovirus-infected and uninfected epithelial cells. *Phytother Res* 2006, 20: 147-152
- Shin DM, Choi KM, Lee YS, Kim W, Shin KO, Oh S, et al. *Echinacea purpurea* root extract enhances the adipocyte differentiation of 3T3-L1 cells. *Arch Pharm Res* 2014, 37(6): 803-812

Skaudickas D, Kondrotas A, Baltrusaitis K. The effect of Echinacea purpurea extract on sexual glands of male rats. *Medicina (Kaunas)* 2004, 40: 1211-1218

Skaudickas D, Kondrotas AJ, Baltrusaitis K, Vaitiekaitis G. Effect of Echinacea (Echinacea purpurea (L.) Moench) preparations on experimental prostate gland. *Medicina (Kaunas)* 2003, 39: 761-766

Skwarek T, Tynecka Z, Glowniak K, Lutostanska E. *Echinacea* L. – Inducer of interferons. *Herba Polonica* 1996, 42: 110-117

Sloley BD, Urichuk LJ, Tywin C, Coutts RT, Pang PK, Shan JJ. Comparison of chemical components and antioxidants capacity of different Echinacea species. *J Pharm Pharmacol* 2001, 53: 849-857

South EH, Exon JH. Multiple immune functions in rats fed *Echinacea* extracts. *Immunopharmacol Immunotoxicol* 2001, 23: 411-421

Speroni E, Govoni P, Guzzardi S, Renzulli C, Guerra MC. Anti-inflammatory and cicatrizing activity of *Echinacea pallida* Nutt. root extract. *J Ethnopharmacol* 2002, 79: 265-272

Stimpel M, Proksch A, Wagner H, Lohmann-Matthes ML. Macrophage activation and induction of macrophage cytotoxicity by purified polysaccharide fractions from the plant *Echinacea purpurea*. *Infect Immun* 1984, 46: 845-849

Sun LZ, Currier NL, Miller SC. The American coneflower: a prophylactic role involving nonspecific immunity. *J Altern Complement Med* 1999, 5: 437-446

Thygesen L, Thulin J, Mortensen A, Skibsted LH, Molgaard P. Antioxidant activity of cichoric acid and alkamides from *Echinacea purpurea*, alone and in combination. *Food Chem* 2007, 101: 74-81

Tiralongo E, Lea RA, Wee SS, Hanna MM, Griffiths LR. Randomised, double blind, placebo-controlled trial of echinacea supplementation in air travellers. *E- Based Complem Altern Med* 2012, 417267

Todd DA, Gullede TV, Britton ER, Oberhofer M, Leyte-Lugo M, Moody AN et al. Ethanolic *Echinacea purpurea* Extracts Contain a Mixture of Cytokine-Suppressive and Cytokine-Inducing Compounds, Including Some That Originate from Endophytic Bacteria. *PLoS One* 2015, 10(5): e0124276

Uluişik D, Keskin E (2012). Effects of ginseng and echinacea on cytokine mRNA expression in rats. *ScieWorld J* 2012: 942025

Wacker A, Hilbig W. Virushemmung mit *Echinacea purpurea*. *Planta Med* 1978, 33: 89-102

Wagner H, Breu W, Willer F, Wierer M, Remiger P, Schwenker G. In vitro inhibition of arachidonate metabolism by some alkamides and prenylated phenols. *Planta Med* 1989, 55: 566-567

Wagner H, Jurcic K. Immunological studies of Revitonil, a phytopharmaceutical containing *Echinacea purpurea* and *Glycyrrhiza glabra* root extract. *Phytomedicine* 2002, 9: 390-397

Wagner H, Proksch A, Riess-Maurer I, Vollmar A, Odenthal S, Stuppner H, et al. Immunostimulating action of polysaccharides (heteroglycans) from higher plants. *Arzneimittelforsch* 1984, 35: 1069-1075

Westendorf J. Carito® – in-vitro Untersuchungen zum Nachweiss spasmolytischer und kontraktiler Einflüsse. *Therapiewoche* 1982, 32: 6291-6297

WHO. Radix echinaceae. In: WHO monographs on selected medicinal plants. Vol. 1. WHO, Geneva 1999, Switzerland: 128-135

Wolters Kluwer Health. *Echinacea*. In: The Review of Natural Products. Wolters Kluwer Health, St. Louis 2004, MO, USA

Wu L, Rowe EW, Jeftinija K, Jeftinija S, Rizshsky L, Nikolau BJ *et al.* *Echinacea*-induced cytosolic Ca²⁺ elevation in HEK293. *BMC Complem Altern Med* 2010, 10: 72

Zhai Z, Haney D, Wu L, Solco A, Murphy PA, Wurtele ES, Kohut ML, Cunnick JE. Alcohol extracts of *Echinacea* inhibit production of nitric oxide and tumor necrosis factor-alpha by macrophages in vitro. *Food Agric Immunol* 2007b, 18: 221-236

Zhai Z, Liu Y, Wu L, Senchina DS, Wurtele ES, Murphy PA, Kohut ML, Cunnick JE. Enhancement of innate and adaptive immune functions by multiple *Echinacea* species. *J Med Food* 2007a, 10: 423-434

Zoutewelle G, Van Wijk R. Effects of *Echinacea purpurea* extracts on fibroblast populated collagen lattice contraction. *Phytother Res* 1990, 4: 77-81