

27 March 2012
EMA/HMPC/143182/2010
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Lavandula angustifolia* Miller, aetheroleum and *Lavandula angustifolia* Miller, flos

Final

The Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Akhondzadeh S, Kashani I, Fotouhi A, Jarvandi S, Mohaseri M, Moin M, et al. Comparison of Lavandula angustifolia Mill. tincture and imipramine in the treatment of mild to moderate depression: a double-blind, randomized trial. *Prog Neuropsychopharmacol Biol Psychiatr* 2003; 27: 123-127

Akutsu H, Kikusui T, Takeuchi Y, Sano K, Hatanaka A, Mori Y. Alleviating effects of plant-derived fragrances on stress-induced hyperthermia in rats. *Physiol Behav* 2002; 75: 355-360

Atassanova-Shopova S, Roussinov KS. On certain central neurotropic effects of lavender essential oil. Bulletin of the Institute of Physiology (Bulgarian Academy of Sciences) 1970; 13: 69-77

Atsumi T, Tonosaki K. Smelling lavender and rosemary increases free radical scavenging activity and decreases cortisol level in saliva. *Psychiatry Research* 2007; 150(1): 89-96

Austin LM, Alford C. The sedating effects of lavender oil and the stimulating effects of jasmine oil on sleep. *J Psychopharmacol Oxford* 1997; 11(3): A55, abstract 2171997

Bickers D, Calow P, Greim h, Hanifin JM, Rogers AE, Saurat JH, Sipes IG, Smith RL, Tagami H. A toxicological and dermatological assessment of linalool and related esters when used as fragrance ingredients. *Food and Chemical Toxicology* 2003; 41: 919-942

Blumenthal M, Goldberg A, Brinckmann J, Foster S. *Herbal Medicine. Expanded Commission E Monographs.* American Botanical Council, Austin Texas 2000; 226-229

Bradley BF, Brown SL, Chu S, Lea RW. Effects of orally administered lavender essential oil on responses to anxiety-provoked film clips. *Hum Psychopharmacol* 2009; 24(4): 319-330

Bradley BF, Starkey NJ, Brown SL, Lea RW. Anxiolytic effects of *Lavandula angustifolia* odour on the Mongolian gerbil elevated plus maze. *J Ethnopharmacology* 2007; 111(3): 517-525

British Herbal Pharmacopoeia, 2nd ed. Part 2. Cowling, British Herbal Medicine Association 1979



- British Herbal Pharmacopoeia. British Herbal Medicine Association, Bournemouth 1983; 128-129
- Bruneton J. Pharmacognosie: Phytochimie Plantes Médicinales. Ed. Tec & Doc, Paris 1999; 529-530
- Buchbauer G, Jirovetz L, Jäger W, Dietrich H, Plank C, Karamat E. Aromatherapy: evidence for sedative effects of the essential oil of lavender after inhalation. *Z Naturforsch* 1991; 46c: 1067-1072
- Buchbauer G, Jirovetz L, Jäger W, Plank C, Dietrich H. Fragrance compounds and essential oils with sedative effects upon inhalation. *J Pharm Sci* 1993a; 82: 660-664
- Buchbauer G, Jäger W, Jirovetz L, Limberger J, Dietrich H. Therapeutic properties of essential oils and fragrances. In: Teranishi R, Buttery RG, Sugisama H, editors. Bioactive Volatile Compounds from Plants. AC Symposium Series 525. Washington DC: American Chemical Society, 1993b
- Büyükokuroglu ME, Gepdiremen A, Hacimüftüoglu A, Oktay M. The effects of aqueous extract of *Lavandula angustifolia* flowers in glutamate-induced neurotoxicity of cerebellar granular cell culture of rat pups. *J Ethnopharmacol* 2003; 84: 91-94
- Coulson IH, Khan ASA. Facial 'pillows' dermatitis due to lavender oil allergy. *Contact Dermatitis* 1999; 41: 111
- Dale A, Cornwell S. The role of lavender oil in relieving perineal discomfort following childbirth: a blind randomized clinical trial. *J Advanced Nurs* 1994; 19: 89-96
- Delaveau P, Guillemain J, Narcisse G, Rousseau A. The Neurodepressive Properties of Essential Oil of Lavender. *C R Seances Soc Biol Ses Fil* 1989; 183(4): 342-348
- Diego MA, Jones NA, Filed T, Hernandez-Reif M, Schanberg S, Kuhn C, et al. Aromatherapy positively affects mood, EEG patterns of alertness and math computations. *Int J Neurosci* 1998; 96: 217-224
- Dodonaeus R. Herbarius oft Cruydt-Boeck. Plantijnsche Druckerije, Antwerpen 1608; 470-473
- Duan X, Tashiro M, Wu D, Yambe T, et al. Autonomic nervous function and localization of cerebral activity during lavender aromatic immersion. *Technol Health Care* 2007; 15(2): 69-78
- Dunn C, Sleep J, Collett D. Sensing an improvement: an experimental study to evaluate the use of aromatherapy, massage and periods of rest in an intensive care unit. *J Advanced Nursing* 1995; 21: 34-40
- Eder E, Neudecker T, Lutz D, Henschler D. Mutagenic potential of allyl and allylic compounds. Structure activity relationship asdetermined by alkylating and direct in vitro mutagenic properties. *Biochem Pharmacol* 1980; 29: 993-998
- Eder E, Henschler D, Neudecker T. Mutagenic properties of allylic and alpha,beta-unsaturated compounds: consideration of alkylating mechanisms. *Xenobiotica* 1982a; 12: 831-848
- Eder E, Neudecker T, Lutz D, Henschler D. Correlation of alkylating and mutagenic activities of allyl and allylic compounds: standard alkylation test vs. kinetic investigation. *Chem-Biol Interact* 1982b; 38: 303-315
- Elhajili M, Baddouri K, Elkabbaj S, Meiouat F, Settaf A. Effet diurétique de l'infusion de fleurs de *Lavandula officinalis*. *Reprod Nutr Dev* 2001; 41: 393-399
- ESCOP Monographs 2nd ed. *Lavandulae Flos, Lavandulae Aetheroleum*. European Scientific Cooperative on Phytotherapy, editor. Supplement 2009. Georg Thieme Verlag, Stuttgart 2009; 147-156
- European Pharmacopoeia 6.0 (2008a) Lavender floweroil. Council of Europe. 1/2008: 1534
- European Pharmacopoeia 6.0 (2008b) Lavender oil. Council of Europe. 1/2008: 1338

Evrandi MG, Battinelli L, Daniele C, Mastrangelo S, Bolle P, Mazzanti G. The antimutagenic activity of *Lavandula angustifolia* (lavender) essential oil in the bacterial reverse assay. *Food Chem. Toxicology* 2005; 43: 1381-1387

Fernandez M, Hernandez Reif M, Field T, Diego M, Sanders C, Roca A. EEG during lavender and rosemary exposure in infants of depressed and non-depressed mothers. *Infant Behavior & Development* 2004; 27(1): 91-100

Field T, Field T, Cullen C, Largie S, Diego M, Schanberg S, Kuhn C. Lavender bath oil reduces stress and crying and enhances sleep in very young infants. *Early Human Development* 2008; 84: 399-401

Fischer G. Heilkräuter und Arzneipflanzen. Reichenberg, Gudetendeutscher Verlag Franz Kraus pp. 1939 124-125Fujii M, Hatakeyama R, Fukukoa Y, et al. Lavender aromatherapy for behaviour and psychological symptoms in dementia patients. *Geriatr Gerontol Int* 2008; 8: 136-138

Gedney JJ, Glover TL, Fillingim RB. Sensory and affective pain discrimination after inhalation of essential oils. *Psychosomatic Med* 2004; 66: 599-606

Ghelardini C, Galeotti N, Salvatore G, Mazzanti G. Local anaesthetic activity of essential oil of *Lavandula angustifolia*. *Planta Medica* 1999; 65: 700-703

Guillemain J, Rousseau A, Delaveau P. Effets neurodépresseurs de l'huile essentielle de *Lavandula angustifolia* Mill. *Ann Pharmaceutiques Françaises* 1989; 47(6): 337-343

Hänsel R, Keller K, Rimpler H, Schneider G. Hager's Handbuch der Pharmazeutischen Praxis. Springer Verlag, Berlin 1993; 630-644

Hardy M, Kirk Smith MD, Stretch DD. Replacement of drug treatment for insomnia by ambient odor. *Lancet* 1995; 346: 701

Hausen BM, Vieluf IK. *Lavandula angustifolia* Miller subsp. *angustifolia*. Echte Lavendel. In: Allergiepflanzen – Handbuch und Atlas. Kontaktallergene, Allergische Frühreaktionen. 2nd ed. Nikol. Verlagsgesellschaft, Hamburg 1997; 175-177

Hawrelak JA, Cattley T, Myers SP. Essential oils in the treatment of intestinal dysbiosis: a preliminary in vitro study. *Altern Med Rev* 2009; 14(4): 380-384

Henley DV, Lipson N, Korach KS, Bloch CA. Prepubertal gynecomastia linked to lavender and tea tree oils. *N Engl J Med* 2007; 356(5): 479-485

Holmes C, Hopkins V, Hensford C, MacLaughlin V, Wilkinson D, Rosenvinge H. Lavender oil as a treatment for agitated behaviour in severe dementia: a placebo controlled study. *Int J Geriatric Psychiatry* 2002; 17: 305-308

Hudson R. The value of lavender for rest and activity in the elderly patient. *Complement Ther Med* 1996; 4(1): 52-57

Hwang JH. The effects of the inhalation method using essential oils on blood pressure and stress responses of clients with essential hypertension. *Taehan Kanho Hakhoe Chi* 2006; 36(7): 1123-1134

Ishidate M, Sofuni T, Yoshikawa K, Hayashi M, Nohmi T, Sawada M, Matsuoka A. Primary mutagenicity screening of food additives currently used in Japan. *Food Chem Toxicol* 1984; 22: 623-636

Jäger W, Buchbauer G, Jirovetz L, Fritzer M. Percutaneous absorption of lavender oil from a massage oil. *J Soc Cosmet Chem* 1992; 43: 49-54

Jirovetz L, Buchbauer G, Jäger W, Raverdinov V, Nikiforov A. Determination of lavender oil fragrance compounds in blood samples. *Fresenius J Anal Chem* 1990; 338: 922-923

Kane FMA, Brodie EE, Coull A, Coyne I, Howd A, Milne A, et al. The analgesic effect of odour and music upon dressing change. *Br J Nurs* 2004; 13(19): 4-12

Kawakami K, Takai-Kawakami K, Okazaki Y, Kurihara H, Shimizu Y, Yanaihara T. The effect of odors on human newborn infants under stress. *Infant Behavior & Development* 1997; 20(4): 531-535

Kim HM, Cho SH. Lavender oil inhibits immediate-type allergic reaction in mice and rats. *J Pharm Pharmacol* 1999; 55: 221-226

Kim JT, Ren CJ, Fielding GA, Pitti A, Kasumi T, Wajda M, et al. Treatment with lavender aromatherapy in the post-anesthesia care unit reduces opioid requirements of morbidly obese patients undergoing laparoscopic adjustable gastric banding. *Obes Surg* 2007; 17: 920-925

Kim JT, Wajda M, Cuff G, Serota D, Schlame M, Axelrod DM, Guth AA, Bekker AY. Evaluation of aromatherapy in treating postoperative pain: pilot study. *Pain Practice: the Official Journal of World Institute of Pain* 2006; 6(4): 273-277

Koto R, Imamura M, Watanabe C, Obayashi S, Shiraishi Mi, Sasaki Y, Azuma H. Linalyl acetate as a major ingredient of lavender essential oil relaxes the rabbit vascular smooth muscle through dephosphorylation of myosin light chain. *J Cardiovasc Pharmacol* 2006; 48(1): 850-856

Kritsidima M, Newton T, Asimakopoulou K. The effects of lavender scent on dental patient anxiety levels: a cluster randomized-controlled trial. *Community Dent Oral Epidemiol* 2010; 38(1): 83-87

Kroeber L. Das neutzeitliche Kräuterbuch. Hippokrates Verlag G.M.B.H. Stuttgart-Leipzig. 1935 pp. 114-117

Kubo I, Muroi H, Kubo A. Antibacterial activity of long-chain alcohols against *Streptococcus mutans*. *J Agric Food Chem* 1993; 41: 2447-2450

Leclerc H. Précis de phytothérapie. Masson Eds, Paris 1966 ; 204-205

Lee IS, Lee GJ. Effects of lavender aromatherapy on insomnia and depression in women college students. *Taehan Kanho Hakhoe Chi* 2006; 36(1): 136-43

Lis-Balchin M, Hart S. A preliminary study of the effect of essential oils on skeletal and smooth muscle in vitro. *J Ethnopharmacol* 1997; 58: 183-187

Lis-Balchin M, Hart S. Studies on the mode of action of the essential oil of lavender (*Lavandula angustifolia* P. Miller). *Phytotherapy Research* 1999; 13: 540-542

Louis M, Kowalski SD. Use of aromatherapy with hospice patients to decrease pain, anxiety, and depression and to promote an increasing sense of well-being. *Am J Hospice Palliative Care* 2002; 19: 381-386

Madaus G. Lehrbuch der biologischen Heilmittel Band II. Georg Thieme Verlag, Leipzig, 1938 pp. 1721-1725

Maddocks-Jennings W. Critical incident: idiosyncratic allergic reactions to essential oils. *Complement Ther Nurs Midwifery* 2004; 10: 58-60

Masago R, Matsuda T, Kikuchi Y, Miyazaki Y, Iwanaga K, Harada H, Katsuura T. Effects of inhalation of essential oils on EEG activity and sensory evaluation. *J Physiol Antropol* 2000; 19: 35-42

Morris N. The effects of lavender (*Lavendula angustifolium*) baths on psychological well-being: two exploratory randomised control trials. *Complementary Therapies in Medicine* 2002; 10(4): 223-228

Moss M, Cook J, Wesnes K, Duckett P. Aromas of rosemary and lavender essential oils differentially affect cognition and mood in healthy adults. *Int J Neurosci* 2003; 113: 15-38

Motomura N, Sakurai A, Yotsuya Y. Reduction of mental stress with lavender odorant. *Percept Motor Skills* 2001; 93: 713-718

Muzzarelli L, Force M, Sebold M. Aromatherapy and reducing preprocedural anxiety: A controlled prospective study. *Gastroenterol Nurs* 2006; 29(6): 466-471

Nord D, Belew J. Effectiveness of the essential oils lavender and ginger in promotion children's comfort in a perianesthesia setting. *J Perianesth Nurs* 2009; 24(5): 307-312

Opdyke DLJ. Monographs on fragrance raw materials: Lavender oil. *Food Cosmet Toxicol* 1976; 14: 451

Pabst G. Koehler's Medizinal-Pflanzen. Hugo Bermüller Verlag, Berlin 1888 pp. 60-62

Parke DV, Rahman KMQ, Walker R. Effect of linalool on hepatic drug-metabolising enzymes in the rat. *Biochem Soc Trans* 1974; 2: 615-618

Peana AT, D'Aquila PS, Panin F, Serra G, Pippia P, Moretti MDL. Anti-inflammatory activity of linalool and linalyl acetate constituents of essential oils. *Phytomedicine* 2002; 9: 721-726

Peana AT, D'Aquila PS, Chessa ML, Moretti MDL, Serra G, Pippia P. (-)-Linalool produces antinociception in two experimental models of pain. *Eur J Pharmacology* 2003; 460: 37-41

Peana AT, De Montis MG, Sechi S, Sircana G, D'Aquila PS, Pippia P. Effects of (-)-linalool in the acute hyperalgesia induced by carrageenan, L-glutamate and prostaglandin E. *Eur J Pharmacol* 2004a; 497: 279-284

Peana AT, De Montis MG, Nieddu E, Spano MT, D'Aquila PS, Pippia P. Profile of spinal and supraspinal antinociception of (-)-linalool. *Eur J Pharmacol* 2004b; 485: 165-174

Pelikan W. Heilpflanzenkunde. Der Mensch und die Heilpflanzen Band I. Philosophisch-Antroposophischer Verlag am Goetheanum. Dornach (Schweiz) 2.Auflage 1958 pp. 51-52

Pulla Reddy AC, Lokesh BR. Studies on anti-inflammatory activity of spice principles and dietary n-3-polyunsaturated fatty acids on carrageenan-induced inflammation in rats. *Ann Nutr Metabol* 1994; 38: 349-358

Rademaker M. Allergic contact dermatitis from lavender fragrance in Difflam® gel. *Contact Derm* 1994; 31: 58

Rahimifard N, Hajimehdipoor H, Pirouz B, Bagheri F, Bagheri O, Mirdamadi KS (2010) The mutagenic and antimutagenic activity of *Lavandula angustifolia* and *Elettaria cardamomum* essential oils in the bacterial reverse mutation assay. *J Medicinal Plants* 2010; 35 (9): 139-142

Rho KH, Han SH, Kim KS, Lee MS. Effects of aromatherapy massage on anxiety and self-esteem in Korean elderly women: a pilot study. *The International Journal of Neuroscience* 2006; 116(12): 1447-1455

Roffey SJ, Walker R, Gibson GG. Hepatic peroxisomal and microsomal enzyme induction by citral and linalool in rats. *Food Chem Toxic* 1990; 28: 403-408

Romine IJ, Bush AM, Geist CR. Lavender aromatherapy in recovery from exercise. *Percept Motor Skills* 1999; 88: 756-758

Saeki Y. The effects of footbath with or without the essential oil of lavender on the autonomic nervous system: a randomized trial. *Complement Ther Med* 2000; 8: 2-7

Sanders C, Diego M, Fernandez M, Field T, Hernandez-Reil M, Roca A. EEG asymmetry responses to lavender and rosemary aromas in adults and infants. *Int J Neurosci* 2002; 112: 305-320

Schulz H, Jobert M, Huebner WD. The quantitative EEG as a screening instrument to identify sedative effects of single doses of plant extracts in comparison with diazepam. *Phytomedicine* 1998; 5(6): 449-458

Shaw D, Annett JM, Doherty B, Leslie JC. Anxiolytic affects of lavender oil inhalation on open-field behavior in rats. *Phytomedicine* 2007; 14(9): 613-620

Shimizu K, Gyokusen M, Kitamura S, Kawabe T, Kozaki T, Ishibashi K, et al. Essential oil of lavender inhibited decreased attention during a long-term task in humans. *Biosci Biotechnol Biochem* 2008; 72: 1944-1947

Snow AL. A controlled trial of aromatherapy for agitation in nursing home patients with dementia. *Journal of Alternative and Complementary Medicine* 2004; 10(3): 431-437

Sweetman S. Martindale the Complete Drug Reference. 36th ed. Pharmaceutical Press, London 2009; 2331

Tašev T, Taléva P, Balabanova V. Effet neuro-psychique des huiles essentielles bulgares de rose, de lavande et de géranium. *Folia Medica* 1969; XI(V): 307-316

Umezawa T, Nagano K, Ito H, Kosakai K, Sakaniwa M, Morita M. Anticonflict effects of lavender oil and identification of its active constituents. *Pharmacology Biochemistry and Behavior* 2006; 85(4): 713-721

Valnet J. L'Aromathérapie. Maloine S.A. Editeur (since 1964), Vigot, Paris 2001; 272-278

von Skramlik E. Über die Giftigkeit und Verträglichkeit von ätherischen Ölen. *Pharmazie* 1959; 14: 435-445

Weiss RF, Fintelmann V. Lehrbuch der Phytotherapie. Hippocrates Verlag, Stuttgart 1999; 298

Williams TI. Evaluating effects of aromatherapy massage on sleep in children with autism: a pilot study. *Evidence-based Complementary and Alternative Medicine* 2006; 3(3): 373-377

Woelk H, Schlafke S. A multi-center, double-blind, randomized study of the lavender oil preparation Silexan in comparison to lorazepam for generalized anxiety. *Phytomedicine* 2010; 17(2): 94-99

Yamada K, Nimaki Y, Sashida Y. Anticonvulsive effects of inhaling lavender oil vapour. *Biol Pharm Bulletin* 1994; 17: 359-360

Yasukawa K, Takido M, Takeuchi M, Nakagawa S. Effect of chemical constituents from plants on 12-O-tetradecanoylphorbol-13-acetate induced inflammation in mice. *Chem Pharm Bull* 1989; 37: 1071-1073

Yip YB, Tse SHM. The effectiveness of relaxation acupoint stimulation and acupressure with aromatic lavender essential oil for non-specific low back pain in Hong Kong: a randomized controlled trial. *Complement Ther Med* 2004; 12: 28-37

Yip YB, Tse SHM. An experimental study on the effectiveness of acupressure with aromatic lavender essential oil for sub-acute, non-specific neck pain in Hong Kong. *Complementary Therapies in Clinical Practice* 2006; 12(1): 18-26