

7 July 2015
EMA/HMPC/55810/2011
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Matricaria recutita L.*, flos and *Matricaria recutita L.*, aetheroleum Final

The European Medicine Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Achterrath-Tuckermann U, Kunde R, Flaskamp E, Isaac O, Thiemer K. [Pharmacological investigations with compounds of chamomile. V. Investigations on the spasmolytic effect of compounds of chamomile and Kamillosan on the isolated guinea pig ileum.] *Planta Med* 1980, 39(1):38-50 [German]

Aertgeerts J. Erfahrungen mit Kamillosan, einem standardisierten Kamilleextrakt in der dermatologischen Praxis. *Ärztl Kosmetol* 1984, 14:502-504

Aertgeerts P, Albring M, Klaschka F, Nasemann T, Patzelt-Wenzler R, Rauhut K, et al. Vergleichende Prüfung von Kamillosan®-Creme gegenüber steroidalen (0,25% Hydrocortison, 0,75% Fluocortinbutylester) und nichtsteroidalen (5% Bufexamac) Externa in der Erhaltungstherapie von Ekzemerkrankungen. *Z Hautkr* 1985, 60(3):270-277

Aggag ME, Yousef RT. Study of antimicrobial activity of chamomile oil. *Planta Med* 1972, 22(2):140-144

Albring M, Albrecht H, Alcorn G, Lücker PW. The measuring of the anti-inflammatory effect of a compound on the skin of volunteers. *Methods Find Exp Clin Pharmacol* 1983, 5(8):575-577

Al-Hindawi MK, Al-Deen IH, Nabi MH, Ismail MA. Anti-inflammatory activity of some Iraqi plants using intact rats. *J Ethnopharmacol* 1989, 26(2):163-168

Al-Ismail KM, Talal A. A Study of the effect of water and alcohol extracts of some plants as antioxidants and antimicrobial on long-term storage of anhydrous butter fat. *Dirasat Agric Sci* 2003, 30(3):330-337

Amsterdam JD, Li Y, Soeller I, Rockwell K, Mao JJ, Shults J. A randomized, double-blind, placebo-controlled trial of oral *Matricaria recutita* (Chamomile) extract therapy for generalized anxiety disorder. *J Clin Psychopharmacol* 2009, 29(4):378-382



Anter J, Romero-Jiménez M, Fernández-Bedmar Z, Villatoro-Pulido M, Analla M, Alonso-Moraga A, et al. Antigenotoxicity, cytotoxicity, and apoptosis induction by apigenin, bisabolol, and protocatechuic acid. *J Med Food* 2011, 14(3):276-283

Arzneimittelverzeichnis der DDR 1983, VEB Verlag Volk und Gesundheit Berlin 1983

Avallone R, Zanolli P, Corsi L, Cannazza G, Baraldi M. Benzodiazepine-like compounds and GABA in flower heads of *Matricaria chamomilla*. *Phytother Res* 1996, 10:177-179

Avallone R, Zanolli P, Puia G, Kleinschnitz M, Schreier P, Baraldi M. Pharmacological profile of apigenin, a flavonoid isolated from *Matricaria chamomilla*. *Biochem Pharmacol* 2000, 59(11):1387-1394

Barene I, Daberte I, Zvirgzdina L, Iriste V. The complex technology on products of German Chamomile. *Medicina (Kaunas)* 2003, 39(Suppl 2):127-131

Barnes J, Anderson L, Phillipson JD. *Herbal Medicines*. 3rd ed. Pharmaceutical Press, London Chicago 2007

Becker B, Kuhn U, Hardewig-Budny B. Double-blind, randomized evaluation of clinical efficacy and tolerability of an apple pectin-chamomile extract in children with unspecific diarrhea.

Arzneimittelforschung 2006, 56(6):387-393

Benedum J, Loew D, Schilcher H. *Arzneipflanzen in der Traditionellen Medizin*. Verlag Kooperation Phytopharmaka GbR, Bonn 2006

Benner MH, Lee HJ. Anaphylactic reaction to chamomile tea. *J Allergy Clin Immunol* 1973, 52(5):307-308

Bishop JL, Northstone K, Green JR, Thompson EA. The use of Complementary and Alternative Medicine in pregnancy: Data from the Avon Longitudinal Study of Parents and Children (ALSPAC). *Complement Ther Med* 2011, 19(6):303-310

Blaschek W, Hilgenfeldt U, Holzgrabe U, Reichling J, Ruth P, Schulz V, editors. *HagerROM 2011: Hagers Enzyklopädie der Arzneistoffe und Drogen*. Springer Medizin Verlag, Berlin Heidelberg 2011

British Herbal Pharmacopoeia. *Matricaria*. 3rd ed. British Herbal Medicine Association, Bournemouth 1983

British Pharmacopoeia. *Matricaria Flowers*. 2008, 1376-1377

Budzinski JW, Foster BC, Vandenhoeck S, Arnason JT. An *in vitro* evaluation of human cytochrome 450 3A4 inhibition by selected commercial herbal extracts and tinctures. *Phytomedicine* 2000, 7(4):273-282

Capasso R, Savino F, Capasso F. Effects of the herbal formulation ColiMil on upper gastrointestinal transit in mice *in vivo*. *Phytother Res* 2007, 21(10):999-1001

Carl W, Emrich LS. Management of oral mucositis during local radiation and systemic chemotherapy. A study of 98 patients. *J Prosthet Dent* 1991, 66(3):361-369

Carle R, Gomaa K. Technologische Einflüsse auf die Qualität von Kamillenblüten und Kamillenöl. *Pharm Ztg*, 1992a, 2(5):71-77

Cemek M, Yilmaz E, Büyükkokuoğlu ME. Protective effect of *Matricaria chamomilla* on ethanol-induced acute gastric mucosal injury in rats. *Pharm Biol* 2010, 48(7):757-763

Cuzzolin L, Francini-Pesenti F, Verlato G, Joppi M, Baldelli P, Benoni G. Use of herbal products among 392 Italian pregnant women: focus on pregnancy outcome. *Pharmacoepidemiol Drug Saf* 2010, 19(11):1151-1158

De La Motte S, Böse-O'Reilly S, Heinisch M, Harrison F. [Double-blind comparison of a preparation of pectin/chamomile extract and placebo in children with diarrhea.] *Arzneimittelforschung* 1997, 47(11):1247-1249 [German]

De la Torre Morin F, Sánchez Machín I, García Robaina JC, Fernández-Caldas E, Sánchez Triviño M. Clinical cross-reactivity between *Artemisia vulgaris* and *Matricaria chamomilla* (Chamomile). *J Investig Allergol Clin Immunol* 2001, 11(2):118-122

Della Loggia RD, Traversa U, Scarcia V, Tubaro A. Depressive effects of *Chamomilla recutita* (L.) Rausch., tubular flowers, on central nervous system in mice. *Pharmacol Res Commun* 1982, 14(2):153-162

Della Loggia R, Carle R, Sosa S, Tubaro A. Evaluation of the anti-inflammatory activity of chamomile preparations. *Planta Med* 1990, 56(6):657-658

Di Carlo G, Autore G, Izzo AA, Maiolino P, Mascolo N, Viola P et al. Inhibition of intestinal motility and secretion by flavonoids in mice and rats: structure-activity relationships. *J Pharm Pharmacol* 1993, 45(12):1054-1059

Dorsch W, Loew D, Meyer E, Schilcher H. Empfehlungen zu Kinderdosierungen von monographierten Arzneidrogen und ihren Zubereitungen. Kooperation Phytopharmaka, Bonn 1993

ESCOP Monographs 2nd ed. Vol 2. *Matricariae flos – Matricaria flower*. European Scientific Cooperative on Phytotherapy, editor. Thieme, Stuttgart 2003, 312-319

European Pharmacopoeia 8th ed. *Matricaria flower - Matricariae flos*. Council of Europe. 2008:0404

European Pharmacopoeia 8th ed. *Matricaria liquid extract - Matricariae extractum fluidum*. Council of Europe. 2008:1544

European Pharmacopoeia 8th ed. *Matricaria oil - Matricariae aetheroleum*. Council of Europe. 2008:1836

Facchinetti F, Pedrielli G, Benoni G, Joppi M, Verlato G, Dante G, et al. Herbal supplements in pregnancy: unexpected results from a multicenter study. *Hum Reprod* 2012, 27(11):3161-3167

Fidler P, Loprinzi CL, O'Fallon JR, Leitch JM, Lee JK, Hayes DL, et al. Prospective evaluation of a chamomile mouthwash for prevention of 5-FU-induced oral mucositis. *Cancer* 1996, 77(3):522-525

Forster HB, Niklas H, Lutz S. Antispasmodic effects of some medicinal plants. *Planta Med* 1980, 40(4):309-319

Förster CF. Proktologische Operationen. Kamille fördert die Heilung. *Ärztliche Praxis* 1987, 74:2122-2123

Forster DA, Denning A, Wills G, Bolger M, McCarthy E. Herbal Medicine use during pregnancy in a group of Australian women. *BMC Pregnancy Childbirth* 2006, 6:21-30

Ganzena M, Schneider P, Stuppner H. Inhibitory effects of the essential oil of chamomile (*Matricaria recutita* L.) and its major constituents on human cytochrome P450 enzymes. *Life Sci* 2006, 78(8):856-861

Gerritsen ME, Carley WW, Ranges GE, Shen CP, Phan SA, Ligon GF, et al. Flavonoids inhibit cytokine-induced endothelial cell adhesion protein gene expression. *Am J Pathol* 1995, 147(2):278-292

Gomes-Carneiro MR, Dias DM, De-Oliveira AC, Paumgartten FJ. Evaluation of mutagenic and antimutagenic activities of alpha-bisabolol in the Salmonella/microsome assay. *Mutat Res* 2005, 585(1-2):105-112

Hausen BM. A 6-year experience with compositae mix. *Am J Contact Dermat* 1996, 7(2):94-99

Hempel B, Hirschelmann R. Kamille-Entzündungshemmende Wirkung von Inhaltsstoffen und Zubereitungen in vivo. *Dtsch Apoth Ztg* 1998, 138(44):43-48

Hernández-Ceruelos A, Madrigal-Bujaidar E, de la Cruz C. Inhibitory effect of chamomile essential oil on the sister chromatid exchanges induced by daunorubicin and methyl methanesulfonate in mouse bone marrow. *Toxicol Lett* 2002, 135(1-2):103-110

Holst L, Wright D, Haavik S, Nordeng H. Safety and efficacy of herbal remedies in obstetrics – review and clinical implications. *Midwifery* 2011, 27(1):80-86

Jablonska S, Rudzki E. Kamillosan® Konzentrat – ein nicht allergisierender Extrakt aus Kamille. *Z Hautkr* 1996, 71(7):542-546

Jarrahi M. An experimental study of the effects of *Matricaria chamomilla* extract on cutaneous burn wound healing in albino rats. *Nat Prod Res* 2008, 22(5):422-427

Jensen-Jarolim E, Reider N, Fritsch R, Breiteneder H. Fatal outcome of anaphylaxis to camomile-containing enema during labour: A case study. *J Allerg Clin Immunol* 1998, 102:1041-1042

Jeschke E, Ostermann T, Lüke C, Tabali M, Kröz M, Bockelbrink A, et al. Remedies containing Asteraceae extracts: A Prospective Observational Study of Prescribing Patterns and Adverse Drug Reactions in German Primary Care. *Drug Saf* 2009, 32(8):691-706

Kalantari H, Dashtearjandi AA, Kalantar E. Genotoxicity study of *Hypiran* and *Chamomilla* herbal drugs determined by *in vivo* supervital micronucleus assay with mouse peripheral reticulocytes. *Acta Biol Hung* 2009, 60(2):177-183

Kerscher MJ. Influence of liposomal encapsulation on the activity of a herbal non-steroidal anti-inflammatory drug. In: Braun-Falco O, Korting HC, Maibach HI, editors. Liposome Dermatics: Griesbach Conference. Springer-Verlag, Berlin Heidelberg 1992, 329-337

Kobayashi Y, Takahashi R, Ogino F. Antipruritic effect of the single oral administration of German chamomile flower extract and its combined effect with antiallergic agents in ddY mice. *J Ethnopharmacol* 2005, 101(1-3):308-312

Kommission E . Aufbereitungsmonographie Matricariae flos. *Bundesanzeiger* (Nr. 228) vom 05/12/1984

Korting HC, Schäfer-Korting M, Hart H, Laux P, Schmid M. Anti-inflammatory activity of hamamelis distillate applied topically to the skin. Influence of vehicle and dose. *Eur J Clin Pharmacol* 1993, 44(4):315-318

Lauber B. Inhalationstherapie mit Kamillosan® liquidum. In: Klaschka F, Maiwald L, Patzelt-Wenzler R, editors. Wirkungsweise und Anwendungsformen der Kamille. Grosse Verlag, Berlin 1988, 99-105

Liang YC, Huang YT, Tsai SH, Lin-Shiau SY, Chen CF, Lin JK. Suppression of inducible cyclooxygenase and inducible nitric oxide synthase by apigenin and related flavonoids in mouse macrophages. *Carcinogenesis* 1999, 20(10):1945-1952

Lis-Balchin M, Deans SG, Eaglesham E. Relationship between bioactivity and chemical composition of commercial essential oils. *Flavour Frag J* 1998, 13(2):98-104

Lundh K, Hindsén M, Gruvberger B, Möller H, Svensson A, Bruze M. Contact allergy to herbal teas derived from Asteraceae plants. *Contact Dermatitis* 2006, 54(4):196-201

Lundh K, Gruvberger B, Möller H, Persson L, Hindsén M, Zimerson E, et al. Patch testing with thin-layer chromatograms of chamomile tea in patients allergic to sesquiterpene lactones. *Contact Dermatitis* 2007, 57(4):218-223

Maiche AG, Gröhn P, Mäki-Hokkonen H. Effect of chamomile cream and almond ointment on acute radiation skin reaction. *Acta Oncol* 1991, 30(3):395-396

Maliakal PP, Wanwimolruk S. Effect of herbal teas on hepatic drug metabolizing enzymes in rats. *J Pharm Pharmacol* 2001, 53(10):1323-1329

Mann C, Staba EJ. The chemistry, pharmacology and commercial formulations of Chamomile. In: Craker LE, Simon JE, editors. *Herbs, Spices and Medicinal Plants: Recent Advances in Botany, Horticulture and Pharmacology*. Vol 1. Oryx press, Phoenix 1986, 235-280

Marti MC. Die Analfissuren. Heilung durch einen einfachen operativen Eingriff. *Moderne Therapie* 1977, 7:94-103

Martins MD, Marques MM, Bussadori SK, Martins MA, Pavesi VC, Mesquita-Ferrari RA, et al. Comparative analysis between *Chamomilla recutita* and corticosteroids on wound healing. An *in vitro* and *in vivo* study. *Phytother Res* 2009, 23(2):274-278

Maschi O, Cero ED, Galli GV, Caruso D, Bosisio E, Dell'Agli M. Inhibition of human cAMP-phosphodiesterase as a mechanism of the spasmolytic effect of *Matricaria recutita* L. *J Agric Food Chem* 2008, 56(13):5015-5020

Matos FJA, Machado MIL, Alencar JW, Craveiro AA. Constituents of Brazilian Chamomile Oil. *J Essent Oil Res* 1993, 5(3):337-339

Medina JH, Viola H, Wolfman C, Marder M, Wasowski C, Calvo D, et al. Neuroactive flavonoids: new ligands for the Benzodiazepine receptors. *Phytomedicine* 1998, 5(3):235-243

Miller T, Wittstock U, Lindequist U, Teuscher E. Effects of some components of the essential oil of chamomile, *Chamomilla recutita*, on histamine release from rat mast cells. *Planta Med* 1996, 62(1):60-61

Mimica-Dukic N, Lukic V, Pavkov R, Gesic O. Study of chemical composition and microbiological contamination of chamomile tea. *Acta hortic* 1993, 333:137-141

Moussally K, Bérard A. Exposure to specific herbal products during pregnancy and the risk of low birth weight. *Alt Ther Health Med* 2012, 18(2):36-43

Moussally K, Oraichi D, Bérard A. Herbal products use during pregnancy: prevalence and predictors. *Pharmacoepidemiol Drug Saf* 2009, 18(6):454-461

Mulinacci N, Romani A, Pinelli P, Vincieri FF, Prucher D. Characterization of *Matricaria recutita* L. flower Extracts by HPLC-MS and HPLC-DAD Analysis. *Chromatografia* 2000, 51(5-6):301-307

Nasemann T. Kamillosan®-Behandlung in der Dermatologie. *Z Allgemeinmed* 1975, 51:1105-1106

Nasemann T, Menzel I. Kamillosan in der Dermatologie. In: Demling L, Nasemann T, Rosch W. Erfahrungstherapie – späte Rechtfertigung. Internationales Symposium Wien, 30-31 Mai. Verlag G. Braun, Karlsruhe 1975, 49-53

Nayak BS, Raju SS, Rao AV. Wound healing activity of *Matricaria recutita* L. extract. *J Wound Care* 2007, 16(7):298-302

- Nissen HP, Biltz H, Kreysel HW. Profilometrie, eine Methode zur Beurteilung der therapeutischen Wirksamkeit von Kamillosan®-Salbe. *Z Hautkr* 1988, 63:184-190
- Nordeng H, Havnen GC. Use of herbal drugs in pregnancy: a survey among 400 Norwegian women. *Pharmacoepidemiol Drug Saf* 2004, 13(6):371-380
- Nowack R, Nowack B. Herbal teas interfere with cyclosporin levels in renal transplant patients. *Nephrol Dial Transplant* 2005, 20(11):2554-2556
- Panés J, Gerritsen ME, Anderson DC, Miyasaka M, Granger DN. Apigenin inhibits tumor necrosis factor-induced intercellular adhesion molecule-1 upregulation *in vivo*. *Microcirculation* 1996, 3(3):279-286
- Patzelt-Wenzler R, Ponce-Pöschl E. Proof of efficacy of Kamillosan® cream in atopic eczema. *Eur J Med Res* 2000, 5(4):171-175
- Paulsen E, Chistensen LP, Andersen KE. Cosmetics and herbal remedies with Compositae plant extracts - are they tolerated by Compositae-allergic patients? *Contact Dermatitis* 2008, 58(1):15-23
- Paulsen E. Contact sensitization from Compositae-containing herbal remedies and cosmetics. *Contact Dermatitis* 2002, 47(4):189-198
- Pereira F, Santos R, Pereira A. Contact dermatitis from chamomile tea. *Contact Dermatitis* 1997, 36(6):307
- Peters H. Suchdiät und Neurodermitis Interdisziplinäres. In: Klaschka F, Maiwald L, Patzelt-Wenzler R, editors. Wirkungsweise und Anwendungsformen der Kamille. Grosse Verlag 1988, 45-51
- Pino JA, Bayat F, Marbot R, Aguero J. Essential oil of Chamomile (*Chamomilla recutita* (L.) Rausch) from Iran. *J Essent Oil Res* 2002, 14(6):407-408
- Pino JA, Marbot R, Agliero J, Fuentes V. Essential oil of Chamomile (*Chamomilla recutita* (L.) Rausch) from Cuba. *J Essent Oil-Bear Pl* 2003, 3:1-3
- Reider N, Sepp N, Fritsch P, Weinlich G, Jensen-Jarolim E. Anaphylaxis to camomile: clinical features and allergen cross-reactivity. *Clin Exp Allergy* 2000, 30(10):1436-1443
- Remme JJ, de Wit RFE. Kamillosan bei der Behandlung von Beingeschwüren. Bericht über eine Untersuchung (Übersetzung aus dem Niederländischen) *Recept* 1984, 4:57-58
- Rocha NF, Rios ER, Carvalho AM, Cerqueira GS, Lopes A, Leal LK, et al. Anti-nociceptive and anti-inflammatory activities of (-)- α -bisabolol in rodents. *Naunyn Schmiedebergs Arch Pharmacol* 2011, 384(6):525-533
- Romero-Jiménez M, Campos-Sánchez J, Analla M, Muñoz-Serrano A, Alonso-Moraga A. Genotoxicity and anti-genotoxicity of some traditional medicinal herbs. *Mutat Res* 2005, 585(1-2):147-155
- Rudzki E, Jalblonska S. Kamillosan® is a safe product of chamomile for topical application: results of patch testing consecutive patients with contact dermatitis. *J Dermatolog Treat* 2000, 11(3):161-163
- Rycroft RJG. Recurrent facial dermatitis from chamomile tea. *Contact Dermatitis* 2003, 48(4):229
- Safayhi H, Sabieraj J, Sailer ER, Ammon HP. Chamazulene: an antioxidant-type inhibitor of leukotriene B4 formation. *Planta Med* 1994, 60(5):410-413
- Salgueiro JB, Ardenghi P, Dias M, Ferreira MB, Izquierdo I, Medina JH. Anxiolytic natural and synthetic flavonoid ligands of the central benzodiazepine receptor have no effect on memory tasks in rats. *Pharmacol Biochem Behav* 1997, 58(4):887-891

Sauer H. Zur Wirksamkeit von Kamillosan® nach operativen Eingriffen/konservativer Behandlung von Nasennebenhöhlenerkrankungen. *Giatros HNO* 1990, 5 11:4-5

Savino F, Cresi F, Castagno E, Silvestro L, Oggero R. A randomized double-blind placebo controlled trial of a standardized extract of *Matricaria recutita*, *Foeniculum vulgare* and *Melissa officinalis* (ColiMil) in the treatment of breastfed colicky infants. *Phytother Res* 2005, 19(4):335-340

Scala G. Acute, short-lasting rhinitis due to camomile-scented toilet paper in patients allergic to compositae. *Int Arch Allergy Immunol* 2006, 139(4):330-331

Schilcher H. Die Kamille. Handbuch für Ärzte, Apotheker und andere Naturwissenschaftler. Wissenschaftliche Verlagsgesellschaft, Stuttgart 1987

Schmidt OP. Erfolge der Phytotherapie in der Pneumologie. In: Demling L, Nasemann T, Rosch W, editors. Erfahrungstherapie – späte Rechtfertigung. Internationales Symposium Wien, 30-31 Mai. Verlag G. Braun, Karlsruhe 1975, 17-27

Segal R, Pilote L. Warfarin interaction with *Matricaria chamomilla*. *CMAJ* 2006, 174(9):1281-1282

Shinomiya K, Inoue T, Utsu Y, Tokunaga S, Masuoka T, Ohmori A, et al. Hypnotic activities of chamomile and passiflora extracts in sleep-disturbed rats. *Biol Pharm Bull* 2005, 28(5):808-810

Shipochliev T, Dimitrov A, Aleksandrova E. [Anti-inflammatory action of a group of plant extracts.] *Vet Med Nauki* 1981, 18(6):87-94 [Bulgarian]

Smolinski AT, Pestka JJ. Modulation of lipopolysaccharide-induced proinflammatory cytokine production in vitro and in vivo by the herbal constituents apigenin (chamomile), ginsenoside Rb(1) (ginseng) and parthenolide (feverfew). *Food Chem Toxicol* 2003, 41(10):1381-1390

Soliman KM, Badeaa RI. Effect of oil extracted from some medicinal plants on different mycotoxicogenic fungi. *Food Chem Toxicol* 2002, 40(11):1669-1675

Stanev S, Zheljazkov V, Janculoff Y. Variation of chemical compounds in the essential oil from some native forms of chamomile (*Chamomilla recutita* L.). *Beitr Züchtungsforschung* 1996, 2:214-217

Stechele U. Kamillosan® Salbe zur Therapie von entzündlichen Hauterkrankungen im Säuglingsalter. In: Nasemann T, Patzelt-Wenzler R, editors. Kamillosan im Spiegel der Literatur. PMI Verlag, 1991, 21

Sticher O, Heilmann J, Zündorf I. Hänsel/Sticher Pharmakognosie – Phytopharmazie. 10th ed. Wissenschaftliche Verlagsgesellschaft, Stuttgart 2015

Stiegelmeier H. Therapie unspezifischer Magenbeschwerden mit Kamillosan. *Kassenarzt* 1978, 18: 3605-3606

Subiza J, Subiza JL, Hinojosa M, Garcia R, Jerez M, Valvidieso R, et al. Anaphylactic reaction after the ingestion of chamomile tea: A study of crossreactivity with other composite pollens. *J Allergy Clin Immunol* 1989, 84(3):353-358

Szelenyi I, Isaac O, Thiemer K. [Pharmacological experiments with compounds of chamomile. III. Experimental studies of the ulcerprotective effect of chamomile (author's transl).] *Planta Med* 1979, 35(3):218-227 [German]

Thien FC. Chamomile tea enema anaphylaxis. *Med J Aust* 2001, 175(1):54

Troll W, Patzelt-Wenzler R. Kamillosan-Therapie bei Erkrankungen im HNO-Bereich. *Giatros HNO* 1990, 5: 3-8

Tubaro A, Zilli C, Redaelli C, Della Loggia R. Evaluation of antiinflammatory activity of a chamomile extract after topical application. *Planta Med* 1984, 50(4):359

United States Pharmacopoeia 29 - National Formulary 24. Rockville, MD: United States Pharmacopeia Convention; 2006

Viegas D, Vaz FC, da Costa M. Utilizacao do extrato etanólico de flor de camomila em base cremosa no tratamento das dermatites amoniacais. *Pediatrica Moderna* 1996, 32:443-448

Vilaginès P, Delaveau P, Vilaginès R. [Inhibition of poliovirus growth by *Matricaria chamomilla* (L.) extract.] *C R Acad Sc* 1985, 301(6):289-294 [French]

Viola H, Wasowski C, Levi de Stein M, Wolfman C, Silveira R, Dajas F, *et al.* Apigenin, a component of *Matricaria recutita* flowers, is a central benzodiazepine receptors-ligand with anxiolytic effects. *Planta Med* 1995, 61(3):213-216

Weizman Z, Alkrinawi S, Goldfarb D, Bitran C. Efficacy of herbal tea preparation in infantile colic. *J Pediatr* 1993, 122(4):650-652

WHO monographs on selected medicinal plants. Vol 2. *Flos Chamomillae*. World Health Organization. Geneva 1999, 86-94

Wichtl M, Bauer R, editors. *Teedrogen und Phytopharmaka*. 5th ed. Wissenschaftliche Verlagsgesellschaft, Stuttgart 2009

Yamada K, Miura T, Mimaki Y, Sashida Y. Effect of inhalation of chamomile oil vapour on plasma ACTH level in ovariectomized-rat under restriction stress. *Biol Pharm Bull* 1996, 19(9):1244-1246

References assessed, but not cited in the assessment report:

- Al-Hashem FH. Gastroprotective effects of aqueous extract of *Chamomilla recutita* against ethanol-induced gastric ulcers. *Saudi Med J* 2010, 31(11):1211-1216
- Abebe W. Herbal medication: potential for adverse interactions with analgesic drugs. *J Clin Pharm Ther* 2002, 27(6):391-401
- Adam B, Liebregts T, Saadat-Gilani K, Vinson B, Holtmann G. Validation of the gastrointestinal symptom score for the assessment of symptoms in patients with functional dyspepsia. *Aliment Pharmacol Ther* 2005, 22(4):357-363
- Adam KP, Thiel R, Zapp J. Incorporation of 1-[1-¹³C] Deoxy-D-xylulose in Chamomile Sesquiterpenes. *Arch Biochem Biophys* 1999, 369(1):127-132
- Ahmad A, Misra LN. Isolation of Herniarin and other Constituents from Matricaria Chamomilla Flowers *Int J Pharmacogn* 1997, 35(2):121-125
- Ahmed AA, Abou Elela M. Highly oxygenated bisabolenes and an acetylene from *Matricaria aurea*. *Phytochemistry* 1999, 51(4):551-554
- Allescher HD. Functional dyspepsia-a multicausal disease and its therapy. *Phytomedicine* 2006, 13(Suppl 5):2-11
- Allescher HD, Wagner H. [STW 5/Iberogast: multi-target-action for treatment of functional dyspepsia and irritable bowel syndrome.] *Wien Med Wochenschr* 2007, 157(13-14):301-307 [German]
- Ammon HP, Kelber O, Okpanyi SN. Spasmolytic and tonic effect of Iberogast (STW 5) in intestinal smooth muscle. *Phytomedicine* 2006, 13(Suppl 5.):67-74
- Ammon HTP, Kaul R. Pharmacology of Chamomile and its Components. *Dtsch Apoth Ztg* 1992, 132(Suppl 27):1-26
- Ammon HTP, Sabieraj J, Kaul R. Kamille Mechanismus der antiphlogistischen Wirkung von Kamillenextrakten und – inhaltsstoffen. *Dtsch Apoth Ztg* 1996, 136(22):1821-1834
- Andreev VC. Anwendung von sebamed compact mit 2% Kamillenextrakt bei verschiedenen Dermatosen. *Ärztl Kosmetol* 1987, 17:452-456
- Anderson C, Lis-Balchin M. The Effect of Essential Oils on Childhood Atopic Eczema, 29th International Symposiumon Essentials Oils, Sept 6-9, 1998 Institut für Lebensmittelchemie Johann-Wolfgang-Goethe-Universität, Frankfurt/Main 1998, Abstract:2-9
- Appel K. Potenzielle Allergene in Kamillentee gefunden. *Act Dermatol* 2008, 34:203
- Appelt GD. Pharmacological aspects of selected herbs employed in Hispanic folk medicine in the San Luis Valley of Colorado, USA: I. *Ligusticum porteri* (Osha) and *Matricaria chamomilla* (Manzanilla). *J Ethnopharmacol* 1985, 13(1):51-55
- Arnold Dr. W. Über Kamille, Pfefferminze und Fenchel: Eine vergleichend, experimentell-pharmakologische und klinische Untersuchung. *Arch Exp Path Pharmak* 1927, 123:129-159
- Ashida Y, Nishimoto M, Matsushima A, Watanabe J, Hirata T. Molecular cloning and mRNA expression of geraniol-inducible genes in cultured shoot primordia of *Matricaria chamomilla*. *Biosci Biotechnol Biochem* 2002, 66(11):2511-2514
- Babenko NA, Shakhova EG. Effects of *Chamomilla recutita* flavonoids on age-related liver sphingolipid turnover in rats. *Exp Gerontol* 2006, 41(1):32-39

Bhaskaran N, Shukla S, Srivastava JK, Gupta S. Chamomile: an anti-inflammatory agent inhibits inducible nitric oxide synthase expression by blocking RelA/p65 activity. *Int J Mol Med* 2010, 26(6):935–940

Bauer K, Biller A, Verspohl EJ. Untersuchungen von Bestandteilen eines pflanzlichen Antidiarrhoikums auf spasmolytische Effekte. *Z Phytother* 2008, 29(4):169-172

Bauer R, Feldmann HU. Matricaria Chamomilla. *Dtsch Apoth* 1980, (32):5

Bauer R, Feldmann HU. Matricaria chamomilla. *Notabene Medici* 1980, 10(3):137-139

Becker H, Reichling J. Ist die Phytotherapie mit Baldrian und Kamille noch zeitgemäß? Teil I: Baldrian. *Z Phytother* 1982, II:2-19

Becker H, Reichling J. Ist die Phytotherapie mit Baldrian und Kamille noch zeitgemäß? Teil II: Kamille. *Dtsch Apoth Ztg* 1981, 121(23):1185-1190

Bedard M. Chamomile: More than a Fragrance? *Alternative Medicine* 2003, 136(1):14-15

Berger F. Synonyma-Lexikon der Heil- und Nutzpflanzen. Chapter: Chamomilla. Österreichische Apotheker Verlags GmbH, Wien 1981 (1955 Reprintdruck)

Berger F. Handbuch der Drogenkunde. Chapter: Flores Chamomillae Vulgaris. Maudrich Verlag, Wien 1954

Bianco MI, Luquez C, de Jong LIT, Fernandez RA. Presence of Clostridium botulinum spores in Matricaria chamomilla (chamomile) and its relationship with infant botulism. *Int J Food Microbiol* 2008, 121(3):357-360

Biller A. Immer trifft es die Kleinen: Pflanzliche Therapie kindlicher Durchfallerkrankungen. *Wien Med Wochenschr* 2007, 157(13-14):308-311

Birt DF, Walker B, Tibbels MG, Bresnick E. Anti-mutagenesis and anti-promotion by apigenin, robinetin and indole-3-carbinol. *Carcinogenesis* 1986, 7(6):959-963

Bisson W, Beiderbeck Reichling J. [Production of Essential Oils by Cell-Suspensions of Matricaria chamomilla in a Two Phase System.] *Planta Med* 1983, 47(3):164-168 [German]

Blaschek W, Ebel S, Hackenthal E, Holzgrabe U, Keller K, Reichling J, Schulz V, editors. HagerROM 2006: Hagers Handbuch der Drogen und Arzneistoffe. Springer Verlag, Berlin Heidelberg 2006

Bleimann H, Hartmann R. Therapie gastrointestinaler Funktionsstörungen mit Iberogast. *Kassenarzt* 1983, 23:52-59

Braun H, Frohne D. Heilpflanzen Lexikon für Ärzte und Apotheker. Chapter: Matricaria chamomilla. 5th ed. Gustav Fischer Verlag, Stuttgart 1987

Breinlich J, Scharnagel K. Pharmakologische Eigenschaften des EN-IN-Dicycloäthers aus Matricaria Chamomilla. Untersuchungen der entzündungshemmenden, antianaphylaktischen, spasmolytischen und bakteriostatischen Wirkung. *Arzneimittelforschung* 1968, 18:429-431

Bremer W, Fischer F, Nicolay K. Gastrointestinal Erkrankungen. *Z Allgemeinmed* 1983, 59(30):1706-1709

Broussard CS, Louik C, Honein MA, Mitchell AA. The National Birth Defect Prevention Study. Herbal use before and during pregnancy. *Am J Obstet Gynecol* 2010, 202:443 e1-6

Brückel MH, Gisevius G. Bei funktionellen Magenbeschwerden hilft Iberis amara. *Ärztliche Praxis* 1984, 36(21):494

Buchert D. Wirkung einer fixen Kombination bei gesicherter Non-Ulcus-Dyspepsie. *Z Phytother* 1994, 15: 24-25

Buck G. Kamilllobad - ein Kamillenextrakt zur äußereren Anwendung. *Z Allg Med* 1988, 64: 112-113

Buckle J. Use of aromatherapy as a complementary treatment for chronic pain. *Altern Ther Health Med* 1999, 5(5): 42-51

Budavari S, editor. The Merck Index. An Encyclopedia of Chemicals, Drugs, and Biologicals. Chapter: 5639 Matricaria. 11th ed. Merck, Rahway NJ 1989

Cahiers de l'Agence N°3 - Médicament à base de plantes - Ministry of Health and Family, Paris 1998

Carle R. Kamillenöl - Gewinnung und Qualitätsbeurteilung. *Dtsch Apoth Ztg* 1996, 136(26): 17-28

Carle R. Pflanzliche Antiphlogistika und Spasmolytika, Zwischen Anspruch und Wirklichkeit. *Dtsch Apoth Ztg* 1987, 127(43): 2206-2208

Carle R. Pflanzliche Antiphlogistika und Spasmolytika. *Z Phytother* 1988, 9: 67-76

Carle R, Dölle B, Müller W, Baumeister U. Thermospray liquid chromatography/mass spectrometry (TSP LC/MS): Analysis of acetylated apigenin-7-glucosides from *Chamomilla recutita*. *Pharmazie* 1993, 48: 304-306

Carle R, Dölle B, Reinhard E. A New Approach to the Production of Chamomile Extracts. *Planta Med* 1989, 55(6): 540-543

Carle R, Fleischhauer I, Beyer J, Reinhard E. Studies on the Origin of (-)-*a*-Bisabolol and Chamazulene in Chamomile Preparations; Part I. Investigations by Isotope Ratio Mass Spectrometry (IHMS). *Planta Med* 1990, 56(5): 455-460

Carle R, Fleischhauer I, Fehr D. Qualitätsbeurteilung von Kamillenölen. *Dtsch Apoth Ztg* 1987, 47: 2451-2457

Carle R, Gomaa K. Chamomile: A pharmacological and clinical profile. *Drugs Today* 1992b, 28(8): 559-565

Casterline CK. Allergy to Chamomile Tea. *JAMA* 1980, 244(4): 330-331

Cellarova E, Rycilova M, Seidelova A, Honcariv R. Comparison of Mitotic Activity and Growth in two Long Term Callus Cultures of *Matricaria recutita* L. *Acta Biotech* 1990, 10(3): 245-251

Cemek M, Kağa S, Simşek N, Büyükkokuoğlu ME, Konuk M. Antihyperglycemic and antioxidative potential of *Matricaria chamomilla* L. in streptozotocin-induced diabetic rats. *J Nat Med* 2008, 62(3): 284-293

Ceska O, Chaudhary SK, Warrington PJ, Ashwood-Smith MJ. Coumarins of chamomile, *Chamomilla recutita*. *Fitoterapia* 1992, 63(5): 387-394

Charuluxananan S, Sumethawattana P, Kosawiboonpol R, Somboonviboon W, Werawataganon T. Effectiveness of Lubrication of Endotracheal Tube Cuff with Chamomile-Extract for Prevention of Postoperative Sore Throat and Hoarseness. *J Med Assoc Thai* 2004, 87(Suppl 2): 185-189

Chizzola R, Michitsch H, Mitteregger US. Extractability of selected mineral and trace elements in infusions of chamomile. *Int J Food Sci Nutr* 2008, 59(6): 451-456

Cinco M, Banfi E, Tubaro A, Della Loggia R. A microbiological survey on the activity of a hydroalcoholic extract of camomile. *Int J Crude Drug Res* 1983, 21(4): 145-151

Czeczot H, Tudek B, Kusztelak J, Szymczyk T, Dobrowolska B, Glinkowska G, et al. Isolation and studies of the mutagenic activity in the Ames test of flavonoids naturally occurring in medical herbs. *Mutation Res* 1990, 240(3):209-216

Cérantola S, Kervarec N, Pichon R, Magné C, Bessieres MA, Deslandes E. NMR characterisation of inulin-type fructooligosaccharides as the major water-soluble carbohydrates from *Matricaria maritima* (L.). *Carbohydr Res* 2004, 339(14):2445-2449

Deutsches Arzneibuch 9th ed. Kamillenblüten. Deutscher Apotheker Verlag, Stuttgart 1986, 2061-2065

Deutsches Arzneibuch 7th ed. Kamillenblüten. Wissenschaftliche Verlagsgesellschaft, Stuttgart 1968, 933-938

Damschun S. Erfassung und Beschreibung vorhandener Sorten und leistungsstarker Herkünfte im Heil- und Gewürzpflanzenanbau. *Drogenreport* 1996, 9(14):31-32

De Jong NW, Vermeulen AM, Gerth van Wijk R, de Groot H. Occupational allergy caused by flowers. *Allergy* 1998, 53:204-209

De Pasquale R, Ragusa S, Iauk L, Barbera R, Galati EM. Effect of cadmium on germination, growth and active principle contents of *Matricaria recutita*, L. *Pharmacol Res Commun* 1988, 20(5):151-154

De Souza Reis LS, Frazatti-Gallina NM, de Lima Paoli R, Giuffrida R, Albas A, Oba E, et al. Efficiency of *Matricaria chamomilla* CH₁₂ and number of doses of rabies vaccine on the humoral immune response in cattle. *J Vet Sci* 2008, 9(4):433-435

De Souza Reis LS, Pardo PE, Oba E, Nascimento Kronka S, Frazatti-Gallina NM. *Matricaria chamomilla* CH₁₂ decreases handling stress in Nelore calves. *J Vet Sci* 2006, 7(2):189-192

Deininger R. Eine neue Methode zum Nachweis der entzündungswidrigen Wirkung des Azulens. *Arzneimittelforschung* 1956, 6(7):394-395

Della Loggia R. Lokale antiphlogistische Wirkung der Kamillen-Flavone. *Dtsch Apoth Ztg*, Suppl 1:9-11

Della Loggia R, Tubaro A, Dri P, Zilli C, Del Negro P. The Role of Flavonoids in the antiinflammatory activity of *Chamomilla recutita*. In: Cody V, Middleton E, Harborne JB. Plant Flavonoids in Biology and Medicine: Biochemical, Pharmacological and Structure-Activity Relationships. Symposium Buffalo NY, July 22-26. 1986, 481-484

Demling L. Erfahrungstherapie - späte Rechtfertigung. In: Demling L, Nasemann T, Rosch W. editors. Erfahrungstherapie - späte Rechtfertigung. Internationales Symposium Wien, 30-31 Mai. Verlag G. Braun, Karlsruhe 1975, 1-8

Detter A. Keimhemmstoffe als Arzneimittel? *Pharm Ztg* 1981, 126(23):1140-1142

Dragland S, Senoo H, Wake K, Holta K, Blomhoff R. Several culinary and medicinal herbs are important sources of dietary antioxidants. *J Nutr* 2003, 133(5):128-190

Eliasová A, Repcák M, Pastírová A. Quantitative changes of secondary metabolites of *Matricaria chamomilla* by abiotic stress. *Z Naturforsch C* 2004, 59(7-8):543-548

EMEA/Working Party on Herbal Medicinal Products Draft position statement on chamomilla containing herbal medicinal products. EMEA/HMPWP/345/03, 2003 4 S

ESCOP. Proposal for a European Monograph on the medicinal use of *Matricariae flos*. European Scientific Cooperative for Phytotherapy 1990, I.2.1-I.2.9

Exner J, Reichling J, Cole TC, Becker H. Methylated flavonoid-aglycones from "matricariae flos". *Planta Med* 1981, 41(2):198-200

Farideh ZZ, Bagher M, Ashraf A, Akram A, Kazem M. Effects of chamomile extract on biochemical and clinical parameters in a rat model of polycystic ovary syndrome. *J Reprod Infertil* 2010, 11(3):169-174

Fischer G. Heilkräuter und Arzneipflanzen. Chapter: Kamille. 5th ed. Karl F. Haug Verlag, Heidelberg 1978

Flaskamp E, Nonnenmacher G, Zimmermann G, Isaac O. Zur Stereochemie der Bisaboloide aus Matricaria chamomilla. *Planta Med* 1981, 42(6):106-107

Florido-Lopez JF, Gonzalez-Delgado P, Saenz de San Pedro B, Perez-Miranda C, Arias de Saavedra JM, Marin-Pozo J. Allergy to natural honeys and camomile tea. *Int Arch Allergy Immunol* 1995, 108(2):170-174

Fonseca FN, Tavares MF. Validation of a capillary electrophoresis method for the quantitative determination of free and total apigenin in extracts of *Chamomilla recutita*. *Phytochem Anal* 2004, 15(1):65-70

Fonseca FN, Tavares MF, Horváth C. Capillary electrochromatography of selected phenolic compounds of *Chamomilla recutita*. *J Chromatogr A* 2007, 1154(1-2):390-399

Förster CF, Süssmann HE, Patzelt-Wenzler R. [Optimization of the Barron ligature treatment of 2nd and 3rd-degree hemorrhoids using a therapeutic troika.] *Praxis* 1996, 85(46):1476-1481 [German]

Foti K, Nettis E, Panebianco R, Cassano N, Diafero A, Pia DP. Contact Urticaria from Matricaria chamomilla. *Contact Dermatitis* 2000, 42(6):360-361

Franz C, Fritz D, Ruhland E. Zur Qualität von Pfefferminz- und Kamillen-Teedroge aus dem Handel. *Planta Med* 1981, 42(6):132

Frohne D, Pfänder HJ. Giftpflanzen Ein Handbuch für Apotheker, Ärzte, Toxikologen und Biologen. Chapter: Asteraceae. 3rd ed. Wissenschaftliche Verlagsgesellschaft, Stuttgart 1987

Füller E, Sosa S, Tubaro A, Franz G, Della Loggia R. Anti-inflammatory activity of Chamomile polysaccharide. *Planta Med* 1993, 59:666-667

Füller E, Sosa S, Tubaro A, Franz G, Della Loggia R. Anti-inflammatory activity of Chamomile polysaccharide. *Pharm Pharmacol Lett* 2000, 2:86-87

Ganzer BM. Phytotherapie in der Kinderheilkunde hoch geschätzt. *PZ* 1990, 135(20):26

Ganzena M, Guggenberger M, Stuppner H, Zidorn C. Altitudinal variation of secondary metabolite profiles in flowering heads of Matricaria chamomilla cv. BONA. *Planta Med* 2008, 74(4):453-457

Gardiner P. Complementary, Holistic, and Integrative Medicine: Chamomile. *Pediatr Rev* 2007, 28(4):e16-e18

Gasic O, Lukic V, Adamovic D, Canak N. Variation in the content and the composition of the essential oils in flower heads of Matricaria chamomilla L. during its ontogenetical development. *Acta Pharm Hung* 1986, 56(6):283-288

Germann I, Hagelauer D, Kelber O, Vinson B, Laufer S, Weiser D, et al. Antioxidative properties of the gastrointestinal phytopharmaceutical remedy STW 5. *Phytomedicine* 2006, 13(Suppl 5):45-50

Gessner O, Orzechowski G. Gift und Arzneipflanzen von Mitteleuropa. 3rd ed. Carl Winter Verlag, Heidelberg 1974

Giligan NP. The palliation of nausea in hospice and palliative care patients with essential oils of *Pimpinella anisum* (aniseed), *Foeniculum vulgare* var. *Dulce* (sweet fennel), *Anthemis nobilis* (Roman chamomile) and *Mentha x piperita* (peppermint). *Int J Aromather* 2005, 15(4):163-167

Giordano-Labadie F, Schwarze HP, Bazex J. Allergic contact dermatitis from camomile used in phytotherapy. *Contact Dermatitis* 2000, 42(4):247

Gläsl H, Wagner H. Gaschromatographische Untersuchung von Arzneibuchdrogen. *Dtsch Apoth Ztg* 1976, 116(3):45-50

Glowania HJ, Raulin C, Swoboda M. Wirkung der Kamille in der Wundheilung – eine klinische Doppelblindstudie. *Z Hautkr* 1987, 62(17):1262-1271

Goldstein LH, Elias M, Ron-Avraham G, Biniaurishvili BZ, Madjar M, Kamargash I, et al. Consumption of herbal remedies and dietary supplements amongst patients hospitalized in medical wards. *Br J Clin Pharmacol* 2007, 64(3):373-80

Gomaa A, Hashem T, Mohamed M, Ashry E. Matricaria chamomilla extract inhibits both development of morphine dependence and expression of abstinence syndrome in rats. *J Pharmacol Sci* 2003, 92(1):50-55

Gould L, Reddy R, Gomprecht RF. Cardiac effects of chamomile tea. *J Clin Pharmacol* 1973, 13(11):475-479

Graf J. Herbal anti-inflammatory agents for skin disease. *Skin Therapy Lett* 2000, 5(4):3-5

Griffiths LA, Barrow A. Metabolism of flavonoid compounds in germ-free rats. *Biochem J* 1972, 130(4):1161-1162

Grochulski A, Borkowski B. Einfluss von Öl. Chamomillae in experimenteller Glomerulonephritis bei Kaninchen. *Planta Med* 1972, 21(3):289-292

Gundermann KJ, Godehardt E, Ulbrich M. Efficacy of a herbal preparation in patients with functional dyspepsia: A meta-analysis of double-blind, randomized, clinical trials. *Adv Ther* 2003, 20(1):43-49

Gundermann KJ, Godehardt E, Ulbrich M. Wirksamkeit eines pflanzlichen Kombinationspräparates bei funktioneller Dyspepsie. Metaanalyse randomisierter Doppelblind-Studien auf Basis eines validen gastrointestinalen Symptomprofils. *MMW Orig* 2004, 146:71-76

Hagenström UH, Spitzkowski A, Winkler U. Kamillenblüten--eine heterogene Droge (Beitrag zur Herstellung von Teegemischen mit Matricariae flos) Chamomile blooms--a heterogenous drug (manufacture of mixed teas with Matricariae flos). *Pharmazie* 1990, 45(3):211-212

Hahn B, Hözl J. Resorption, Verteilung und Metabolismus von 14-C-Levomentol in der Haut. *Arzneim.-Forsch./Drug Res* 1987, 37(I), (6):716-720

Hamon NW. The Chamomiles. *CPJ-RPC* 1989, 612-615

Hänsel R, Haas H. Therapie mit Phytopharmaka. Chapter: Kamillenblüten. Springer Verlag, Berlin 1983

Hänsel R, Haas H. Therapie mit Phytopharmaka. Chapter: Kamillenblüten. Springer-Verlag, Berlin 1984

Hänsel R, Keller K, Rimpler H, Schneider G. editors. Hagers Handbuch der Pharmazeutischen Praxis. Vol 5. 4th ed. Springer-Verlag, Berlin 1975

Hänsel R, Keller K, Rimpler H, Schneider G. editors. Hagers Handbuch der Pharmazeutischen Praxis. Drogen A-D. Vol 4. 5th ed. Springer-Verlag, Berlin 1992

Harnischfeger G, Stolze H. Kamille. Notabene medici. *Journal für Ärzte* 1982, Reprint 12(10):831-840

Hatzky K. Über die Anwendung der Kamille in der inneren Medizin als Chamomillysatum Bürger. *Med Klin Klin* 1930, 22:819-820

Hauben BM. Sesquiterpene Lactones - *Chamomilla recutita*. In: De Smet PAGM. Adverse Effects of Herbal Drugs. Vol 1. Springer Verlag 1992, 243-248

Hauben BM. Allergiepflanzen, Pflanzenallergene: Handbuch und Atlas der allergieinduzierenden Wild- und Kulturpflanzen. Chapter: *Chamomilla recutita*. 2nd ed. Ecomed, Landsberg 1988

Hauben BM. Kompositen-Allergie. Eine Auswertung der Testergebnisse von 1985-1990. *Akt Dermatol* 1992, 18:96-101

Hauben BM. The sensitizing capacity of Compositae plants. III. Test results and cross-reactions in Compositae-sensitive patients. *Dermatologica* 1979, 159(1):1-11

Hauben BM, Busker E, Carle E. Über das Sensibilisierungsvermögen von Compositenarten: VII. Experimentelle Untersuchungen mit Auszügen und Inhaltsstoffen von *Chamomilla recutita* (L.) Rauschert und *Anthemis corula* L. *Planta Med* 1984, 50(3):229-234

Heidari MR, Dadollahi Z, Mehrabani M, Mehrabi H, Pourzadeh-Hosseini M, Behravan E, et al. Study of antiseizure effects of *Matricaria recutita* extract in mice. *Ann N Y Acad Sci* 2009, 1171:300-304

Hegyi E, Doležalová A, Danilla T, Manovská M, Gejdoš M, Danyszová J, et al. Inzidenz von Kontaktallergien in der West-Slowakei. *Derm Beruf Umwelt* 1994, 42:148-152

Hegyi E, Hudaková G. Inzidenz der Kontaktensitivität auf Kontaktallergene der Standardreihe bei Patienten über 40 Jahre mit besonderer Berücksichtigung der Frauen. *Derm Beruf Umwelt* 1997, 45(1):6-11

Heilman J, Merfort I, Hagedorn U, Lippold BC. *In vivo* skin penetration studies of Chamomile Flavonoids. *Planta Med* 1993, 59:638

Heinle H, Hagelauer D, Pascht U, Kelber O, Weiser D. Intestinal spasmolytic effects of STW 5 (Iberogast) and its components. *Phytomedicine* 2006, 13(Suppl 5):75-79

Heinrich U, Tronnier H. Hautglättungsstudie mit einem Hautfunktionsöl. *PZ* 1992, 137(25):36-38

Hempel B. Co₂-extract from German Chamomile-proposed medicinal use. *Act Hort* 1999, 503:15-19

Heuskin S, Godin B, Leroy P, Capella Q, Wathelet JP, Verheggen F, et al. Fast gas chromatography characterisation of purified semiochemicals from essential oils of *Matricaria chamomilla* L. (Asteraceae) and *Nepeta cataria* L. (Lamiaceae). *J Chromatogr A* 2009, 1216(14):2768-2775

Hinz HL, Müller-Schärer H. Suitability of two root-mining weevils for the biological control of scentless chamomile, *Tripleurospermum perforatum*, with special regard to potential non-target effects. *Bull Entomol Res* 2000, 90(6):497-508

Hitziger T, Höll P, Ramadan M, Dettmering D, Imming P, Hempel B. Die alte junge Kamille. *Phytopharmazie* 2003, 5:22-30

Hofmann HA. Kamillosan liquidum bei Dickdarmerkrankungen. *Fortschritte der Therapie* 1926, 5:156-157

Hohenester B, Rühl A, Kelber O, Schemann M. The herbal preparation STW 5 (Iberogast) has potent and region-specific effects on gastric motility. *Neurogastroenterol Motil* 2004, 16(6):765-773

Holtmann G, Adam B, Vinson B. Evidenz-basierte Medizin und Phytotherapie bei funktioneller Dyspepsie und Reizdarmsyndrom: Eine systematische Analyse der verfügbaren Evidenz zum Präparat Iberogast. *Wiener Med Wochenschr* 2004, 154:528-534

Höltzel C. Tee und Teezubereitungen in der Apotheke. *Dtsch Apoth Ztg* 1984, 124(48):2479-248

Holz W, Miethling H. Wirkstoffe in wässrigen Kamillenaufgüßen. *Pharmazie* 1994, 49(1):53-55

Hölzl J, Hahn B. Präparation von [¹⁴C](-)- α -Bisabolol mit Chamomilla recutita (L.) Rauschert. *Dtsch Apoth Ztg* 1985, Suppl 1:31-38

Hölzl J, Ghassemi N, Hahn B. Preparation of ¹⁴C-Spiro Ethers by Chamomile and their Use by an Investigation of Absorption. *Planta Med* 1986, 52(6):553

Homöopathisches Arzneibuch. Chamomilla recutita. Amtliche Ausgabe. 4th Nachtrag. Deutscher Apotheker Verlag, 1985, 131-133

Hope MF. Kapitel: Matricaria. 692-694

Hoppe HA. Drogenkunde. Chapter: Matricaria chamomilla. Vol 1. 8th ed. Walter de Gruyter Verlag, Berlin 1975

Ianoli P, Avallone R, Baraldi M. Behavioral Characterisation of the flavonoids apigenin and chrysin. *Fitoterapia* 2000, 71(Suppl 1):5117-23

Illing G, Sajthy G. Zur antiemetischen Wirkung von Iberogast. *Kassenarzt* 1984, 17:46-49

Isaac O. Fortschritte in der Kamillenforschung. *Physikalische Medizin und Rehabilitation* 1975, 16(11):223-229

Isaac O. Die Kamillentherapie - Erfahrung und Bestätigung. *Dtsch Apoth Ztg* 1980, 120(13):567-570

Isaac O. Fortschritte in der Kamillenforschung. *Dtsch Apoth Ztg* 1974, 114(7):255-260

Isaac O. Pharmakologische Untersuchungen von Kamillen-Inhaltsstoffen: I. Zur Pharmakologie des (-)- α -Bisabolols und der Bisabololoxide. *Planta Med* 1979, 35(2):118-124

Isaac O, Kristen G. Alte und neue Wege der Kamillentherapie. Die Kamille als Beispiel für moderne Arzneipflanzenforschung. *Med Welt* 1980, 31(31-32):1145-1149

Isaac O, Schimpke H. Wissenswertes für die pharmazeutische Praxis. *Mitt Dtsch Pharm Ges DDR* 1965, 35(8):133-147

Isaac O, Schneider Dr. H, Eggenschwiler H. Beitrag zur Kenntnis des Kamillen-Bisabolos. *Dtsch Apoth Ztg* 1968, 108(9):293-298

Isar HJ. Alte und neue Anwendungsmöglichkeiten der deutschen Kamille in der dermatologischen und urologischen Praxis. *Dermatol Wschr* 1930, 21:712-5

Izumi S, Takashima O, Hirata T. Geraniol is a potent inducer of apoptosis-like cell death in the cultured shoot primordia of Matricaria chamomilla. *Biochem Biophys Res Commun* 1999, 259(3):519-522

Jakovlev V, Isaac O, Thiemer K, Kunde R. Pharmakologische Untersuchungen von Kamillen-Inhaltsstoffen: II. Neue Untersuchungen zur antiphlogistischen Wirkung des (-)- α -Bisabolols und der Bisabololoxide. *Planta Med* 1979, 35(2):125-140

Jakovlev V, von Schlichtegroll A. Zur entzündungshemmenden Wirkung von(-)-a-Bisabolol, einem wesentlichen Bestandteil des Kamillenöls. *Arzneimittelforschung/Drug Res* 1969, 19:615-616

Jakovlev V, Isaac O, Flaskamp E. [Pharmacologic studies on chamomile compounds. VI. Studies on the antiphlogistic effect of chamazulene and matricine.] *Planta Med* 1983, 49(2):67-73 [German]

Jarrahi M, Vafaei AA, Taherian AA, Miladi H, Rashidi Pour A. Evaluation of topical Matricaria chamomilla extract activity on linear incisional wound healing in albino rats. *Nat Prod Res* 2008, 22(14):1197-1202

Johri AK, Srivastava LJ, Singh JM, Rana RC. Effect of Row Spacings and Nitrogen Levels of Flower and Essential Oil Yield in German Chamomile. *Indian Perfumer* 1991, 35(2):93-96

Kai H, Kimoto A, Ishibashi A, Marsuzaki M, Takahama K, Miyara T. Involvement of divalent cation in the basal secretion of mucus glycoconjugates by rabbit trachea and dog gastric mucosa. *Eur J Pharmacol* 1990, 183(3):728-729

Kaiser CS, Hömpf H, Schmidt PC. Supercritical Carbon Dioxide Extraction of Chamomile Flowers: Extraction Efficiency, Stability, and In-line Inclusion of Chamomile-Carbon Dioxide Extract in β-cyclodextrin. *Phytochem Anal* 2004, 15(4):249-256

Kanda Y, Richards RG, Handwerger S. Apolipoprotein A-1 stimulates human placental lactogen release by activation of MAP Kinase. *Mol Cell Endocrinol* 1998, 143(1-2):125-131

Karbalay-Doust S, Noorafshan A. Antiulcerogenic Effects of Matricaria Chamomilla Extract in Experimental Gastric Ulcer in Mice. *Iran J Med Sci* 2009, 34(3):198-203

Kato A, Minoshima Y, Yamamoto J, Adachi I, Watson AA, Nash RJ. Protective effects of dietary chamomile tea on diabetic complications. *J Agric Food Chem* 2008, 56(17):8206-8211

Katz R. Therapeutische Erfahrungen mit Kamillosan. *Fortschr Med* 1928, 46:388-391

Kelber O, Wittwer A, Lapke C, Kroll U, Weiser D, Okpanyi SN, et al. Ex vivo/in vitro absorption of STW 5 (Iberogast) and its extract components. *Phytomedicine* 2006, 13(Suppl 5):107-113

Kesmati M, Barfinejad N, Moghadam HF. Effect of matricaria recutita on acute pain in the presence and absence of sex hormones. *J Res Med Sci* 2007, 12(4):190-197

Kesmati M, Zadeh A, Moghaddam F. Study of Benzodiazepine-like effects of Matricaria recutita on Morphine withdrawal syndrome in adult male rats. *Pak J Med Sci* 2008, 24(5):735-739

Ketel WG. Allergy to Matricaria chamomilla. *Contact Dermatitis* 1987, 16(1):50-51

Khayyal MT, Seif-EI-Nasr M, EI-Ghazaly MA, Okpanyi SN, Kelber O, Weiser D. Mechanisms involved in the gastro-protective effect of STW 5 (Iberogast) and its components against ulcers and rebound activity. *Phytomedicine* 2006, 13(Suppl 5):56-66

Kobayashi Y, Nakano Y, Inayama K, Sakai A, Kamiya T. Dietary intake of the flower extracts of German Chamomile (*Matricaria recutita* L.) inhibited compound 48/80-induced itch-scratch responses in mice. *Phytomedicine* 2003, 10(8):657-664

Kobayashi Y, Suzuki A, Kohayashi A, Kasai A, Ogata Y, Kumada Y. Suppression of Sensory Irritation by Chamomile Essential Oil und Its Active Component - Bisabololoxide A. *Niigata University of Pharmacy and Applied Sciences* 2007, 163-174

Koch C, Reichling J, Kehm R, Scharaf MM, Zentgraf H, Schneele J, et al. Efficacy of anise oil, dwarf-pine oil and chamomile oil against thymidine-kinase-positive and thymidine-kinase-negative herpesviruses. *J Pharm Pharmacol* 2008, 60(11):1545-1550

Koßowicz J, Siebert A, Fiedler W, Böhm R. Kamille und Schafgarbe "unter Beobachtung".
Therapiewoche 1996, 34:1857-1861

Kováčik J, Bačkor M, Kaduková J. Physiological Responses of *Matricaria chamomilla* to Cadmium and Copper Excess. *Environ Toxicol* 2007, 23(1):123-130

Kováčik J, Bačkor M. Oxidative status of *Matricaria chamomilla* plants related to cadmium and copper uptake. *Ecotoxicology* 2008, 17(6):471-479

Kováčik J, Bačkor M. Phenylalanine Ammonia-Lyase and Phenolic Compounds in Chamomile Tolerance to Cadmium and Copper Excess. *Water Air Soil Pollut* 2007, 185:185-193

Kováčik J, Klejdus B, Bačkor M. Nitric oxide signals ROS scavenger-mediated enhancement of PAL activity in nitrogen-deficient *Matricaria chamomilla* roots: side effects of scavengers. *Free Radic Biol Med* 2009, 46(12):1686-93

Kováčik J, Klejdus B. Dynamics of phenolic acids and lignin accumulation in metal-treated *Matricaria chamomilla* roots. *Plant Cell Rep* 2008, 27(3):605-615

Kováčik J, Klejdus B, Hedbavny J, Bačkor M. Salicylic acid alleviates NaCl-induced changes in the metabolism of *Matricaria chamomilla* plants. *Ecotoxicology* 2009, 18(5):544-554

Kováčik J, Klejdus B, Kaduková J, Bačkor M. Physiology of *Matricaria chamomilla* exposed to nickel excess. *Ecotoxicol Environ Saf* 2009, 72(2):603-609

Kováčik J, Grúz J, Bačkor M, Strnad M, Repčák M. Salicylic acid-induced changes to growth and phenolic metabolism in *Matricaria chamomilla* plants. *Plant Cell Rep* 2009, 28(1):135-143

Kowalzig H. Erfahrungen mit "Kamillosan", einer neuen Anwendungsart der Kamille. *Münch Med Wochenschr* 1922, 2:49-50

Král'ová K, Masarovičová E. *Hypericum perforatum* L. and *Chamomilla recutita* (L.) Rausch.-- accumulators of some toxic metals. *Pharmazie* 2003, 58(5):359-360

Kraft K. Aromatherapie - Was ist gesichert? *EHK* 2008, 57:273-278

Kraft K. Wunden und stumpfe Traumen. *Z Phytother* 2008, 29:178-180

Kroon LA. Drug interactions with smoking. *Am J Health Syst Pharm* 2007, 64(18):1917–1921

Kunde R, Isaac O. Über die Flavone der Kamille und ein neues acetyliertes Apigenin-7-glucosid. *Planta Med* 1979, 37(2):124-130

Kupfersztain C, Rotem C, Fagot R, Kaplan B. The immediate effect of natural plant extract, Angelica Sinensis and Matricaria Chamomilla (Clirnex) for the treatment of hot flushes during menopause. A preliminary report. *Clin Exp Obstet Gynecol* 2003, 30(4):203-206

Kuo ML, Yang NC. Reversion of V-H-Ras-transformed NIH 3T3 cells by apigenin through inhibiting mitogen activated protein-kinase and its downstream oncogenes. *Biochem Biophys Res Commun* 1995, 212(3):767-775

Kurita N, Miyaji M, Kurane R, Takahara Y. Antifungal activity of components of essential oils. *Agric Biol Chem* 1981, 45(4):945-952

Kyokong O, Charuluxananan S, Muangmingsuk V, Rodanant O, Subornsug K, Punyasang W. Efficacy of chamomile-extract spray for prevention of post-operative sore throat. *J Med Assoc Thai* 2002, 85(Suppl 1):180-185

Lahijani MSS, Kateb HRR, Heady R, Yazdani D. The effect of German chamomile (*Marticaria recutita* L.) extract and tea tree (*Melaleuca alternifolia* L.) oil used as irrigants on removal of smear layer: a scanning electron microscopy study. *Int Endod J* 2006, 39(3):190-195

Latz B. Über meine Erfahrungen mit Kamille in besonderer Anwendungsform. *Fortschritte der Therapie* 1927, 22: 796-799

Lenk S. Behandlung mit Kamillenblütenextrakt. *TW Urologie Kompakt* 1996, 11-12

Lepley DM, Li B, Birt DF, Pelling JC. The chemopreventive flavonoid apigenin induces G2/M arrest in keratinocytes. *Carcinogenesis* 1996, 17(11):2367-2375

Leslie GB, Salmon G. Repeated dose toxicity studies and reproductive studies on nine biostrath herbal remedies. *Swiss Med* 1979, 1: 43-45

Linde H, Cramer G. (-)-alpha-Bisabolol und cis-2-[Hexadiin-(2,4)-yilden]-1,6-dioxaspiro-[4,4]-nonen-(3) in handelsüblichen Kamillen-Extrakten. *Arzneimittelforschung* 1972, 22(3):583-585

Liu CY, Müller MH, Glatzle J, Weiser D, Kelber O, Enck P, et al. The herbal preparation STW 5 (Iberogast) desensitizes intestinal afferents in the rat small intestine. *Neurogastroenterol Motil* 2004, 16(6): 759-764

Liu JP, Yang M, Liu YX, Wei ML, Grimsbaard S. Herbal medicines for treatment of irritable bowel syndrome (Review). *Cochrane Database Syst Rev* 2006, (1):CD004116.

Lu T, Cantrell CL, Robbs SL, Franzblau SG, Fische NH. Antimycobacterial Matricaria Esters and Lactones from Astereae Species. *Planta Med* 1998, 64:665-667

Luo LH, Kim HJ, Nguyen DH, Lee HB, Lee NH, Kim EK. Depigmentation of melanocytes by (2Z,8Z)-matricaria acid methyl ester isolated from *Erigeron breviscapus*. *Biol Pharm Bull* 2009, 32(6):1091-1094

Luppold E. Matricaria Chamomilla-eine alte und neue Arzneipflanze. *Pharm Unserer Zeit* 1984, 13(3):65-70

Ma CM, Winsor L, Daneshbalab M. Quantification of spiroether isomers and herniarin of different parts of Matricaria matricarioides and flowers of Chamaemelum nobile. *Phytochem Anal* 2007, 18(1):42-49

Mac Lean N, Hübner-Steiner U. Behandlung arzneimittelbedingter Magen-Darm-Beschwerden. *Fortschr Med* 1987, 105(12):239-242

Mac Lean N, Hübner-Steiner U. Behandlung gastrointestinaler Beschwerden aufgrund von Arzneimittelunverträglichkeit mit dem Phytotherapeutikum Iberogast. *Gastro-Enterो-Hepatol* 1985, 3(6):3-8

Macander PJ. Flavonoids Affect Acetylcholine, Prostaglandin E2 and Antigen-Mediated Smooth Muscle Contraction. *Prog Clin Biol Res* 1986, 213:489-492

Madaus G. Lehrbuch der biologischen Heilmittel. Vol 3. Georg Olms Verlag, Hildesheim New York 1938 (1979 2nd reprint)

Maday E, Marczal G, Lemberkovics E, Muskath Z, Szöke E. *In vitro* and *in vivo* study of the essential oil of chamomile. 46th Kongress der Gesellschaft für Arzneipflanzeforschung "Quality of Medicinal Plants and Herbal Medicinal Products" Wien C08. 1998

Maday E, Szöke E, Muskáth Z, Lemberkovics E. A study of the production of essential oils in chamomile hairy root cultures. *Eur J Drug Metab Pharmacokinet* 1999 24(4):303-308

Madisch A, Holtmann G, Mayr G, Vinson B, Hotz J. Treatment of Functional Dyspepsia with a Herbal Preparation. *Digestion* 2004, 69:45-52

Madisch A, Holtmann G, Plein K, Hotz J. Treatment of irritable bowel syndrome with herbal preparations: results of a double-blind, randomized, placebo-controlled, multi-centre trial. *Aliment Pharmacol Ther* 2004, 19(3):271-279

Madisch A, Melderis H, Mayr G, Sassin I, Hotz J. Ein Phytotherapeutikum und seine modifizierte Rezeptur bei funktioneller Dyspepsie. *Z Gastroenterol* 2001, 39:511-517

Magiatis P, Michaelakis A, Skaltsounis AL, Haroutounian SA. Volatile secondary metabolite pattern of callus cultures of *Chamomilla recutita*. *Nat Prod Lett* 2001, 15(2):125-130

Maier R, Carle R, Kreis W, Reinhard E. Purification and Characterization of a Flavone 7 -O-Glucoside-Specific Glucosidase from Ligulate Florets of *Chamomilla recutita*. *Planta Med* 1993, 59(5):436-441

Maier R, Kreis W, Carle R, Reinhard E. A Flavone-Glucoside-Cleaving beta-Glucosidase from *Chamomilla recutita*. *Planta Med* 1991, 57(3):297-298

Mares D, Romagnoli C, Bruni A. Antidermatophytic Activity of Herniarin in Preparations of *Chamomilla recutita* (L.) Rauschert. *Plantes medicinales et phytotherapie* 1993, 26(2):91-100

Martindale W. The Extra Pharmacopoeia. Chapter: 4668 Matricaria Flowers, Essential Oils and Aromatic Carminatives. 28th ed. The Pharmaceutical Press, London 1982

Mayer JG, Uehlecke B, Saum K OSB. Das große Handbuch der Klosterheilkunde. Weltbild, Augsburg 2008

Mazokopakis EE, Vrentzos GE, Papadakis JA, Babalis DE, Ganotakis ES. Wild chamomile (*Matricaria recutita* L.) mouthwashes in methotrexate-induced oral mucositis. *Phytomedicine*. 2005, 12(1-2):25-27

McGeorge BC, Steele MC. Allergic contact dermatitis of the nipple from Roman chamomile ointment. *Contact Dermatitis* 1991, 24(2):139-140

McKay DL, Blumberg JB. A Review of the Bioactivity and potential Health Benefits of Chamomile Tea (*Matricaria recutita* L.). *Phytother Res* 2006, 20(7):519-530

Medić-Sarić M, Stanić G, Bosnjak I. The use of information theory and numerical taxonomy methods for evaluating the quality of thin-layer chromatographic separations of flavonoid constituents of *Matricariae flos*. *Pharmazie* 2001, 56(2):156-159

Melzer J, Iten F, Reichling J, Saller R. Iberis amara L. and Iberogast – Results of a systematic review concerning functional dyspepsia. *J Herb Pharmacother* 2004, 4(4):51-59

Melzer J, Rösch W, Reichling J, Brignoli R, Saller R. Meta-Analysis: phytotherapy of functional dyspepsia with the herbal drug preparation STW 5 (Iberogast). *Aliment Pharmacol Ther* 2004, 20(11-12):1279-1287

Melzig MF. Hemmung der neutrophilen Elastase durch ätherische Öle. *Z Phytother* 2006, 27:227-230

Messerschmidt W. Vergleichende Untersuchungen von wässrigen und äthanolischem-wässrigen Kamillenpräparaten des Handels. *Dtsch Apoth Ztg* 1973, 20: 745-748

Meyer A, Zimmermann S, Hempel B, Imming P. Anthecotulide: purification, analytical data, absence from chamomile preparations, stability and reactivity, and anti-infective testing. *J Nat Prod* 2005, 68(3):432-434

Michael S, Kelber O, Hauschmidt S, Spanel-Borowski K, Nieber K. Inhibition of inflammation-induced alterations in rat small intestine by the herbal preparations STW 5 and STW 6. *Phytomedicine* 2009, 16(2-3):161-171

Miethe H, Holz W. Wirkstoffe in wäßrigen Kamillenaufgüssen, 1. Mitteilung: Freisetzungskinetiken ausgewählter Flavonoidglycoside. *Pharmazie* 1989, 44:784-785

Mohr T, Hecht H, Eichhorn H. Vergleichende Untersuchungen einer verbesserten Pflückmaschine zur Gewinnung von Blütendrogen der Echten Kamille, Ringelblume sowie Johanniskraut. *Drogenreport* 1996, 9(14):15-23

Morck H. Drogenkunde. Chapter: Kamillenblüten. Georg Thieme Verlag, Stuttgart 1996

Morgan N. Strategies for Colic. *Birth Gaz* 1996, 12(4):18-23

Moss M, Howarth R, Wilkinson L, Wesnes K. Expectancy and the aroma of Roman chamomile influence mood and cognition in healthy volunteers. *Int J Aromather* 2006, 16:63-73

Moussally K, Bérard A. Exposure to herbal products during pregnancy and the risk of preterm birth (Letter to the editors). *Eur J Obstet Gynecol Reprod Biol* 2010, 150(1):107-108

Müller MH, Liu CY, Glatzle J, Weiser D, Kelber O, Enck P, et al. STW 5 (Iberogast) reduces afferent sensitivity in the rat small intestine. *Phytomedicine* 2006, 13(Suppl 5):100-106

Münzel M. Okkulte Sinusitis des Kindes. *Selecta* 1975, 24:2258-2260

Mulinacci Neander KD, Hesse F. Kamillentees in der Mundpflege. *Pflege Aktuell* 2004, 58:350-351

Nerseseyn AK. Influence of chamomile essential oil (CO) on SCE induced by two strong mutagens. (Letter to the Editor). *Toxicology Letters* 2004, 150:229-230

Newsholme SJ, Kellerman TS, Welman WG. Pathology of a nervous disorder ("pushing disease or "stootsiekte") in cattle caused by the plant Matricaria nigellifolia DC. (Asteraceae). *Onderstepoort J Vet Res* 1984, 51(2):119-127

Nicolay K. Funktionelle Gastroenteropathien im therapeutischen Blindvergleich von Metoclopramid mit dem Phytopharmacon Iberogast. *Gastro-Enterो-Hepatol* 1984, 2:24-28

Niederhofer H. Erste Ergebnisse einer Anwendung von echter Kamille bei autistischen Patienten. *Z Phytother* 2008, 29:275-276

Niederhofer H. Observational study: *Matricaria chamomilla* may improve some symptoms of attention-deficit hyperactivity disorder. *Phytomedicine* 2009, 16(4):284-286

Nissen HP, Patzelt-Wenzler R, Hose. Scavenger and antioxidant properties of special chamomile extracts. 2nd International Congress Phytomedicine and the 7th Congress of the Society of Phytotherapy in Cooperation with ESCOP, 1996

Noonan K, Arensman R, Hoover JD. Herbal Medication Use in the pediatric surgical patient. *J Pediatr Surg* 2004, 39(3):500-503

O'Hara M, Kiefer D, Farrel K, Kemper K. A review of 12 commonly used medicinal herbs. *Arch Fam Med* 1998, 7(6):523-536 Comment in: *Arch Fam Med* 1999, 8(5):376

Ohms P. Jeden Tag Tabletten: Wie lange macht der Magen mit. *Ärztliche Praxis* 1983, 35(99):3109-3110

Okpanyi SN, Mark M, Wahl MA. Gastrointestinal motility modulation with Iberogast. *Acta Hortic* 1993, 332:227-235

- Olivares M, Pizarro F, Hertrampf E, Fuenmayor G, Estévez E. Iran absorption from wheat flour: effects of lemonade and chamomile infusion. *Nutrition* 2007, 23:296-300
- Ożarowski A, Łaćucki J, Gąsiorowska K, Azulan. Herbal medicines, Guidebook 1978
- Pahlow M. Das große Buch der Heilpflanzen. Bechtermünz Verlag, München 2001
- Pauli A. α-Bisabolol from Chamomile - A specific ergosterol biosynthesis inhibitor? *Int J Aromather* 2006, 16:21-25
- Paulsen E, Andersen KE, Hausen BM. Compositae dermatitis in a Danish dermatology department in one year. Results of routine patch testing with the sesquiterpene lactone mix supplemented with aimed patch testing with extracts and sesquiterpene lactones of Compositae plants. *Contact Dermatitis* 1993, 29(1):6-10
- Pavesi VC, Lopez TC, Martins MA, Sant'Ana Filho M, Bussadori SK, Fernandes KP, et al. Healing action of topical chamomile on 5-fluoracil induced oral mucositis in hamster. *Support Care Cancer* 2011, 19(5):639-646
- Pavlovič A, Masarovičová E, Králl'ová K, Kubová J. Response of Chamomile Plants (*Matricaria recutita* L.) to Cadmium Treatment. *Bull Environ Contam Toxicol* 2006, 77(5):763-771
- Pereira RP, Fachilletto R, de Souza Prestes A, Puntel RL, Santos da Silva GN, Heinzmann BM, et al. Antioxidant Effects of Different Extracts from *Melissa officinalis*, *Matricaria recutita* and *Cymbopogon citratus*. *Neurochem Res* 2009, 34(5):973-983
- Petroianu G, Szőke E, Kalz H, Szegi P, Laufer R, Benko B, et al. Monitoring by HPLC of chamomile flavonoids exposed to rat liver microsomal metabolism. *Open Med Chem J* 2009, 3:1-7
- Pevny I. Kamillen -und Zitronenöl-Allergie. *Akt Dermatol* 1984, 10:56-57
- Pharmacopoeia Helvetica. 9th German ed. Kamillenblüten. 1993
- Pilichiewicz AN, Horowitz M, Russo A, Maddox AF, Jones KL, Schemann M, et al. Effects of Iberogast on proximal gastric volume, antropyloroduodenal motility and gastric emptying in healthy men. *Am J Gastroenterol* 2007, 102:1-8
- Pinto SA, Bohland E, Coelho Cde P, Morgulis MS, Bonamin LV. An animal model for the study of Chamomilla in stress and depression. pilot study. *Homeopathy* 2008, 97(3):141-144
- Pirali-Kheirabadi K, Razzaghi-Abyaneh M. Biological activities of chamomHe (*Matricaria chamomile*) flowers extract against the survival and egg laying of the cattle fever tick (Acari Ixodidae). *J Zhejiang Univ Sci B* 2007, 8(9):693-696
- Pirzad A, Alyari H, Shakiba MR, Zehtab-Salrnasi S, Mohammadi A. Essential Oil Content and Composition of Gennan Chamomile (*Matricaria chamomilla* L.) at Different Irrigation Regimes. *J Agron* 2006, 5(3):451-455
- Presibella MM, Santos CAM, Weffort-Santos AM. In Vitro Anticancer Activity of Chamomilla recutita Hydroethanol Extract. *Pharm Biol* 2007, 45(2):124-130
- Radaelli C, Formentini L, Santaniello E. Evaluation of Coumarins and Dicycloethers in *Matricaria chamomilla* Flowers and Chamomile Extracts by Reverse-Phase HPLC. Poster. International Research Congress on Natural Products as Medicinal Agents, 1980
- Radaelli C, Formentini L, Santaniello E. Evaluation of Coumarins and Dicycloethers in *Matricaria chamomilla* Flowers and Chamomile Extracts by Reverse-Phase HPLC. *Planta Med* 1981, 42(6):130

Raedsch R, Hanisch J, Bock P, Sibaev A, Vinson B, Gundermann KJ. Wirksamkeit und Unbedenklichkeit des Phytopharmakons STW 5 versus Metoclopramid oral bei funktioneller Dyspepsie unter Praxisbedingungen. *Z Gastroenterol* 2007, 45:1041

Ramadan M, Goeters S, Watzer B, Krause E, Lohmann K, Bauer R, et al. Chamazulene carboxylic acid and matricin: a natural profen and its natural prodrug, identified through similarity to synthetic drug substances. *J Nat Prod* 2006, 69(7):1041-1045

Ramirez-Durón R, Ceniceros-Almaguer L, Salazar-Aranda R, Salazar-Cavazos L, Waksman de Torres N. Evaluation of thin-layer chromatography methods for quality control of commercial products containing *Aesculus hippocastanum*, *Turnera diffusa*, *Matricaria recutita*, *Passiflora incarnata*, and *Tilia occidentalis*. *J AOAC Int* 2007, 90(4):920-924

Ramos-e-Silva M, Ferreira AF, Bibas R, Carneiro S. Clinical evaluation of fluid extract of *Chamomilla recutita* for oral aphthae. *J Drugs Dermatol* 2006, 5(7):612-617

Razmjoo K, Heydarizadhe P, Sabzalian MR. Effect of Salinity and Drought Stresses on Growth Parameters and Essential Oil Content of *Matricaria chamomilla*. *Int J Agric Biol* 2008, 10(4):451-454

Redaelli C, Formentini L, Santaniello E. HPLC Determination of Coumarins in *Matricaria chamomilla*. *Planta Med* 1981, 43(12):412-413

Redaelli C, Formentini L, Santaniello E. Reversed-Phase High-Performance Liquid Chromatography Analysis of Apigenin and its Glucosides in Flowers of *Matricaria chamomilla* and Chamomile Extracts. *Planta Med* 1981, 42(7):288-292

Reicher K. Zur Therapie der akuten und chronischen Dysenterie, Kolitis, Sigmoiditis und Proktitis. *Münch Med Wschr* 1925, 7:261-262

Reichling J, Bisson W, Becker H. Vergleichende Untersuchungen zur Bildung und Akkumulation von etherischem Öl in der intakten Pflanze und in der Calluskultur von *Matricaria chamomilla*. *Planta Med* 1984, 50(4):334-337

Repčák M, Eliašová A, Ruscancinová A. Production of herniarin by diploid and tetraploid *Chamomilla recutita*. *Pharmazie* 1998, 53(4):278-279

Repčák M, Šmajda B, Kováčik J, Eliašová A. Circadian rhythm of (Z)- and (E)-2-β-D:-glucopyranosyloxy-4-methoxy cinnamic acids and herniarin in leaves of *Matricaria chamomilla*. *Plant Cell Rep* 2009, 28(7):1137-1143

Riepelmeier F. Über das Kamillenpräparat Kamillosan und seine Brauchbarkeit in der Hals-, Nasen- und Ohrenheilkunde. *Monatsschr Ohrenheilkd Laryngorhinol* 1933, 67:483-487

Rivera IG, Martins MT, Sanchez PS, Sato MIZ, Coelho MC, Akisue M, et al. Genotoxicity assessment through the Ames test of medicinal plants commonly used in Brazil. *Environm Toxicol Water Quality* 1994, 9(2):87-93

Rodríguez-Serna M, Sánchez-Motilla JM, Ramón R, Aliaga A. Allergic and systemic contact dermatitis from *Matricaria chamomilla* tea. *Contact Dermatitis* 1998, 39(4):192-209

Rodríguez-Serna M, Sánchez-Motilla JM, Ramón R, Aliaga A. Short Communications Allergic and systemic contact dermatitis from *Matricaria chamomilla* tea. *Contact Dermatitis*. 1998 39(4):192-193

Rösch W, Vinson B, Sassin I. A randomized clinical trial comparing the efficacy of a herbal preparation STW 5 with the prokinetic drug cisapride in patients with dysmotility type of functional dyspepsia. *Z Gastroenterol* 2002, 40:401-408

Rösch W, Vinson BR, Gundermann KH, Holtmann G. Phytotherapy for functional dyspepsia. A review of the clinical evidence for the herbal preparation STW 5. *Phytomedicine* 2006, 13(Suppl 5):114-121

Ross SM. Chamomile: A Spoonful of Medicine. *Holist Nurs Pract* 2008, 22(1):56-57

Rossmann H, Patzelt-Wenzler R. Prüfung der Wirksamkeit, Verträglichkeit und Akzeptanz von Kamilllosan® Mundspray bei entzündlichen Erkrankungen in Mundhöhle und Rachenraum. In: Klaschka F, Maiwald L, Patzelt-Wenzler R, editors. Wirkungsweise und Anwendungsformen der Kamille. Grosse Verlag, Berlin 1988, 106-110

Roth L, Daunderer M, Kormann K. Giftpflanzen-Pfanzengifte, Vorkommen Wirkung Therapie. Chapter: Matricaria recutita. 2nd ed. Ecomed Verlagsgesellschaft mbH, Landsberg 1984

Roth L, Daunderer M, Kormann K. Giftpflanzen-Pfanzengifte, Vorkommen Wirkung Therapie. Chapter: Matricaria recutita. 4th ed. Ecomed Verlagsgesellschaft mbH, Landsberg 1984

Rubiolo P, Belliardo F, Cordero C, Liberto E, Sgorbini B, Bicchi C. Headspace-solid-phase microextraction fast GC in combination with principal component analysis as a tool to classify different chemotypes of chamomile flower-heads (Matricaria recutita L.). *Phytochem Anal* 2006, 17(4):217-225

Rudzki E, Rebandel P. Positive patch test with Kamilllosan® in a patient with hypersensitivity to camomile. *Contact Dermatitis* 1998, 38(3):164-184

Rudzki E, Zawisza E, Rapiejko P, Rebandel P, Jaworski E. Oral allergy syndrome with contact urticaria from cosmetic creams. *Contact Dermatitis* 1999, 40(6):326

Rücker G, Mayer R, Lee KR. Peroxide als Pflanzeninhaltsstoffe, 6. Mitt.: Hydroperoxide aus den Blütenköpfchen der Römischen Kamille, Anthemis Nobilis L. Peroxides as plant constituents. 6. Hydroperoxides from the blossoms of Roman camomile, Anthemis nobilis L. *Arch Pharm* 1989, 322(11):821-826

Sabieraj J. Zum Mechanismus der entzündungshemmenden Wirkung der Kamille, ihrer Extrakte und Inhaltsstoffe Bisaboloide, Chamazulen, Matricin, der Spiroether und Apigenin. PhD Thesis, Eberhard-Karls University Tübingen 1993

Saderi H, Owlia P, Hosseini A, Semiyari H. Antimicrobial Effects of Chamomile Extract and Essential Oil on Clinically Isolated *Porphyromonas gingivalis* from Periodontitis. *Acta Hortic* 2005, 680:145-146

Sadr Lahijani MS, Raoof Kateb HR, Heady R, Yazdani D. The effect of German chamomile (*Matricaria recutita* L.) extract and tea tree (*Melaleuca alternifolia* L.) oil used as irrigants on removal of smear layer: a scanning electron microscopy study. *Int Endod J* 2006, 39(3):190-195

Saller R, Beschorner M, Hellenbrecht D, Bühring M. Dose-dependency of symptomatic relief of complaints by camomile steam inhalation in patients with common cold. *Eur J Pharmacol* 1990, 183:728-729

Sato F, Matsukawa Y, Matsumoto K, Nishino H, Sakai T. Apigenin induces morphological differentiation and G2/M arrest in rat neuronal cells. *Biochem Biophys Res Commun* 1994, 204:578-584

Scalia S, Giuffreda L, Pallado P. Analytical and preparative supercritical fluid extraction of chamomile flowers and its comparison with conventional methods. *J Pharm Biomed Anal* 1999, 21(3):549-558

Schemann M, Michel K, Hohenester B, Rühl A. Region-specific effects of STW 5 and its components in gastric fundus, corpus and antrum. *Phytomedicine* 2006, 13(Suppl 5):90-99

Schempp H, Toth A, Weiser D, Elstner EF. Antioxidative properties of Iberis amara extracts in biochemical model reactions. *Arzneimittelforschung* 2003, 53(8):568-577

Schempp H, Weiser D, Kelber O, Elstner EF. Comparison of the inhibition of myeloperoxidase-catalyzed hypochlorite formation by different plant extracts contained in the drug Iberogast.

Arzneimittelforschung 2004, 54(7):389-95

Schempp H, Weiser D, Kelber O, Elstner EF. Radical scavenging and anti-inflammatory properties of STW 5 and its components. *Phytomedicine* 2006, 13(Suppl 5):36-44

Schilcher H. Standardisierte Kamillenzubereitungen – eine Voraussetzung für die medizinische Anwendung von Kamillenblüten. In: Klaschka F, Maiwald I, Patzelt-Wenzler R, editors. Wirkungsweise und Anwendungsformen der Kamille. Grosse Verlag, Berlin 1988, 13-18

Schilcher H. Neuere Erkenntnisse bei der Qualitätsbeurteilung von Kamillenblüten bzw. Kamillenöl, 2. Teil des Referates. 20th Vortragstagung der Gesellschaft für Arzneipflanzenforschung Helsinki, 22-29 Juli 1972, 23/2:132-144

Schilcher H. Phytopharmaka zur äußeren Anwendung in der Kinderheilkunde. Wissenschaftliche Verlagsgesellschaft, Stuttgart 1991

Schmid F. Kamillenwirkstoffe und deren Indikationen in der Kinderheilkunde aus Kamillentherapie - in der Praxis bewährt, naturwissenschaftlich begründet? „Erfahrungstherapie – späte Rechtfertigung. Internationales Symposium Wien 30-31 Mai. Karlsruhe Verlag, G. Braun 1975

Schmidt M. Die Kamille als Heilpflanze. *PTA heute* 1988, 2(10):312-315

Schmidt PC, Soyke B. Entwicklung eines matricinreichen Kamillenpräparates. *Sci Pharm* 1991, 60:111-123

Schmidt PC, Weibler K, Soyke B. Kamillenblüten -und extrakte. *Dtsch Apoth Ztg* 1991, 131(5):175-81

Schreiber A, Carle R, Reinhard E. On the Accumulation of Apigenin in Chamomile Flowers. *Planta Med* 1990, 56(2):179-181

Schröder EM. Fenchel, Kamille und Anis - Klassiker der Kräutertees. *Dtsch Apoth Ztg* 2006, 146(35):58-59

Schubert HJ. Allergie auf Asteraceae (Compositae) im Gartenaugegebiet Erfurt. *Dermatosen* 1995, 43(6):257-261

Schultze W. Composition of the essential oil of the flower heads of *Matricaria perforata*. *Planta Med* 1996, 62(4):332-325

Schulz V. Apfelpektin-Kamillenextrakt im Placebovergleich wirksam bei akuter Diarrhoe. *Z Phytother* 2006, 27:286-287

Schulz V. Chamazulencarbonsäure (CHCS) aus Kamillenblüten, *Z Phytother* 2007, 28:22-23

Schulz V. Risiko von Arzneimittelinteraktionen durch Echte Kamille (*Matricaria recutita* L.) *Z Phytother* 2009, 30:162-168

Seidler-Lozykowska K. Determination of the ploidy level in chamomile (*Chamomilla recutita* (L.) Rausch.) strains rich in alpha-bisabolol. *J Appl Genet* 2003, 44(2):151-155

Senff H, Kuhlwein A, Kalveram KJ, Hausen BM. Asthma bronchiale durch Inhalation mit echter Kamille. *Allergologie* 1989, 12(2):51-53

Shabanloei R, Ahmadi F, Vaez J, Ansarin K, Hajizadeh E, Javadzadeh Y, et al. Alloporeinol, Chamomile and Normal Saline Mouthwashes for the Prevention of Chemotherapy-Induced Stomatitis. *J Clin Diagn Res* 2009, 3:1537-1542

Shamseer L, Charrois TL, Vohra S. Chamomile: Practical management of adverse effects and drug interactions. *Clinical Brief Peer Review* 2006, 139(6):1281-1282

Shikov AN, Pozharitskaya ON, Makarov VG, Kvetnaya AS. Antibacterial Activity of *Chamomilla recutita* on Extract against *Helicobacter pylori*. *Phytother Res* 2008, 22(2):252-253

Shoukry IF. Toxicological deteriorations of two volatile oils of *Matricaria chamomilla* and *Clerodendron inerme* on the adult house fly *Musca domestica* L. *J Egypt Soc Parasitol* 1997, 27(3):893-904

Sibaev A, Yuece B, Kelber O, Weiser D, Schirra J, Göke B, et al. STW 5 and its individual herbal components modulate intestinal electrophysiology of mice. *Phytomedicine* 2006, 13(Suppl 5):80-89

Simmen U, Kelber O, Okpanyi SN, Jaeggi R, Bueter B, Weiser D. Binding of STW 5 and its components to intestinal 5-HAT, muscarinic M₃, and opioid receptors. *Phytomedicine* 2006, 13(Suppl 5):51-55

Srivastava JK, Gupta S. Antiproliferative and Apoptotic Effects of Chamomile Extract in Various Human Cancer Cells. *J Agric Food Chem* 2007, 55(23):9470-9478

Skórska C, Sitkowska J, Krysińska-Traczyk E, Cholewa G, Dutkiewicz J. Exposure to Airborne Microorganisms, Dust and Endotoxin during Processing of Peppermint and Chamomile Herbs on Farms. *Ann Agric Environ Med* 2005, 12(2):281-288

Spaich W. Moderne Phytotherapie. Chapter: Chamomilla. Karl F Haug Verlag, Heidelberg 1978

Srivastava JK, Gupta S. Antiproliferative and apoptotic effects of chamomile extract in various human cancer cells. *J Agric Food Chem* 2007, 55(23):9470-9478

Srivastava JK, Gupta S. Health benefits of chamomile. *Recent Progress in Medicinal Plants* 2010, 27: 33-53

Srivastava JK, Shankar E, Gupta S. Chamomile: a herbal medicine of the past with a bright future (review). *Mol Med Rep* 2010, 3(6):895-901

Standen MD, Myers SP. The roles of essential oils in the modulation of immune function and inflammation: survey of aromatherapy educators. *Int J Aromather* 2004, 14:150-161

Steimer P. Iberogast-Therapie in der Gastroenterologie. *Krankenhausarzt* 1983, 56:1005-1008

Sticker A. Deutsche Kamillen, Kamillosan als Hilfsmittel in der Krebsbehandlung. *Med Welt* 1932, 37:1320

Subiza J, Subiza JL, Alonso M, Hinojosa M, Garcia R, Jerez M, et al. Allergic conjunctivitis to chamomile tea. *Ann Allergy* 1990, 65:127-32

Svehliková V, Bennett RN, Mellon FA, Needs PW, Piacente S, Kroon PA, et al. Isolation, identification and stability of acylated derivatives of apigenin 7-O-glucoside from chamomile (*Chamomilla recutita* [L.] Rauschert). *Phytochemistry* 2004, 65(16):2323-2332

Swoboda M. Stark in der Wundheilung. *Pharm Rundschau* 1988, 4:26

Szőke E, Máday E, Tyihák E, Kuzokina IN, Lemberkovics E. New terpenoids in cultivated and wild chamomile (in vivo and in vitro), *J Chromatogr B Analyt Technol Biomed Life Sci* 2004, 800(1-2):231-238

Szőke E, Máday E, Kiss SA, Sonnewend L, Lemberkovics E. Effect of magnesium on essential oil formation of genetically transformed and non-transformed chamomile cultures. *J Am Coll Nutr* 2004, 23(6):763S-767S

Szőke E, Máday E, Gershenson J, Allen JL, Lemberkovics E. β -eudesmol, a new sesquiterpene component in intact and organized root of chamomile (*Chamomilla recutita*). *J Chromatogr Sci* 2004, 42(5):229-233

Tamasdan S, Cristea E, Mihele D. The action upon gastric secretion of robiniae flores, chamomillae flores and strobuli lupuli extracts. *Farmacia* 1981, 29(2):71-75

Thiemer K, Stadler R, Isaac O. Biochemische Untersuchungen von Kamillen-Inhaltsstoffen II. *Arzneimittelforschung* 1973, 23:756-759

Tissot H. Über Kamillosan und Kamillosept. *Schweiz Med Wschr* 1929, 39:992-993

Torrado S, Torrado S, Agis A, Jimenez ME, Cadorniga R. Effect of dissolution profile and (-)- alpha-Bisabolol on the gastrototoxicity of acetylsalicylic acid. *Pharmazie* 1995, 50(2):141-143

Tschiersch K, Hözl J. [Absorption and excretion of apigenin, apigenin-7-glycoside and herniarin after oral administration of extracts of Matricaria recutita (L.) (syn. Chamomilla recutita (L.) Rauschert).] *Pharmazie* 1993, 48(7):554-555 [German]

Uda Y, Price KR, Williamson G, Rhodes MJ. Induction of the anticarcinogenic marker enzyme, quinone reductase, in murine hepatoma cells in vitro by flavonoids. *Cancer Lett* 1997, 120(2):213-216

Vanderplas O, Pirson F, D'Alapos V, Vander-Borgh T, Thimpong J, Filette C. Occupational asthma caused by chamomile. *Allergy* 2008, 63(8):1090-1092

Vogelsang-Tschiersch K. Apigenin, Apigenin-7- glucosid und Acrylderivate aus *Matricaria recutita* L.: Analytik, ^{14}C - Markierung und Beiträge zur Pharmakokinetik. PhD Thesis, Philipps University Marburg 1994

von Arnim U, Peitz U, Vinson B, Gundermann KJ, Malfertheiner P. STW 5, a phytopharmacon for patients with functional dyspepsia. Results of a multicenter, placebo-controlled double-blind study. *Am J Gastroenterol* 2007, 102(6):1268-1275

Wagner C, Marquard R, Friedt W, Ordon F. [Investigations on the genetic diversity of camomile (*Chamomilla recutita* (L.) Rausch.) by PCR-based marker techniques.] *Z Arzn Gew Pfl* 2001, 6/4:216-221 [German]

Wagner H. Pharmazeutische Biologie, Drogen und ihre Inhaltsstoffe. Chapter: Chamomillae (Matricariae) Flos. 2nd ed. Gustav Fischer Verlag, Stuttgart 1980

Wagner H. Multitarget therapy – the future of treatment for more than just functional dyspepsia. *Phytomedicine* 2006, 13(Suppl 5):122-29

Wagner H, Wiesenauer M. Phytotherapie: Phytopharmaka und pflanzliche Homöopathika. Gustav-Fischer Verlag, Stuttgart Jena New York 1995

Wang Y, Tang H, Nicholson JK, Hylands PJ, Sampson J, Whitcombe I, et al. Metabolomic strategy for the classification and quality control of phytomedicine: A case study of chamomile flower (*Matricaria recutita* L.). *Planta Med* 2004, 70(3):250-255

Wang Y, Tang H, Nicholson JK, Hylands PJ, Samson J, Holmes E. A metabonomic strategy for the detection of the metabolic effects of chamomile (*Matricaria recutita* L.) ingestion. *J Agric Food Chem* 2005, 53(2):191-196

Weber-Stahl K. Lehrbuch der Pharmakognosie. Chapter 3: Flores Chamomillae - Echte kamillenblüten. 9th ed. Gustav Fischer Verlag, Stuttgart 1962

- Wegener T, Wagner H. The active components and the pharmacological multi-target principle of STW 5 (Iberogast). *Phytomedicine* 2006, 13(Suppl 5):20-35
- Wehba C, Fernandes F, Oppi EC. Application of the ointment of chamomile extract as adjunctive in reducing symptoms of the painful, ulcerated lesions of the oral mucosa. *RBM* 2008, 129-132
- Weinsheimer K. Über Kamillosan-Salbe. *Klin Wschr* 1923, 2(5) Reprint
- Weiβ RF. Lehrbuch der Phytotherapie. Chapter: 1.1 Matricaria chamomilla. 6th ed. Hippokrates Verlag, Stuttgart 1985
- Weiβ RF. Moderne Pflanzenheilkunde. Neues über Heilpflanzen und ihre Anwendung in der Medizin. 7th ed. Kneipp-Verlag, Bad Worishofen 1982
- Weiβ RF, Fintelmann V. Lehrbuch der Phytotherapie. Matricaria chamomilla, Echte Kamille. 8th ed. Hippokrates Verlag, Stuttgart 1997
- Weiβ, RF. Kamille- "Heilpflanze 1987". Kneipp-Blatter 1987, 1:4-8
- Weitgasser H. Dermatologische Erfahrungen mit Kamillosan. *ZFA* 1979, 5:340-352
- Wichtl M. Kapitel. Kamillenblüten. In: Teedrogen, Wissenschaftliche Verlagsgesellschaft, Stuttgart
- Wichtl M, editor. Teedrogen und Phytopharmaka. 4th ed. Wissenschaftliche Verlagsgesellschaft, Stuttgart 2002
- Wiedemann B. [Gluteal pruritus... Not necessarily a mycotic etiology and especially don't recommend camomile baths!] *MMW Fortschr Med* 2001, 143(37):6-8 [German]
- Williamson EM. Interactions between herbal and conventional medicines: the role of cytochrome P450 and P-Glycoprotein. *Pharmacologyonline* 2006, 2:200-205
- Wölle J, Hill RR, Ferguson E, Devall LJ, Trivedi BK, Newton RS, et al. Selective inhibition of tumor necrosis factor-induced vascular cell adhesion molecule-1 gene expression by a novel flavonoid. Lack of effect on transcription factor NF-kappa B. *Arterioscler Thromb Vasc Biol* 1996, 16(12):1501-1508
- Woollard AC, Tatham KC, Barker S. The influence of essential oils on the process of wound healing: a review of the current evidence. *J Wound Care* 2007, 16(6):255-257
- Wunderer H. Mund- und Rachentherapeutika. *Dtsch Apoth Ztg* 1986, 126(42):2281-2292
- Yoshinari T, Yaguchi A, Takahashi-Ando N, Kimura M, Takahashi H, Nakajima T, et al. Spiroethers of German chamomile inhibit production of aflatoxin C1 and trichothecene mycotoxin by inhibiting cytochrome P450 monooxygenases involved in their biosynthesis. *FEMS Microbiol Lett* 2008, 284(2):184-190
- Zamani AA, Talebi A, Fathipour Y, Baniameri V. Effect of temperature on life history of *Aphidius colemani* and *Aphidius matricariae* (Hymenoptera: Braconidae), two parasitoids of *Aphis gossypii* and *Myzus persicae* (Homoptera: Aphididae). *Environ Entomol* 2007, 36(2):263-271
- Zander R. Handwörterbuch der Pflanzennamen. Chapter: Chamomilla recutita. Eugen Ulmer Verlag, Stuttgart 1979
- Zepernick B, Langhammer L, Lüdke JBP. Lexikon der offizinellen Arzneipflanzen. Chapter: Chamomilla. Walter de Gruyter Verlag, Berlin 1984
- Zwaving JH. Echte en roomse kamille, hun verschil en overeenkomst in samenstelling en toepassing. *Pharm Weekbl* 1982, 117:157-165