

12 November 2013
EMEA/HMPC/321182/2012
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Pimpinella anisum* L., fructus and *Pimpinella anisum* L., aetheroleum

Final

The Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Abraham SK. Anti-genotoxicity of trans-anethole and eugenol in mice. Food Chem Toxicol 2001; 39: 493-498.

Aburjai T, Darwish RM, Al-Khalil S, Mahafzah A, Al-Abbad A. Screening of antibiotic resistant inhibitors from local plant materials against two different strains of *Pseudomonas aeruginosa*. J Ethnopharmacol 2001; 76: 39-44.

Al-Bataina BA, Maslat AO, Al-Kofahi MM. Element analysis and biological studies on ten oriental spices using XRF and Ames test, JTrace Elem Med Biol 2003; 17: 85-90.

Albert-Puleo MJ. Fennel and anise as estrogenic agents. J Ethnopharmacol 1980; 2: 337-344.

Al-Harbi MM, Qureshi S, Raza M, Ahmed MM, Giangreco AB, Shah AH. Influence of anethole treatment on the tumour induced by Ehrlich ascites carcinoma cells in paw of Swiss albino mice. Eur J Cancer Prev 1995; 4: 307-318.

Andersen KE, Contact allergy to toothpaste flavours. Contact Dermatitis 1978; 4: 195-198.

Arslan N, Gürbüz B, Saruhan EO, Bayrak A, Gümüşçü A. Variation in essential oil content and composition in Turkish anise (*Pimpinella anisum* L.) populations. Turkish J Agricul Forestry 2004; 28: 173-177.



Bellakhdar J, Claisse R, Fleurentin J, Younos C. Repertory of standard herbal drugs in the Moroccan Pharmacopoeia. *J Ethnopharmacol* 1991; 35(2): 123-143.

Blumenthal M, Busse WR, Goldberg A, et al. editors. The Complete German Commission E Monographs: Therapeutic Guide to Herbal Medicines. Austin: American Botanical Council and Boston: Integrative Medicine Communications 1998, 82-83.

Boskabady MH, Ramazani-Assari M. Relaxant effect of *Pimpinella anisum* on isolated guinea pig tracheal chains and its possible mechanism(s). *J Ethnopharmacol* 2001, 74: 83-88.

Bounds SV, Caldwell J. Pathways of Metabolism of [1'-14C] trans-anethole in the rat and mouse. *Drug Metab Dispos* 1996; 24: 717-724.

Boyd EM. Expectorants and respiratory tract fluid. *Pharmacol Rev* 1954, 6: 521-452.

Boyd EM, Pearson GL. On the expectorant action of volatile oils. *Am J Med Sci* 1946; 211: 602-610.

Boyd EM, Sheppard EP. The effect of steam inhalation of volatile oils on the output and composition of respiratory tract fluid. *J Pharmacol Exp Therap* 1968, 163: 250-256.

British Herbal Pharmacopoeia. Pimpinella. British Herbal Medicine Association, Bournemouth 1983, 160-161.

Buechi S, Vogelin R, von Eiff MM, Ramos M, Melzer J. Open trial to assess aspects of safety and efficacy of a combined herbal cough syrup with ivy and thyme. *Forsch Komplementarmed Klass Naturheilkd* 2005, 12: 328-332.

Burkhardt G, Reichling J, Martin R, Becker H. Terpene hydrocarbons in *Pimpinella anisum* L. *Pharm Weekbl Sci* 1986, 8: 190-193.

Caldwell J, Chan VSW, Marshall AD, Hasheminejad G, Bounds SJV. 1'-hydroxylation is the only metabolic pathway of simple alkylbenzenes involved in their genotoxicity. *Toxicologist* 1992, 12: 56.

Caldwell J, Sutton JD. Influence of dose size on the disposition of trans-[methoxy-14C] anethole in human volunteers. *Food Chem Toxicol* 1988, 26: 87-91.

Caldwell J. Perspective on the usefulness of the mouse lymphoma assay as an indicator of a genotoxic carcinogen: ten compounds which are positive in the test but are not genotoxic carcinogens. *Teratogen Carcin Mut* 1993, 13: 185-190.

Cambar PJ, Aviado DM. Bronchopulmonary effects of paraquat and expectorants. *Arch Env Health* 1970; 20: 488-494.

Ceska O, Chaudhary SK, Warrington PJ, Ashwood-Smith MJ. Photoactive furocoumarins in fruits of some umbellifers. *Phytochemistry* 1987, 26: 165-169.

Chainy GBN, Manna SK, Chaturvedi MM, Aggarwal BB. Anethole blocks both early and late cellular responses transduced by tumor necrosis factor: effect on NF-KB, AP-1, JNK, MAPKK and apoptosis. *Oncogene* 2000, 19: 2943-2950.

Czygan F-C, Hiller K. Anisi fructus - Anis. In: Wichtl M, editor. Teedrogen and Phytopharma. Ein Handbuch für die Praxis auf wissenschaftlicher Grundlage, 4th ed. Stuttgart: Wissenschaftliche Verlagsgesellschaft 2002, 42-44.

Czygan F-C. Anisi fructus DAB 10 - *Pimpinella anisum* L. Z Phytotherapie 1992, 13: 101-106.

Dhar SK. Anti-fertility activity and hormonal profile of trans-anethole in rats. Indian J Physiol Pharmacol 1995, 39: 63-67.

Dirks U, Herrmann K. 4-(13-D-glucopyranosyloxy)-benzoic acid, a characteristic phenolic constituent of the Apiaceae. Phytochemistry 1984, 23: 1811-1812.

Dirks U, Herrmann K. Hochleistungsflüssigkeits chromatographie der Hydroxycinnamoylchinasäuren und der 4-O-D-glucopyranosyloxy)-benzoësäure in Gewürzen. Z Lebensm Unters Forsch 1984; 179: 12-16.

Dorsch W, Loew D, Meyer-Buchtela E, Schilcher H. Teil I. Empfehlungen zur Anwendung und Dosierung von Phytopharma, monographierte Arzneidrogen und ihren Zubereitungen in der Pädiatrie: Foeniculi fructus (Fenchelfrüchte) In: Kooperation Phytopharma, editor. Kinderdosierung von Phytopharma, 3rd ed. Kooperation Phytopharma Bonn 2002: 70-71.

Drinkwater NR, Miller EC, Miller JA, Pitot HC. Hepatocarcinogenicity of estragole (1-allyl-4-methoxybenzene) and 1'-hydroxyestragole in the mouse and mutagenicity of 1'-acetoxyestragole in bacteria. J Natl Cancer Inst 1976, 57: 1323-1331.

EI-Moghazi AM, Ali AA, Ross SA, Mottaleb MA. Flavonoids of *Pimpinella anisum* L. growing in Egypt. Fitoterapia 1979, 50: 267-268.

EI-Wakeil F, Khairy M, Morsi S, Farag RS, Shihata AA, Badel ZMA. Biochemical studies on the essential oils of somefruits of Umbelliferaefamily. Seifen-Ole-Fette-Wachse 1986, 112: 77-80.

Elgayyar M, Draughon FA, Golden DA, Mount JR. Antimicrobial activity of essential oils from plants against selected pathogenic and saprophytic microorganisms. J Food Prot 2001, 64: 1019-1024.

EMEA, HMPC: Committee for Herbal Medicinal Products: 'Public statement on the use of herbal medicinal products containing estragole' (EMEA/HMPC/137212/2005).

EMEA, CVMP: Committee for Veterinary Medicinal Products: Anisi aetheroleum, summary report. (EMEA/MRL/413/98-FINAL)

ESCOP (.) Monographs: Aniseed. European Scientific Cooperative on Phytotherapy, editors. Argyle House, Exeter, UK 1996-1999.

ESCOP Monographs : Aniseed. European Scientific Cooperative on Phytotherapy.. Argyle House, ESCOP, Exeter, UK 2003.

European Commission: Scientific Committee on Food. Opinion of the Scientific Committee on food on estragole (1-allyl-4-methoxybenzene). 26 September 2001.

European Commission Report from the Commission on Dietary Food Additive Intake in the European Union. COM/2001/0542 Final.

European Pharmacopoeia 5th ed., Anise oil - Anisi aetheroleum. Council of Europe. 1/2005: 0804; 1004.

European Pharmacopoeia 4th ed. Aniseed - Anisi fructus. Council of Europe. 1997

European Pharmacopoeia 5th ed., Aniseed - Anisi fructus. Council of Europe. 1/2005: 0262; 1006.

European Pharmacopoeia 7th ed., Anise oil - Anisi aetheroleum. Council of Europe. 1/2008: 0804 corrected 7; 1050-1052.

European Pharmacopoeia 7th ed., Aniseed - Anisi fructus. Council of Europe. 1/2012: 0262; 3858-3859.

Farag R, Daw Z, Abo-Raya S. Influence of some spice essential oils on *A. parasiticus* growth and production of aflatoxins in a synthetic medium. J Food Sci 1989, 54: 74-76.

Fraj J, Lezaun A, Colas C, Duce F, Dominguez MA, Alonso MD. Occupational asthma induced by aniseed. Allergy 1996, 51(5): 337-339.

Franks A. Contact allergy to anethole in toothpaste associated with loss of taste. Contact Dermatitis 1998, 38: 354-355.

Friedman M, Henika PR, Mandrell RR. Bactericidal activities of plant essential oils and some of their isolated constituents against *Campylobacter jejuni*, *Escherichia coli*, *Listeria monocytogenes* and, *Salmonella enterica*. J Food Protect 2002, 65: 1545-1560.

Fujimatu E, Ishikawa T, Kitajima J. Aromatic compound glucosides, alkyl glucoside and glucide from the fruit of anise. Phytochemistry 2003, 63: 609-616.

Fyfe L, Armstrong F, Stewart J. Inhibition of *Listeria monocytogenes* and *Salmonella enteriditis* by combinations of plant oils and derivatives of benzoic acid: the development of synergistic antimicrobial combinations. Int J Antimicrobial Agents 1998, 9: 195-199.

García-Bravo B, Pérez Bernal A, García-Hernández MJ, Camacho F. Occupational contact dermatitis from anethole in food handlers. Contact Dermatitis 1997, 37: 38.

García-González JJ, Bartolomé-Zavala B, Fernández-Meléndez S, Barceló-Muñoz JM, Miranda Páez A, Carmona-Bueno MJ et al. Occupational rhinoconjunctivitis and food allergy because of aniseed sensitization. Ann Allergy Asthma Immunol 2002, 88: 518-522.

Gázquez García V, Gaig Jané P, Bartolomé Zavala B. Aniseed-induced nocturnal tongue angioedema. J Investig Allergol Clin Immunol 2007, 17: 406-408.

Gershbein LL. Regeneration of rat liver in the presence of essential oils and their components. Food Cosmet Toxicol 1977, 15: 173-181.

Ghelardini C, Galeotti N, Mazzanti G. Local anaesthetic activity of monoterpenes and phenylpropanes of essential oils. *Planta Med* 2001, 67: 564-566.

Gorelik NJ. Genotoxicity of trans-anethole *in vitro*. *Mutation Res* 1995, 326: 199-209.

Gülçin I, Oktay M, Kireçci E, İrfan Küfrevoğlu Ö. Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extracts. *Food Chemistry* 2003, 83: 371-382.

Guneyli A, Karacali I. Effects of storage conditions on the quality of aniseed. *Ege Universitesi Ziraat Fakultesi Dergisi* 2002, 39: 17-24.

Hagan EC, Hansen WH, Fitzhugh OG, Jenner PM, Jones WI, Taylor JM et al. Food flavourings and compounds of related structure. II. Subacute and chronic toxicity. *Food Cosmet Toxicol* 1967, 5: 141-157.

Haggag EG, Abou-Moustafa MA, Boucher W, Theoharides TC. The effect of a herbal water-extract on histamine release from mast cells and on allergic asthma. *J Herb Pharmacother* 2003, 3: 41-54.

Hammer KA, Carson CF, Riley TV. Antimicrobial activity of essential oils and other plant extracts. *J Applied Microbiol* 1999, 86: 985-990.

Hänsel R, Keller K, Rimpler H, Schneider G, editors. *Pimpinella*. In: *Hagers Handbuch der Pharmazeutischen Praxis*, Vol 6.5th ed.: Drogen P-Z. Springer-Verlag, Berlin-Heidelberg-New York-London, 1994, 135-156.

Heck AM, DeWitt BA, Lukes AL: Potential interactions between alternative therapies and warfarin. *Am J Health Syst Pharm* 2000, 57: 1221-1227.

Heck JD, Vollmuth TA, Cifone MA, Jagannath DR, Mhyr B. and Current RD. An evaluation of food flavouring ingredients in a genetic toxicity screening battery. *Toxicologist* 1989, 9: 257.

Howes AJ, Chan VSW, Caldwell J. Structure-specificity of the genotoxicity of some naturally occurring alkenylbenzenes determined by the unscheduled DNA synthesis assay in rat hepatocytes. *Food Chemical Toxicol* 1990, 28: 537-542.

Howes MJ, Houghton PJ, Barlow DJ, Pocock VJ, Milligan SR. Assessment of estrogenic activity in some common essential oil constituents. *J Pharm Pharmacol* 2002, 54: 1521-1528.

Hsia MTS, Adamovics JA, Kremer BL. Microbial mutagenicity studies of insect growth regulators and other potential insecticidal compounds in *Salmonella typhimurium*. *Chemosphere* 1979, 8: 521-529.

Ibrahim YKE, Ogunmodede MS. Growth and survival of *Pseudomonas aeruginosa* in some aromatic waters. *Pharm Acta Helv* 1991, 66: 286-288.

Ishidate M, Sofuni T, Yoshikawa K, Hayashi M, Nohmi T, Sawada M et al. Primary mutagenicity screening of food additives currently used in Japan. *Food Chem Toxicol* 1984, 22: 623-636.

Ishikawa T, Fujimatu E, Kitajima J. Water-soluble components of anise: new glucosides of anethol glycol and its related compounds. *Chem Pharm Bull* 2002; 50: 1460-1466.

Janahmadi M, Farajnia S, Vatanparast J, Abbasipour H, Kamalinejad M. The fruit essential oil of *Pimpinella anisum* L. (Umbelliferae) induces neuronal hyperexcitability in snail partly through attenuation of after-hyperpolarization. *Journal of Ethnopharmacology* 2008; 120: 360-365.

Jensen-Jarolim E, Leitner A, Hirschwehr R, Kraft D, Wuthrich B, Scheiner O, et al. Characterization of allergens in Apiaceae spices: anise, fennel, coriander and cumin. *Clin Exp Allergy* 1997; 27: 1299-1306.

JECFA :Joint FAO/WHO Expert Committee on Food Additives .Safety evaluation of certain food additives. Trans-anethole. WHO Geneva, 1999.

Kassi E, Papoutsi Z, Fokialakis N, Messari, J., Mitakou S., Moutsatsou P: Greek plant extracts exhibit selective estrogen receptor modulator (SERM)-like properties. *J Agric Food Chem* 2004; 52: 6956-6961.

Kitajima J, Ishikawa T, Fujimatu E, Kondho K, Takayanagi, T. Glycosides of 2-C-methyl-D-erythritol from the fruits of anise, coriander and cumin. *Phytochemistry* 2003; 62: 115-120.

Kreydiyyeh SI, Usta J, Knio K, Markossian S, Dagher S. Aniseed oil increases glucose absorption and reduces urine output in the rat. *Life Sci* 2003; 74: 663-673.

Kreydiyyeh SI, Usta J, Copti R. Effects of cinnamon, clove and some of their constituents on the Na(+) - K(+) - ATPase activity and alanine absorption in the rat jejunum. *Food ChemToxicol* 2000; 38: 755-762.

Kubeczka KH. Grundlagen der Qualitätsbeurteilung arzneilich verwendeter atherischer Ole. *Acta Horticulturae* 1978; 73: 85-93.

Kunzemann J, Herrmann K. Isolierung und Identifizierung der Flavon(ol)-O-glykoside in Kiimmel (*Carum carvi* L.), Fenchel (*Foeniculum vulgare* Mill.), Anis (*Pimpinella anisum* L.) und Koriander (*Coriandrum sativum* L.) und von Flavon-C-Glykosiden im Anis. *Z Lebensm Unters Forsch* 1977; 164: 194-200.

Le Bourhis B. Propriétés biologiques du trans-anethole. Essai de détermination de la dose journalière acceptable. *Parfums Cosmet Sav Fr* 1973, 3: 450-456.

Leclerc H. Anis. In: *Précis de Phytothérapie. Essai de thérapeutique par les plantes françaises*. 5 ed. Paris, Masson 1983.

Lee JB, Yamagishi C, Hayashi K and Hayashi T. Antiviral and immunostimulating effects of lignin-carbohydrate protein complexes from *Pimpinella anisum*. *Biosci Biotechnol Biochem* 2011; 75(3): 459-465.

Lin FSD. Trans-anethole. In: Joint FAO/WHO Expert Committee on Food Additives. Toxicological evaluation of certain food additives and contaminants. *WHO Food Additives Series 28*. WHO Geneva, 1991: 135-152.

Linares E, Bye RA. A study of four medicinal plant complexes of Mexico and adjacent United States. *J Ethnopharmacol* 1987, 19: 153-183.

Marcus C, Lichtenstein EP. Interactions of naturally occurring food plant components with insecticides and pentobarbital in rats and mice. *J Agric Food Chem* 1982, 30: 563-568.

Marshall AD, Caldwell J. Lack of influence of modulators of epoxide metabolism on the genotoxicity of trans-anethole in freshly isolated rat hepatocytes assessed with the unscheduled DNA synthesis assay. *Food Chem Toxicol* 1996, 34(4):337-45.

Martindale. The complete drug reference, 33rd ed. Aniseed, Anise oil. Sweetmann SE. Pharmaceutical Press, London 2002, 1580.

Maruzzella JC, Freundlich M. Antimicrobial substances from seeds. *J Am Pharm Assoc* 1959, 48: 356-358.

Miller EC, Swanson AB, Phillips DH, Fletcher L, Liem A, Miller JA. Structure-activity studies of the carcinogenicities in the mouse and rat of some naturally occurring and synthetic alkenylbenzene derivatives related to safrole and estragole. *Cancer Res* 1983, 43: 1124-1134.

Mortelmans K, Haworth S, Lawlor T, Speck W., Tainer B, Zeiger E. Salmonella mutagenicity tests: II. Results from the testing of 270 chemicals. *Envirnm Mutagenensis* 1986; 6 (7): 1-119.

Moyler DA. Spices-Recent Advances. In Spices, Herbs and Edible Fungi; Charalambous G. (Ed). Elsevier Science; London, UK. 1994, 1-70.

Müller-Riebau F, Berger B, Yeken O. Chemical composition and fungitoxic properties to phytopathogenic fungi of essential oils of selected aromatic plants growing wild in Turkey. *J Agric Food Chem* 1955, 43: 2262-2266.

Müller L, Kasper P, Müller-Tegethoff K, Petr T. The genotoxic potential in vitro and in vivo of the alkylbenzene etheric oils estragole, basil oil and trans-anethole. *Mutation Res* 1994; 325: 129-136.

Müller-Limmroth W, Fröhlich HH. Wirkungsnachweis einiger phytotherapeutischer Expektorantien auf den mukoziliaren transport. *Fortschr Med* 1980, 98: 95-101.

Nestmann ER, Lee EGH, Matula TI, Douglas GR, Meuller JC. Mutagenicity of constituents of pulp and paper mill effluent using the salmonella/mammalian-microsome assay. *Mutation Res* 1980, 79: 203-212.

Nestmann ER, Lee EGH. Mutagenicity of constituents of pulp and paper mill effluent in growing cells of *Saccharomyces cerevisiae* *Mutation Res* 1983, 119: 273-280.

Newall CA, Anderson LA, Phillipson JD: Herbal Medicines. A Guide for Health Care Professionals. The Pharmaceutical Press, London, England, 1996.

Newberne P, Smith RL, Doull J, Goodman JI, Munro IC, Portoghese PS *et al.* The FEMA GRAS assessment of trans-anethole used as a flavouring substance. *Food Chem Toxicol* 1999, 37: 789-811.

Newberne PM, Cartlton WW, Brown WR. Histopathological evaluation of proliferative lesions in rats fed trans-anethole in chronic studies. *Food Chem Toxicol* 1989, 27: 21-26.

Okazaki K, Nakayama S, Kawazoe K, Takaishi Y. Antiaggregant effect of human platelets of culinary herbs. *Phytother Res* 1998, 12: 603-605.

Omidbaigi R, Hadjiakhoondi A, Saharkhiz M. Changes in the content and chemical composition of *Pimpinella anisum* oil at various harvest time. *J. Essent Oil-bear Plants* 2003, 6: 46-50.(Abstract)

Ondarza M, Sanchez A. Steam distillation and supercritical fluid extraction of some Mexican spices. *Chromatographia* 1990, 30: 16-18.

Phillips DH, Reddy MV, Randerath K. 32-P-post labelling analysis of DNA adducts formed in the livers of animals treated with safrole, estragole and other naturally-occurring alkenylbenzenes.II. Newborn male B6C3F1 mice. *Carcinogenesis* 1984, 5: 1623-1628.

Picon PD, Picon RV, Costa AF, Sander GB, Amaral KM, Aboy AL et al. Randomized ized clinical trial of a phytotherapeutic compound containing *Pimpinella anisum*, *Foeniculum vulgare*, *Sambucus nigra*, and *Cassia angustifolia* for chronic constipation. *BMC Complement Altern Med* 2010, 10: 17.

Pourgholami MH, Majzoob S, Javadi M, Kamalinejad M, Fanaee GH, Sayyah M. The fruit essential oil of *Pimpinella anisum* exerts anticonvulsant effects in mice. *J Ethnopharmacol* 1999, 66: 211-215.

Ramadan FM, El-Zanfaly RT, El-Waked FA, Alian AM. On the antibacterial effects of some essential oils. I. Use of agar diffusion method. *Chem Mikrobiol Technol Lebensm* 1972, 2: 51-55.

Reed PM, Caldwell J. Induction of hepatic cytochrome P450 and related activities following dietary administration of trans-anethole in SD-CD rats. *Hum. Experimen Toxicol* 1992, 11: 580-581.

Reiter M, Brandt W. Relaxant effects on tracheal and ileal smooth muscles of the guinea pig. *Arzneim-Forsch. Drug Res* 1985, 35: 408-414.

Rodrigues VM, Paulo T, Rosa V, Marques MMO, Petenate A, Meireles MAA. Supercritical extraction of essential oil from aniseed (*Pimpinella anisum* L.) using CO₂: solubility, kinetics and composition data. *J Agric Food Chem* 2003, 51: 1518-1523.

Sahraei H, Ghoshooni H, Hossein SS, Mohseni AA, Shafaghi B, Falahi M et al.. The effects of fruit essential oil of the *Pimpinella anisum* on the acquisition and expression of morphine induced conditioned place preference in mice. *J Ethnopharmacol* 2002, 80: 43-47.

Samojlik I, Mijatović V, Petković S, Skrbić B, Božin B. The influence of essential oil of aniseed (*Pimpinella anisum*, L.) on drug effects on the central nervous system. *Fitoterapia* 2012.

Sangster SA, Caldwell J, Hutt AJ, Anthony A, Smith RL. The metabolic disposition of [methoxy-14C]-labelled trans-anethole, estragole and p-propylanisole in human volunteers. *Xenobiotica* 1987; 17: 1223-1232.

Sangster SA, Caldwell J, Smith RL, Farmer PB. Metabolism of anethole. I. Pathways of metabolism in the rat and mouse. *Food Chem Toxicol* 1984b, 22: 695-706.

Sangster SA, Caldwell J, Smith RL. Metabolism of anethole II. Influence of dose size on the route of metabolism of trans-anethole in the rat and mouse. *Food Chem Toxicol* 1984a, 22: 707-713.

Sathiyamoorthy P, Lugasi Evgil H, Schlesinger P, Kedar I, Gopas J, Pollack Y et al.. Screening for cytotoxic and antimalarial activities in desert plants of the Negev and Bedouin market plant products. Pharm Biol 1999, 37: 188-195.

Schultze W, Lange G, Kubeczka K-H. Direkte massenspektrometrische analyse von *Pimpinella anisum* L. - Anispulver and Anisi 1. Dtsch Apoth Ztg 1987, 127: 372-378.

Schulz JM, Herrmann K. Analysis of hydroxybenzoic and hydroxycinnamic acids in plant material. II. Determination by gas-liquid chromatography. J Chromatogr 1980, 195: 95-104.

Schulz V, Hansel R, Tyler VE. Rational Phytotherapy. A Physician's Guide to Herbal Medicine. Springer, New York; 1998: 159-160.

Sekizawa J, Shibamoto T. Genotoxicity of safrolerelated chemicals in microbial test systems. Mutation Res 1982, 101: 127-140.

Shashikanth KN, Hosono A. *In vitro* mutagenicity of tropical spices to streptomycin-dependent strains of *Salmonella typhimurium* TA 98. Agric Biol Chem 1986, 50: 2947-2948.

Shojaei A, Fard MA. Review of Pharmacological Properties and Chemical Constituents of *Pimpinella anisum*. ISRN Pharm 2012, 2012: 510795.

Shukla HS, Tripathi SC. Antifungal substance in the essential oil of anise (*Pimpinella anisum* L.). Agric Biol Chem 1987; 51: 1991-1993.

Singh G, Kapoor IPS, Pandey SK, Singh UK, Singh RK. Studies on essential oil: Part 10; Antibacterial activity of volatile oils of some spices. Phytother Res 2002, 16: 680-682.

Sokmen A, Jones BM, Erkork M. The in vitro antibacterial activity of Turkish Medicinal Plants. J Ethnopharmacol 1999, 67: 79-86.

Soliman KM, Badeaa RI. Effect of oil extracted from some medicinal plants on different mycotoxigenic fungi. Food Chem Toxicol 2002, 40: 1669-1675.

Steinberger E, Hänsel R. Anise and Anise oil. In: Lehrbuch der Pharmakognosie und Phytopharmazie, 4th ed. Springer-Verlag, Berlin-Heidelberg 1988, 304-306.

Swanson AB, Chambliss DD, Blomquist JC, Miller EC, Miller JA. The mutagenicities of safrole, estragole, eugenol, trans-anethole and some of their known or possible metabolites for *Salmonella typhimurium* mutants. Mutation Res 1979, 60: 143-145.

Teissedre PL, Waterhouse AL. Inhibition of oxidation of human Low-Density Lipoproteins by phenolic substances in different essential oil varieties. J. Agric. Food Chem 2000, 48: 3801-3805.

To LP, Hunt TP, Andersen ME. Mutagenicity of trans-anethole, estragole, eugenol and safrole in the Ames *Salmonella typhimurium* assay. Bull Environ Contam Toxicol 1982, 28: 647-654.

Oil of anise. HSDB Database.National Library of Medicine. Oil of anise. Available at:
(<http://toxnet.nlm.nih.gov/cgi-bin/sis/search/a?dbs+hsdb:@term+@DOCNO+1938>)

Truhaut R, Le Bourhis B, Attia M, Glomot R, Newman J, Calwell J. Chronic toxicity/carcinogenicity of trans-anethole in rats. *Food Chem Toxicol* 1989, 27; 11-19.

Tuckler V, Peck C, Nesbit C, Coleman M, Weimer S, Martinez J et al.. Seizure in an infant from aniseed oil toxicity. *J Toxicol Clin Toxicol* 2002, 40: 689.

Tunc I, Berger B.M., Erler F., Dagli F. Ovicultural activity of essential oil from five plants against two stored-product insects. *J Stored Prod Res* 2000, 36:161-168.

Valnet J. Anis. In: Aromathérapie. Traitement des maladies par les essences de plantes. 11 ed., Paris, Maloine, 1990.

Van Dongen K, Leusink H. The action of opium-alkaloids and expectorants on the ciliary movements in the air passages. *Arch Int Pharmacodyn* 1953, 93: 261-276.

Van Loon J. Die Zusammensetzung von einigen Umbelliferenölen. *Z Lebensm Unters Forsch* 1973, 153; 289-293.

Von Skramlik E. Über die Giftigkeit und Verträglichkeit von ätherischen Ölen. *Pharmazie* 1959, 14: 435-445.

Weiss RF. Herbal Medicine. Gothenberg, Sweden: Ab Arcanum and Beaconsfield: Beaconsfield Publishers Ltd, 1985: 203-204.

Weiss RF. In: Lehrbuch der Phytotherapie. 8th ed. Stuttgart: Hippocrates, 1997: 83 and 443.

Wichtl M. Herbal drugs and phytopharmaceuticals, Bisset NG editor., CRC Press. Stuttgart 1994, ,73–75.Wiseman RW, Miller EC, Miller JA, Liem A. Structure-activity studies of the hepatocarcinogenicities of alkenylbenzene derivatives related to estragole and safrole on administration to preweaning male C57BL/6J x C3H/HeJ F1 mice. *Cancer Res* 1987, 47: 2275-2283.

Wüthrich B, Dietschi R. The celery carrot mugwort condiment syndrome: skin test and RAST results. *Schweiz Med Wochenschr* 1985, 115: 258-264.

Other consulted bibliographic references, not cited in the AR.

Abdul-Ghani AS et al. Anticonvulsant effects of some Arab medicinal plants. *Int J Crude Drug Res* 1987, 25: 39-43.

Anisi fructus Monographie DAB 1996.

Anisi fructus Monographie DAB10 Kommentar Grundlfg. 1991, 2 Lfg. 1993.

Boyd EM, A review of studies on the pharmacology of the expectorants and inhalants. *Int J Clin Pharmacol* 1970, 3:55–60.

Boyd EM, Sheppard EP. Inhaled anisaldehyde and respiratory tract fluid. *Pharmacology* 1970, 3:345–352.

Boyd EM, Sheppard EP. An autumn-enhanced mucotropic action of inhaled terpenes and related volatile agents. *Pharmacology* 1971, 6(2): 65–80.

Caldwell J. Anise. In: Tisserand R, Balacs T: *Essential Oil Safety. A Guide for Health Care Professionals.* Edinburgh: Churchill Livingstone, 1995; 117.

De Smet PAGM. Legislative outlook on the safety of herbal remedies: *Pimpinella anisum*. In: De Smet PAGM, Keller K, Hansel R, Chandler RF, editors. *Adverse effects of herbal drugs, Vol 2.* Berlin Springer, 1993, 62.

Fintelmann V, Weiss RF. In: *Lehrbuch der Phytotherapie*, 10th ed. Stuttgart: Hippokrates, 2002, 85: 215.

Kraus A, Hammerschmidt FJ Untersuchungen an Fenchelölen, Dragoco report 1980, 27: 31-40.

Miething H, Seger V, Hänsel R. Determination of photoanethole from a stored essential oil of anise fruits as 4,4-dimethoxystilbene by high performance liquid chromatography-ultraviolet coupling. *Phytother Res* 1990, 4: 121-123.

Koller WD Z. *Lebensm Unters Forsch* 1976, 160: 143–147.

Leclerc A, Brugere J, Luce D, Point D, Guenel P . Type of alcoholic beverage and cancer of the upper respiratory and digestive tract. *Eur J Cancer Clin Oncol* 1987, 23: 529–534.

Schulte KE, Rücker G, Backe W. Polyacetylene from *pimpinella* species. *Arch Pharm* 1970, 303: 912–919.

Sharaf A, Goma N. Phytoestrogens and their antagonism to progesterone and testosterone. *J Endocrinol* 1965, 31: 289–290.

Zondek B, Bergman E. LXXXIV. Phenol methyl ethers as oestrogenic agents. *Biochm J* 1938, 32: 641–645.