



EUROPEAN MEDICINES AGENCY  
SCIENCE MEDICINES HEALTH

31 January 2017  
EMA/HMPC/80629/2016  
Committee on Herbal Medicinal Products (HMPC)

## List of references supporting the assessment of *Salix* [various species including *S. purpurea* L., *S. daphnoides* Vill., *S. fragilis* L.], cortex

Final

**The European Medicines Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.**

Adamkiewicz VM, Fortier AA. Passage of salicin and saligenin across the wall of the rat ileum. *Can J Biochem Physiol* 1961, 39:1097-99

Akao T, Yoshino T, Kobashi K, Hattori M. Evaluation of salicin as an antipyretic prodrug that does not cause gastric injury. *Planta Med* 2002, 68:714-18

Albrecht M, Nahrstedt, A. Luepke, N.P. Theisen, N.L. Baron, G. Anti-inflammatory activity of flavonol glycosides and salicin derivatives from the leaves of *Populus tremuloides*. *Planta Med* 1990, 56:660

American Herbal Pharmacopoeia. Willow bark. Scotts Valley 1999

Aronson JK (Ed). Meyler's side effects of drugs. *Elsevier Science*, Amsterdam 2006

Baker S, Thomas PS. Herbal medicine precipitating massive haemolysis [letter]. *Lancet* 1987, 1:1039-40

Barnes J, Anderson LA, Phillipson JD. Herbal Medicines, 3<sup>rd</sup> edition. *Pharmaceutical Press*. London 2007, 598-600

Beer A-M, Loew D. Phytopharmaka und Rückenschmerzen. *Fortschritte der Medizin* 2008, 150:33-37

Beer A-M, Helleman M, Schübel H, Wegener T. Weidenrinden Extract bei Gon- und Coxarthrose. *Zeitschrift für. Phytotherapie* 2008, 29:215-222

Beer A-M, Wegener T. Willow bark extract (Salicis cortex) for gonarthrosis and coxarthrosis – Results of a cohort study with a control group. *Phytomedicine* 2008, 15:907-913

Benedum J, Loew D, Schilcher H. Arzneipflanzen in der traditionelle Medizin. Kooperation Phytopharmaka, Bonn. *Salicis cortex Monographie* 2006, 179:320-321



- Biegert C, Wagner I, Lüdtkke R, Kötter I, Lohmüller C, Günaydin I *et al.* Efficacy and safety of willow bark extract in the treatment of OA and RA: results of 2 randomised db controlled trials. *J Rheumatol* 2004, 31:2121-30
- Bisset NG. Herbal Drugs and Phytopharmaceuticals. *Medpharm Scientific Publishers*, Stuttgart 1994, 437-439
- Blaschek *et al.* In Hagers Handbuch der Pharmazeutischen Praxis. Folgenband 3. Drogen L-Z. *Springer Verlag*, Berlin Heidelberg 1998, 469-96
- Bogduk N. Pharmacological alternatives for the alleviation of back pain. *Expert Opin Pharmacother* 2004, 5(10):2091-98
- Boullata J, McDonnell PJ, Oliva CD. Anaphylactic reaction to a dietary supplement containing willow bark. *Ann Pharmacother* 2003, 37:832-35
- BHMA. *British Herbal Pharmacopoeia*. Bournemouth 1983, 184-85
- BHMA. *British Herbal Compendium*. Bournemouth 1992, 224-26
- Bruneton J. Phytothérapie: les données de l'évaluation. *Tec et Doc Ed médicales internationales*. Paris 2002, 99-100
- Cheng GF, Liu DP, Yang DX, He KQ, Bai JY, Zhu XY. Anti-inflammatory effects of tremulacin, a salicin-related substance isolated from *Populus tomentosa* Carr leaves. *Phytomedicine* 1994, 1:209-11
- Chrubasik S, Eisenberg E, Balan E, Weinberger T, Luzzati R, Conradt C., Treatment of low back pain exacerbations with willow bark extract: a randomized double-blind study. *Am J Med* 2000, 109:9-14
- Chrubasik S, Künzel O, Model A, Conradt C, Black A. Treatment of low back pain with a herbal or synthetic anti-rheumatic: a randomized controlled study. Willow bark extract for low back pain. *Rheumatology* 2001a, 40:1388-93
- Chrubasik S, Kunzel O, Black A, Conradt C, Kerschbaumer F. Potential economic impact using a proprietary willow bark extract in outpatient treatment of low back pain: an open non-randomized study. *Phytomedicine* 2001b, 8(4):241-51
- Chrubasik S, Pollak S. Wirksamkeit von Weidenrindenextrakt bei Schmerzen. *Z Phytother* 2002, 23:263-66
- Chrubasik S. Weidenrindenextrakt verstärkt nicht die Regelblutung. *Naturamed* 2002, 17(12):13
- Chrubasik S. Effects of willow bark and therapeutic activity. *Clin Pharmacol Ther.* 2003, 74:95-97
- Chrubasik JE, Roufogalis BD, Chrubasik S. Evidence of effectiveness of herbal anti-inflammatory drugs in the treatment of painful osteoarthritis and chronic low back pain. *Phytother Res* 2007, 21:675-83
- Clauson KA, Santamarina ML, Buettner CM, Cauffield JS. Evaluation of presence of aspirin-related warnings with willow bark. *Ann Pharmacother* 2005, 39:1234-37
- Commission E. *Salicis cortex* (Weidenrinde). BAnz N° 228, 1984
- CPMP/EMEA. Points to consider on clinical investigation of medicinal products used in the treatment of osteoarthritis. CPMP/EWP/784/97, 1998
- CPMP/EMEA. Points to consider on clinical investigation of medicinal products other than NSAIDs for treatment of rheumatoid arthritis. CPMP/EWP/556/95 rev 1, 2003
- DAB 10 Deutsche *Apotheker Verlag*, Stuttgart 1991

- DAB 10. Kommentar. Deutsche *Apotheker Verlag*, Stuttgart 1995
- Dingerman T, Hänsel R, Zündorf I. Pharmazeutische Biologie. *Springer Verlag*, Berlin 2002, 65-78
- Drummond EM, Harbourne N, Marete E, Martyn D, Jacquier JC, O’Riordan D, Gibney ER. Inhibition of pro-inflammatory biomarkers in THP1 macrophages by polyphenols derived from chamomile, meadowsweet and willow bark. *Phytotherapy Research* 2013, 27:588-596
- Durak A, Gawlik-Dziki U. The Study of Interactions between Active Compounds of Coffee and Willow (*Salix* sp.) Bark Water Extract. *BioMed Research International* Volume 2014, Article ID 386953, 11 pages <http://dx.doi.org/10.1155/2014/386953>, 2014
- EDQM Willow bark dry extract. *Pharmeuropa* 2006, 18 (1):158-60
- EDQM. European Pharmacopoeia 5<sup>th</sup> edition, Willow bark. *Pharmeuropa* 2006, 18 (3):497-99
- Enayat S, Banerjee S. The Ethanolic Extract of Bark from *Salix Aegyptiaca* L. Inhibits the Metastatic Potential and Epithelial to Mesenchymal Transition of Colon Cancer Cell Lines. *Nutrition and Cancer* 2014, 66(6):999-1008
- ESCOP. Monographs; second edition: *Salicis cortex*, 2003. 445-51
- Esatbeyoglu T, Wray V, Winterhalter P. Dimeric Procyanidins: Screening for B1 to B8 and Semisynthetic Preparation of B3, B4, B6, and B8 from a Polymeric Procyanidin Fraction of White Willow Bark (*Salix alba*). *J. Agric. Food Chem.* 2010, 58:7820–7830
- Fiebig BL, Appel K. Anti-inflammatory effects of willow bark extract. *Clinical pharmacology & therapeutics* 2003, 96
- Fiebig BL, Müller J, Kelber O, Okpanyi S, Weiser D, Appel K. Does willow bark have a sildenafil-like action? Effect of different fraction on the synthesis of c-GMP and NO in human umbilical vein endothelial cell (HUVEC). *Planta Medica* 2010, 76:1190
- Fiebich BL, Chrubasik S. Effects of an ethanolic *Salix* extract on the release of selected inflammatory mediators *in vitro*. *Phytomed* 2004, 11:135-38
- Förster N, Ulrichs C, Zander M, Kätzel R, Mewis I. Salicylatreiche Weiden für die Arzneimittelherstellung. *Gesunde Pflanzen* 2009, 61:129–134
- Förster N, Ulrichs C, Zander M, Kätzel R, Mewis I. Factors influencing the variability of antioxidative phenolic glycosides in *Salix* species. *Agricultural and food chemistry* 2010; 58: 8205-8210.
- Fötsch G, Gründemann E, Pfeifer S, Hiller K, Salzwedel D. Biotransformation der Phenolglycosid Leiocarposid und Salicin. *Pharmazie* 1989a, 44:555-58
- Fötsch G, Pfeifer S. Die Biotransformation der Phenolglycosid Leiocarposid und Salicin – Beispiele für Besonderheiten von Absorption und Metabolismus glycosidischer Verbindungen. *Pharmazie* 1989b, 44:710-2
- Fötsch G, Pfeifer S. Vergleichende Serumspiegel-untersuchung von Salicylsäure nach oraler Applikation von Salicin bzw Natriumsalicylat in Ratten. *Pharmazie* 1990, 45:535-36
- Freischmidt A, Jürgenliemk G, Kraus B, Okpanyi SN, Müller J, Kelber O, *et al.* Contribution of flavonoids to the reduction of ICAM-1 expression in endothelial cells by a standardized willow bark extract. *Phytomedicine* 2012, 19:245-252
- Freischmidt A, Untergehrer M, Ziegler M, *et al.* Quantitative analysis of flavanones and chalcones from willow bark. *Pharmazie* 2015, 70:565-568

- Gagnier JJ, van Tulder MW, Berman B, Bombardier C. Herbal medicine for low back pain: a Cochrane Review. *Spine* 2007, 32(1):82-92
- Gagnier JJ, Oltean H, MPH, van Tulder MW, Berman BM, Bombardier C, Robbins CB. Herbal Medicine for Low Back Pain. *Spine* 2016, 41:116–133
- Gawlik-Dziki U, Sugier D, Dziki D, Sugier P. Bioaccessibility *In Vitro* of Nutraceuticals from Bark of Selected *Salix* Species. Hindawi Publishing Corporation Scientific World Journal Volume 2014, Article ID 782763, 10 pages <http://dx.doi.org/10.1155/2014/782763>
- Gopalan V, Pastuszyn A, Galey WR Jr, Glew RH. Exolytic hydrolysis of toxic plant glucosides by guinea pig liver cytosolic beta-glucosidase. *J Biol Chem* 1992, 267:14027-32
- Hänsel. Weidenrinde, *Salicis cortex*. In: *Phytopharmaka*, 2<sup>nd</sup> ed. Berlin-Heidelberg-New York: *Springer-Verlag* 1991, 96-97
- HagerRom. *Salicis cortex (Salix purpurea-Rinde)*, 2001
- Hostanska K, Jürgenliemk G, Abel G, Nahrstedt A, Saller R. Willow bark extract (BNO 1455) and its fractions suppress growth and induce apoptosis in human colon and lung cancer cells. *Cancer Detection and Prevention* 2007, 31:129-131
- Ishikado A, Sono Y, Matsumoto M, *et al.* Willow bark extract increases antioxidant enzymes and reduces oxidative stress through activation of Nrf2 in vascular endothelial cells and *Caenorhabditis elegans*. *Free Radical Biology and Medicine* 2013, 65:1506–1515
- Julkunen-Tiitto R, Meier B. The enzymatic decomposition of salicin and its derivatives obtained from *Salicaceae* species. *J Nat Prod* 1992a, 55:1204-12
- Julkunen-Tiitto R, Gebhardt K. Further studies on the drying of willow twigs: the effect of low drying temperature on labile phenolics. *Planta Med* 1992b, 58:385-86
- Julkunen-Tiitto R, Sorsa S. Testing the effects of drying methods on willow flavonoids, tannins and salicylates. *J Chem Ecol* 2001, 27(4):779-89
- Kammerer B, Kahlich R, Biegert C, Gleiter CH, Heide L. HPLC-MS/MS analysis of willow bark extracts contained in pharmaceutical preparations. *Phytochem Anal* 2005, 16:470-78
- Kaufeld AM, Pertz HH, Kolodziej H. A Chemically Defined 2,3-trans Procyanidin Fraction from Willow Bark Causes Redox-Sensitive Endothelium-Dependent Relaxation in Porcine Coronary Arteries. *J. Nat. Prod.* 2014, 77:1607–1614
- Kaul R, Lagoni N. Weidenrinde. Renaissance eines Phytoanalgetikums. *Deutsche Apothekerzeitung* 1999, 19(37):43-50
- Kelber O, Bonnaterra G, Kinscherf R, Weiser D, Metz J. Inhibitorische Effekte von Weidenrindenextrakten auf proinflammatorische Prozesse in LPS aktivierten Humanmonozyten. *Zeitschrift für Rheumatologie* 2006, 65 Suppl 1:S31
- Kelber O, Okpanyi SN, Freischmidt A, *et al.* Antidepressant mechanism of action of willow bark extract STW I-33. *Planta Medica* 2011, 77:1444
- Keusgen M, Algauer-Lechner C. Vielstoffgemisch zwischen Entzündungen und Schmerzen. *Pharmazeutische Zeitung* 2007, 152:16-22
- Khayyal MT, El-Ghazaly MA, Abdallah DM, Okpanyi SN, Kelber O, Weiser D. Mechanisms involved in the anti-inflammatory effect of a standardised willow bark extract. *Arzneim. Forsch / Drug Research* 2005, 55(11):677-87

- Kim CS, Subedi L, Jin Park K, *et al.* Salicin derivatives from *Salix glandulosa* and their biological activities. *Fitoterapia* 2015, 106:147–152
- Knuth S, Abdelsalam RM, Khayyal MT, *et al.* Catechol conjugates are *in vivo* metabolites of *Salicis cortex*. *Planta Med* 2013, 79:1489–1494
- Kong CS, Kim KH, Choi JS, Kim JE, Park C, Jeong JW. Salicin, an Extract from White Willow Bark, inhibits Angiogenesis by Blocking the ROS-ERK Pathways. *Phytother. Res.* 2014, 28:1246–1251
- Krivoy N, Pavlotzky E, Chrubasik S, Eisenberg E, Brook G. Effect of *Salicis cortex* extract on human platelet aggregation. *Planta Med* 2001, 67:209-12
- Kuppsamy UR, Khoo HE, Das NP. Structure-activity studies of flavonoids as inhibitors of hyaluronidase. *Biochem Biophys* 1990, 140:397-401
- Lardos A, Schmidlin C.B., Fischer M., Ferlas-Chlodny E., Loniewski I., Samochowiec L., *et al.* Wirksamkeit und Verträglichkeit eines wässrig ausgezogenen Weidenrindenextraktes bei Patienten mit Hüft-und Kniearthrose. *Z Phytotherap* 2004, 25:275-81
- Leporatti ML, Pavesi A. New or uncommon uses of several medicinal plants in some areas of central Italy. *J Ethnopharmacol* 1990, 29:213-23
- Madaus A. Lehrbuch biologische Heilmittel 1935, 2394-2399
- März RW, Kemper F. Weidenrindenextrakt – Wirkungen und Wirksamkeit. Erkenntnisstand zur Pharmacologie, Toxikologie und Klinik. Wien. Med. *Wochenschrift* 2002, 152:354-359
- Matsumoto Y, Ohsako M, Takadate A, Goto S. Reduction of erythrocyte membrane permeability and protein binding of LMW drugs following glycoside derivatisation. *J Pharm Sci* 1993, 82:399-403.
- Mayer RA, Mayer M. Biologische Salicyltherapie mit Cortex Salicis (Weidenrinde). *Pharmazie* 1949, 4:77-81
- McGuffin M, Hobbs C, Upton R. Botanical Safety Handbook. CRC Press Boca Raton Florida 1997 , 154-155
- Meier B, Lehmann D, Sticher O, Bettschart A . Identification and determination of 8 phenol glycosides each in *Salix purpurea* and *S. daphnoides* by modern HPLC. *Pharm Acta Helv* 1985a, 60:269-75
- Meier B, Sticher O, Zürich, Bettschart A Weidenrinde – Qualität. Gesamtsalicinbestimmung in Weidenrinden und Weidenpräparaten mit HPLC. *Deutsche Apoth Ztg* 1985b, 125:341-47
- Meier B, Libi M. Salizinhaltige pflanzliche Arzneimittel. Überlegungen zur Wirksamkeit und Unbedenklichkeit. *Z Phytother* 1990, 11:50-58
- Mills SY, Jacoby RK, Chacksfield M, Willoughby M. Effect of a proprietary herbal medicine on the relief of chronic arthritic pain: a double-blind study. *Br J Rheumatol* 1996, 35:874-78
- Morgan M, Bone K, Mills S, McMillan J. Part 2: Safety Monographs, in: Mills S and Bone K, The Essential Guide Herbal Safety, Elsevier Churchill Livingstone, St. Louis 2005, 626-628
- Moro PA, Flacco V, Cassetti F, Clementi V, Colombo ML, Chiesa GM, Menniti-Ippolito F, Raschetti R, Santucci C. Hypovolemic shock due to severe gastrointestinal bleeding in a child taking an herbal syrup. *Ann Ist Super Sanità*, 2011, Vol. 47, 3:278-283
- Müller-Fassbender H, Müller J, Bach G. Wirksamkeit und Verträglichkeit eines Weidenrindenpräparates bei Erkrankungen des Bewegungsapparates. *Arthritis und Rheuma* 2007, 27:267-271

- Müller J, Kelber O, Weiser D, Stange S. Willow bark extract STW 33-1 in the long term treatment of osteoarthritis and low back pain. *Planta Medica* 2010, 76:1361
- Oltean H, Robbins C, van Tulder MW, Berman BM, Bombardier C, Gagnier JJ. Herbal medicine for low back pain. Cochrane Database of Systematic Reviews 2014, Issue 12. Art. No.: CD004504 DOI: 10.1002/14651858.CD004504.pub4
- Nahrstedt A, Schmidt M, Jäggi R, Metz J, Khayyal MT. Willow bark extract: the contribution of polyphenols to the overall effect. *Wien Med Wochenschr* 2007, 157(13-14):348-51
- Pentz R, Busse HG, König R, Siegers CP. Bioverfügbarkeit von Salicylsäure und Coffein aus einem phytoanalgetischen Kombinationspräparat. *Z Phytother* 1989, 10:92-96
- Pugliese A, Beltramo T, Torre D. Reye's and Reye's-like syndromes. *Cell Biochem Function* 2008; 26:741-746
- Rice-Evans CA, Miller NJ, Bolwell PG, Bramley PM, Pridham JB. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Rad Ser* 1995, 22(4):375-83
- Rohnert U, Koske D, Schneider W, Elstner EF. Inhibition by *Salix*-extracts and Phytodolor® of copper-catalyzed oxidative destructions. *Z Naturforsch* 1998, 53:233-40
- Rotblatt M. Evidence-based herbal medicine. Hanley and Belfus, Philadelphia 2002, 360-63
- Saller R, Melzer J, Felder M, the Swiss Assalix Study Group. Pain relief with a proprietary extract of willow bark in rheumatology. An open trial. *Schweiz Zschr GanzheitsMedizin* 2008, 20(3):156-62
- Samochowiec L. Efficacy of Cortex *Salix* Extract (*Salix* (r)) in patients with arthrosis in double blind, randomized, verum controlled clinical study 2001 (unpublished)
- Schmid B, Kötter I, Heide L. Pharmacokinetics of salicin after oral administration of a standardized willow bark extract. *Eur J Clin Pharmacol* 2001a, 57:387-91
- Schmid B, Lüdtke R, Selbmann HK, Kötter I, Tschirdewahn B, Schaffner W, et al. Efficacy and tolerability of a standardized willow bark extract in patients with osteoarthritis: randomized placebo-controlled double blind clinical trial. *Phyt Res* 2001b, 15:344-50
- Schmid B, Biegert C, Kötter I, Heide L. Pharmakinetik von Salicin nach oraler Gabe eines standardisierten Weidenrindenextraktes, in: *Phytopharmaka VII*. Schulz V, Rietbrock N, Roots I, Loew D, *Steinkopf*, Darmstadt 2002: 93-100
- Setty AR, Sigal LH. Herbal medications commonly used in the practice of rheumatology: mechanisms of action, efficacy and side effects. *Semin Arthritis Rheum* 2005, 34:773-84
- Shalansky S, Lynd L, Richardson K, Ingaszewski A, Kerr C. Risk of warfarin-related bleeding events and supratherapeutic INRs associated with CAM: a longitudinal analysis. *Pharmacother* 2007, 27(9):1237-47
- Shakibaei M, Allaway D, Nebrich S, Mobasheri A. Botanical Extracts from Rosehip (*Rosa canina*), Willow Bark (*Salix alba*), and Nettle Leaf (*Urtica dioica*) Suppress IL-1 $\beta$ -Induced NF- $\kappa$ B Activation in Canine Articular Chondrocytes. Evidence-Based Complementary and Alternative Medicine Volume 2012, Article ID 509383, 16 pages doi:10.1155/2012/509383
- Shara M, Stohs SJ. Efficacy and Safety of White Willow Bark (*Salix alba*) Extracts. *Phytother. Res.* 2015, 29:1112–1116

Shrivastava R, Pechadre JC, John GW. *Tanacetum parthenium* and *Salix alba* (Mig-RL) combination in migraine prophylaxis. *Clin Drug Invest* 2006, 26(5):287-96

SPC Aspirin Junior; May 2016

Stange R, Müller J, Kelber O, Uehleke B. Willow bark extract STW 33-I is safe and effective in the long term treatment of outpatients with chronic musculoskeletal pain. *J Alternative Complementary Med* 2014, 20:A83

Steinegger E, Hövel H. Analytische and biologische Untersuchungen and Salicaceen-Wirkstoffen, insbesondere Salicin. *Pharm Acta Helv* 1972, 47:222-34

Tunón H1, Olavsdotter C, Bohlin L. Evaluation of anti-inflammatory activity of some Swedish medicinal plants. Inhibition of prostaglandin biosynthesis and PAF-induced exocytosis. *J Ethnopharmacol* 1995, 48:61-76

Uehleke B, Preis S, Müller J, Stange R, Kelber O, Melzer J. Willow bark extract STW 33-I in the long-term treatment of outpatients with rheumatic pain mainly osteoarthritis or back pain. *Phytomedicine* 2013a, 980–984

Uehleke B, Müller J, Stange R, Kelber O, Melzer J. Willow bark extract STW 33-I is safe and effective in the long term treatment of outpatients with rheumatic pain esp. osteoarthritis or backpain - a subgroup analysis. *Planta Medica* 2013b, 79:1273 (abstract)

Ulrich-Merzenich G, Kelber O, Freischmidt A, et al. Evidences for an antidepressant effect of willow bark in the Porsolt swimming test. *European Journal of Integrative Medicine* 2010, 2:188 (abstract)

Ulrich-Merzenich G, Koptina A, Kelber O, et al. Comparative analysis of gene expression profiles from rats treated with a standardized willow bark preparation, its salicin rich ethanol fraction and imipramine for toxicological outcomes. *Planta Medica* 2011, 77:1251

Ulrich-Merzenich G, Koptina A, Kelber O, et al. Prediction of adverse events by in vivo gene expression profiling exemplified for phytopharmaceuticals containing salicylates and the antidepressant imipramine. *Phytomedicine* 2012, 19:322–329

Ulrich-Merzenich G, Hartbrod F, Koptina A. Synergie-Effekte als Grundprinzip der Wirkung von Phytopharmaka am Beispiel der Weidenrinde (*Salix spec.*) *Z Phytother* 2013, 34:34

Vlachojannis JE, Cameron M, Chubasick S. A systematic review on the effectiveness of willow bark for musculoskeletal pain. *Phytotherapy Research* 2009, 23:897-900

Vlachojannis J, Magora F, Chrubasik S. Willow Species and Aspirin: Different Mechanism of Actions. *Phytother. Res.* 2011, 25:1102–1104

Volmink J. The willow as a Hottentot (Khoikhoi) remedy for rheumatic fever. *J R Soc Med* 2008, 101:321-323

Wagner I, Greim I, Laufer C, Heide S, L, Gleiter CH. Influence of willow bark extract on cyclooxygenase activity and on TNF alpha or interleukin 1 beta release in vitro and in vivo. *Clin Pharmacol Therap* 2003a, 73(3):272-74

Wagner I, Biegert C, und Lutz Heide, Aktuelle Forschungsergebnisse zur Weidenrinde. *Pharm. Ztg* 2003b, 138:21-32

Wagner I., Effects of willow bark. *Clin Pharmacol Therap* (letters to editor)2003c, 96-97

Wegener T. Zur Studienlage der Weidenrindenextrakten. *Zietschrift für Phytotherapie* 2009, 30:232-237

Werner G, Scheithe K. Willow bark extract (Assalix ®) for chronic back pain and arthralgia, a post-authorisation study. *Phytopharmaka und Phytotherapie* 2004, Forschung und Praxis, Berlin (poster abstract)

Wichtl M (ed) Teedrogen und Phytopharmaka 4<sup>th</sup> ed, Stuttgart: *Wissenschaftliche Verlagsgesellschaft*, 2002, 534-7

Wiesneth S, Petereit F, Jürgenliemk G. *Salix daphnoides*: a screening for oligomeric and polymeric proanthocyanidins. *Molecules* 2015, 20:13764-13779

Williamson E, Driver S, Baxter K. Stockley's herbal medicines interactions. *Pharmaceutical Press* 2<sup>nd</sup> ed. London 2013, 485

Wuthold K, Germann I, Roos G, Kelber O, Weiser D, Heinle H, Kovar KA. TLC and multivariate data analysis of willow bark extracts. *J Chromatogr Sci* 2004, 42(6)306-09

Wutzke KD. Radioisotopenmarkierung und Pharmakokinetik von Phenolglycosiden. Dissertation Phillipsuniversität Marburg 1991