



EUROPEAN MEDICINES AGENCY  
SCIENCE MEDICINES HEALTH

12 November 2013  
EMA/HMPC/317318/2012  
Committee on Herbal Medicinal Products (HMPC)

## List of references supporting the assessment of *Phaseolus vulgaris* L., fructus sine semine

Final

**The Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.**

Braun H. Heilpflanzen – Lexikon für Ärzte und Apotheker, Anwendung, Wirkung und Toxicologie. 4<sup>th</sup> ed. Gustav Fisher Verlag, Stuttgart, New York 1981, 162

Cerović A, Miletić I, Konić-Ristić A, Baralić I, Djordjević B, Djurčić I, Radusinović M. The dry plant extract of common bean seed (*Phaseoli vulgaris pericarpium*) does not have an effect on postprandial glycaemia in healthy human subject. *Bosn J Basic Med Sci* 2006, 6(3):28-33

Czech Pharmaceutical Codex. *Phaseoli fructus sine semine*. Český farmaceutický kodex (Codex Pharmaceuticus Bohemicus). Editio prima, X-EGEM Publishers, Praha 1993

DAC. Deutscher Arzneimittel Codex. *Phaseoli fructus sine semine*. Deutscher Apotheker-Verlag, Stuttgart, 1986

Delfosse M. Drogues végétales et plantes médicinales. APB Service scientifique, Brussels, 1998, 249

Dodoens R. Herbarius oft Cruydt-Boeck van Rembertus Dodonaeus. Plantijn – Moretus, Antwerpen 1608, 910

Hagers Handbuch der Pharmazeutischen Praxis (CDRom) 2012

Helmstädter A. Beans and diabetes: *Phaseolus vulgaris* preparations as antihyperglycemic agents. *J Med Food* 2010, 13(2):251-254

Martinez Alonso JC, Callejo Melgosa A, Fuentes Gonzalo MJ, Martin Garcia C. Angioedema induced by inhalation of vapours from cooked white bean in a child. *Allergol Immunopathol (Madr)* 2005, 33(4):228-30

Neef H, Declercq P, Laekeman G. Hypoglycaemic activity of selected European plants. *Phytother Res* 1995, 9:45-48



- Ocho-Anin Atchibri AL, Kouakou TH, Brou KD, Kouadio YJ, Gnakri D. Evaluation of bioactive components in seeds of *Phaseolus vulgaris* L. (Fabaceae) cultivated in Côte d'Ivoire. *J Appl Biosci* 2010, 31:1928-1934
- Pari L, Venkateswaran S. Effect of an aqueous extract of *Phaseolus vulgaris* on plasma insulin and hepatic key enzymes of glucose metabolism in experimental animals. *Pharmazie* 2003, 58(12):916-919
- Pari L, Venkateswaran S. Protective role of *Phaseolus vulgaris* on changes in the fatty acid composition in experimental diabetes. *J Med Food* 2004, 7(2):204-209
- Reuter H.D. Therapie mit Phytopharmaka (Pharmakologie – Indikationen Dosierungen) 1<sup>st</sup> ed. Gustav Fisher Verlag, Stuttgart, 1997, 153-154
- Reynoso-Camacho R, Ramos-Gomez M, Loarca-Pina G. Advances in Agricultural and Food Biotechnology. Bioactive components in common beans (*Phaseolus vulgaris* L.), 2006, 217-236
- Roman-Ramos R, Flores-Saenz JL, Alarcon-Aguilar FJ. Anti-hyperglycaemic effect of some edible plants. *J Ethnopharmacol* 1995, 48(1):25-32
- Rougé P, Culerrier R, Thibau F, Didier A, Barre A. A case of severe anaphylaxis to kidney bean: phaseolin (vicilin) and PHA (lectin) identified as putative allergens. *Allergy* 2011, 66(2):301-302
- Van Hellemont J. Phytotherapeutisch compendium. Algemene Pharmaceutische Bond. 1985, 429-430
- Verhelst G. Groot Handboek Geneeskrachtige Planten. 4<sup>th</sup> ed. BVBA Mannavita, 2010, 427-428
- Wichtl M. Teedrogen (Ein Handbuch für Apotheker und Ärzte). Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, 1984, 81-82, 376
- Wichtl M. Herbal Drugs and Phytopharmaceuticals. Medpharm GmbH Scientific Publishers, Stuttgart, 1994, 373-374

*References not supporting the assessment report*

Arija I, Centeno C, Viveros A, Brenes A, Marzo F, Illera JC et al. Nutritional evaluation of raw and extruded kidney bean (*Phaseolus vulgaris* L. var. *pinto*) in chicken diets. *Poult Sci* 2006, 85(4):635-644 [Epub 2006/04/18]

Available at: [http://www.phase2info.com/pdf/Phase2\\_Study10.pdf](http://www.phase2info.com/pdf/Phase2_Study10.pdf). Accessed 2012/03/07

Barrett ML, Udani JK. A proprietary alpha-amylase inhibitor from white bean (*Phaseolus vulgaris*): a review of clinical studies on weight loss and glycemic control. *Nutr J* 2011, 10: 24

Carai MA, Fantini N, Loi B, Colombo G, Gessa GL, Riva A et al. Multiple cycles of repeated treatments with a *Phaseolus vulgaris* dry extract reduce food intake and body weight in obese rats. *Br J Nutr* 2011, 106(5):762-768 [Epub 2011/05/04]

Carai MAM, Fantini N, Loi B, Colombo G, Riva A, Morazzoni P. Potential efficacy of preparations derived from *Phaseolus vulgaris* in the control of appetite, energy intake, and carbohydrate metabolism. *Diabetes Metab Syndr Obes* 2009, 2: 145-153

Carmalt J, Rosel K, Burns T, Janzen E. Suspected white kidney bean (*Phaseolus vulgaris*) toxicity in horses and cattle. *Aust Vet J*. 2003, 81(11):674-676 [Epub 2004/04/17]

Celleno L, Tolaini MV, D'Amore A, Perricone NV, Preuss HG. A Dietary supplement containing standardized *Phaseolus vulgaris* extract influences body composition of overweight men and women. *Int J Med Sci* 2007, 4(1):45-52

Chokshi D. Subchronic oral toxicity of a standardized white kidney bean (*Phaseolus vulgaris*) extract in rats. *Food Chem Toxicol* 2007, 45(1):32-40 [Epub 2006/10/19]

Chokshi D. Toxicity studies of Blockal™, a dietary supplement containing Phase 2 Starch Neutralizer (Phase 2), a standardized extract of the common white kidney bean (*Phaseolus vulgaris*). *Int J Toxicol* 2006, 25(5):361-371 [Epub 2006/08/31]

Dilawari JB, Kamath PS, Batta RP, Mukewar S, Raghavan S. Reduction of postprandial plasma glucose by Bengal gram dal (*Cicer arietinum*) and rajmah (*Phaseolus vulgaris*). *Am J Clin Nutr* 1981, 34(11):2450-2453 [Epub 1981/11/01]

Donatucci DA, Liener IE, Gross CJ. Binding of navy bean (*Phaseolus vulgaris*) lectin to the intestinal cells of the rat and its effect on the absorption of glucose. *J Nutr* 1987, 117(12):2154-2160 [Epub 1987/12/01]

Erner S, Meiss D. The Effect of TheraSlim™ (with Phase 2) on Weight, Body Composition, and Select Laboratory Parameters in Adults with Overweight and Mild-Moderate Obesity, 2003  
Available at: [http://www.phase2info.com/pdf/Phase2\\_Study8.pdf](http://www.phase2info.com/pdf/Phase2_Study8.pdf). Accessed 2012/03/07

Fantini N, Cabras C, Lobina C, Colombo G, Gessa GL, Riva A, et al. Reducing effect of a *Phaseolus vulgaris* dry extract on food intake, body weight, and glycemia in rats. *J Agric Food Chem* 2009, 57(19):9316-9323 [Epub 2009/09/08]

Grant G, Dorward PM, Buchan WC, Armour JC, Pusztai A. Consumption of diets containing raw soya beans (*Glycine max*), kidney beans (*Phaseolus vulgaris*), cowpeas (*Vigna unguiculata*) or lupin seeds (*Lupinus angustifolius*) by rats for up to 700 days: effects on body composition and organ weights. *Br J Nutr* 1995, 73(1):17-29 [Epub 1995/01/01]

Harikumar KB, Jesil AM, Sabu MC, Kuttan R. A preliminary assessment of the acute and subchronic toxicity profile of phase2: an alpha-amylase inhibitor. *Int J Toxicol* 2005, 24(2):95-102 [Epub 2005/07/23]

Kik MJ, Huisman J, van der Poel AF, Mouwen JM. Pathologic changes of the small intestinal mucosa of pigs after feeding *Phaseolus vulgaris* beans. *Vet Pathol* 1990, 27(5):329-334 [Epub 1990/09/01]

Kilpatrick DC, Green C, Yap PL. Lectin content of slimming pills. *Br Med J (Clin Res Ed)* 1983, 286(6361):305 [Epub 1983/01/22]

Kluh I, Horn M, Hyblova J, Hubert J, Doleckova-Maresova L, Voburka Z et al. Inhibitory specificity and insecticidal selectivity of alpha-amylase inhibitor from *Phaseolus vulgaris*. *Phytochemistry* 2005, 66(1):31-39 [Epub 2005/01/15]

Koike T, Koizumi Y, Tang L, Takahara K, Saitou Y. The anti-obesity effect and the safety of taking "Phaseolamin™ 1600 diet". *Journal of New Remedies and Clinics (Japanese)* 2005, 54:1-16 [Japanese]

Lara-Diaz VJ, Gaytan-Ramos AA, Davalos-Balderas AJ, Santos-Guzman J, Mata-Cardenas BD, Vargas-Villarreal J et al. Microbiological and toxicological effects of Perla black bean (*Phaseolus vulgaris* L.) extracts: in vitro and in vivo studies. *Basic Clin Pharmacol Toxicol* 2009, 104(2):81-86

Martin-Cabrejas MA, Jaime L, Karanja C, Downie AJ, Parker ML, Lopez-Andreu FJ, et al. Modifications to physicochemical and nutritional properties of hard-to-cook beans (*Phaseolus vulgaris* L.) by extrusion cooking. *J Agric Food Chem* 1999, 47(3):1174-1182 [Epub 1999/12/20]

Marzo F, Alonso R, Urdaneta E, Arricibita FJ, Ibanez F. Nutritional quality of extruded kidney bean (*Phaseolus vulgaris* L. var. *pinto*) and its effects on growth and skeletal muscle nitrogen fractions in rats. *J Anim Sci* 2002, 80(4):875-879 [Epub 2002/05/11]

Marzolo MP, Amigo L, Nervi F. Hepatic production of very low density lipoprotein, catabolism of low density lipoprotein, biliary lipid secretion, and bile salt synthesis in rats fed a bean (*Phaseolus vulgaris*) diet. *J Lipid Res* 1993, 34(5):807-814 [Epub 1993/05/01]

Obiro WC, Zhang T, Jiang B. The nutraceutical role of the *Phaseolus vulgaris* alpha-amylase inhibitor. *Br J Nutr* 2008, 100(1):1-12 [Epub 2008/03/12]

Onakpoya I, Aldaas S, Terry R, Ernst E. The efficacy of *Phaseolus vulgaris* as a weight-loss supplement: A systematic review and meta-analysis of randomised clinical trials. *Br J Nutr* 2011, 106(2):196-202

Opala T, Rzymiski P, Pischel I, Wilczak M, Wozniak J. Efficacy of 12 weeks supplementation of a botanical extract-based weight loss formula on body weight, body composition and blood chemistry in healthy, overweight subjects - A randomised double-blind placebo-controlled clinical trial. *Eur J Med Res* 2006, 11(8):343-350

Osorio L, Gamboa J. Random multicenter evaluation to test the efficacy of *Phaseolus vulgaris* (Precarb™) in obese and overweight individuals. Internal Report, 2005  
Available at: [http://www.phase2info.com/pdf/Phase2\\_Study10.pdf](http://www.phase2info.com/pdf/Phase2_Study10.pdf). Accessed 2012/03/07

Preuss HG. Bean amylase inhibitor and other carbohydrate absorption blockers: effects on diabetes and general health. *J Am Coll Nutr* 2009, 28(3):266-276 [Epub 2010/02/13]

Pusztai A, Grant G, Buchan WC, Bardocz S, de Carvalho AF, Ewen SW. Lipid accumulation in obese Zucker rats is reduced by inclusion of raw kidney bean (*Phaseolus vulgaris*) in the diet. *Br J Nutr* 1998, 79(2):213-221 [Epub 1998/04/16]

Pusztai A, Grant G, Duguid T, Brown DS, Peumans WJ, Van Damme EJ et al. Inhibition of starch digestion by alpha-amylase inhibitor reduces the efficiency of utilization of dietary proteins and lipids and retards the growth of rats. *J Nutr* 1995, 125(6):1554-1562 [Epub 1995/06/01]

- Pusztai A, King TP, Clarke EM. Recent advances in the study of the nutritional toxicity of kidney bean (*Phaseolus vulgaris*) lectins in rats. *Toxicol* 1982, 20(1):195-197 [Epub 1982/01/01]
- Reeves PG, Nielsen FH, Fahey GC Jr. AIN-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the AIN-66A rodent diet. *J Nutr* 1993, 123(11): 1939-1951
- Rondanelli M, Giacosa A, Orsini F, Opizzi A, Villani S. Appetite control and glycaemia reduction in overweight subjects treated with a combination of two highly standardized extracts from *Phaseolus vulgaris* and *Cynara scolymus*. *Phytother Res* 2011, 25(9):1275-82
- Rothacker D. Reduction in body weight with a starch blocking diet aid: StarchAway™ comparison with placebo. *Leiner Health Products*, 2003. Available at: [http://www.phase2info.com/pdf/Phase2\\_Study6.pdf](http://www.phase2info.com/pdf/Phase2_Study6.pdf). Accessed 2012/03/07
- Santidrian S, de Moya CC, Grant G, Fruhbeck G, Urdaneta E, Garcia M, et al. Local (gut) and systemic metabolism of rats is altered by consumption of raw bean (*Phaseolus vulgaris* L. var *athropurpurea*). *Br J Nutr* 2003, 89(3):311-319 [Epub 2003/03/12]
- Santimone M, Koukietolo R, Moreau Y, Le Berre V, Rouge P, Marchis-Mouren G, et al. Porcine pancreatic alpha-amylase inhibition by the kidney bean (*Phaseolus vulgaris*) inhibitor (alpha-AI1) and structural changes in the alpha-amylase inhibitor complex. *Biochim Biophys Acta*. 2004, 1696(2):181-190 [Epub 2004/02/12]
- Tegua A, Fru SF. The growth performances of broiler chickens as affected by diets containing common bean (*Phaseolus vulgaris*) treated by different methods. *Trop Anim Health Prod* 2007, 39(6):405-410 [Epub 2007/10/31]
- Thom E. A randomized, double-blind, placebo-controlled trial of a new weight-reducing agent of natural origin. *J Int Med Res* 2000, 229-233
- Tormo MA, Gil-Exojo I, Romero de Tejada A, Campillo JE. Hypoglycaemic and anorexigenic activities of an alpha-amylase inhibitor from white kidney beans (*Phaseolus vulgaris*) in Wistar rats. *Br J Nutr* 2004, 92(5):785-790 [Epub 2004/11/10]
- Tormo MA, Gil-Exojo I, Romero de Tejada A, Campillo JE. White bean amylase inhibitor administered orally reduces glycaemia in type 2 diabetic rats. *Br J Nutr* 2006, 96(3):539-544 [Epub 2006/08/24]
- Udani J, Hardy M, Madsen DC. Blocking carbohydrate absorption and weight loss: a clinical trial using Phase 2 brand proprietary fractionated white bean extract. *Altern Med Rev* 2004, 9(1):63-69 [Epub 2004/03/10]
- Udani J, Singh BB. Blocking carbohydrate absorption and weight loss: a clinical trial using a proprietary fractionated white bean extract. *Altern Ther Health Med* 2007, 13(4):32-37 [Epub 2007/07/31]
- Udani JK, Singh BB, Barrett ML, Preuss HG. Lowering the glycaemic index of white bread using a white bean extract. *Nutr J*. 2009, 8:52 [Epub 2009/10/29]
- Vinson J, Al Kharrat H, Shuta D. Investigation of an amylase inhibitor on human glucose absorption after starch consumption. *The Open Nutraceuticals Journal* 2009, 2:88-91
- Wu X, Xu X, Shen J, Perricone N, Preuss H. A dietary supplement containing standardized *Phaseolus vulgaris* extract influences body composition of overweight men and women. *Journal of Applied Research* 2010, 10:73-79
- Yamada J, Yamamoto T, Yamaguchi T. Effects of combination of functional food materials on body weight, body fat percentage, serum triglyceride and blood glucose. 2007