

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Thalidomide BMS 50 mg hard capsules

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each capsule contains 50 mg of thalidomide.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Hard capsule.

White opaque capsules marked “Thalidomide BMS 50 mg”.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Thalidomide BMS in combination with melphalan and prednisone is indicated as first line treatment of patients with untreated multiple myeloma, aged  $\geq 65$  years or ineligible for high dose chemotherapy.

Thalidomide BMS is prescribed and dispensed according to the Thalidomide BMS Pregnancy Prevention Programme (see section 4.4).

### 4.2 Posology and method of administration

Treatment must be initiated and monitored under the supervision of physicians with expertise in managing immunomodulatory or chemotherapeutic agents and a full understanding of the risks of thalidomide therapy and monitoring requirements (see section 4.4).

#### Posology

The recommended dose of thalidomide is 200 mg orally per day.

A maximum number of 12 cycles of 6 weeks (42 days) should be used.

**Table 1: Starting doses for thalidomide in combination with melphalan and prednisone**

Age (years)	ANC* (/ $\mu$ L)		Platelet Count (/ $\mu$ L)	Thalidomide <sup>a,b</sup>	Melphalan <sup>c,d,e</sup>	Prednisone <sup>f</sup>
$\leq 75$	$\geq 1,500$	AND	$\geq 100,000$	200 mg daily	0.25 mg/kg daily	2 mg/kg daily
$\leq 75$	$< 1,500$ but $\geq 1,000$	OR	$< 100,000$ but $\geq 50,000$	200 mg daily	0.125 mg/kg daily	2 mg/kg daily
$> 75$	$\geq 1,500$	AND	$\geq 100,000$	100 mg daily	0.20 mg/kg daily	2 mg/kg daily
$> 75$	$< 1,500$ but $\geq 1,000$	OR	$< 100,000$ but $\geq 50,000$	100 mg daily	0.10 mg/kg daily	2 mg/kg daily

\* ANC: Absolute Neutrophil Count

<sup>a</sup> Thalidomide dosed once daily at bedtime on Days 1 to 42 of each 42-day cycle.

<sup>b</sup> Due to the sedative effect associated with thalidomide, administration at bedtime is known to generally improve tolerability.

<sup>c</sup> Melphalan dosed once daily on Days 1 to 4 of each 42-day cycle.

<sup>d</sup> Melphalan dosing: reduce by 50 % for moderate (creatinine clearance:  $\geq 30$  but  $< 50$  mL/min) or severe (CrCl:  $< 30$  mL/min) renal insufficiency

<sup>e</sup> Maximum daily melphalan dose: 24 mg (subjects  $\leq 75$  years old) or 20 mg (subjects  $> 75$  years old).

<sup>f</sup> Prednisone dosed once daily on Days 1 to 4 of each 42-day cycle.

Patients should be monitored for: thromboembolic events, peripheral neuropathy, severe skin reactions, bradycardia, syncope, somnolence, neutropenia and thrombocytopenia (see sections 4.4 and 4.8). Dose delay, reduction or discontinuation, dependent upon the NCI CTC (National Cancer Institute Common Toxicity Criteria) grade, may be necessary.

If less than 12 hours has elapsed since missing a dose, the patient can take the dose. If more than 12 hours has elapsed since missing a dose at the normal time, the patient should not take the dose, but take the next dose at the normal time on the following day.

#### *Thromboembolic events*

Thromboprophylaxis should be administered for at least the first 5 months of treatment especially in patients with additional thrombotic risk factors. Prophylactic antithrombotic medicinal products, such as low molecular weight heparins or warfarin, should be recommended. The decision to take antithrombotic prophylactic measures should be made after careful assessment of an individual patient's underlying risk factors (see sections 4.4, 4.5 and 4.8).

If the patient experiences any thromboembolic events, treatment must be discontinued and standard anticoagulation therapy started. Once the patient has been stabilised on the anticoagulation treatment and any complications of the thromboembolic event have been managed, the thalidomide treatment may be restarted at the original dose dependent upon a benefit-risk assessment. The patient should continue anticoagulation therapy during the course of thalidomide treatment.

#### *Neutropenia*

White blood cell count and differential should be monitored on an ongoing basis, in accordance with oncology guidelines, especially in patients who may be more prone to neutropenia. Dose delay, reduction or discontinuation, dependent upon the NCI CTC grade, may be necessary.

#### *Thrombocytopenia*

Platelet counts should be monitored on an ongoing basis, in accordance with oncology guidelines. Dose delay, reduction or discontinuation, dependent upon the NCI CTC grade, may be necessary.

#### *Peripheral neuropathy*

Dose modifications due to peripheral neuropathy are described in Table 2.

**Table 2: Recommended dose modifications for thalidomide -related neuropathy in first line treatment of multiple myeloma**

<b>Severity of neuropathy</b>	<b>Modification of dose and regimen</b>
Grade 1 (paraesthesia, weakness and/or loss of reflexes) with no loss of function	Continue to monitor the patient with clinical examination. Consider reducing dose if symptoms worsen. However, dose reduction is not necessarily followed by improvement of symptoms.
Grade 2 (interfering with function but not with activities of daily living)	Reduce dose or interrupt treatment and continue to monitor the patient with clinical and neurological examination. If no improvement or continued worsening of the neuropathy, discontinue treatment. If the neuropathy resolves to Grade 1 or better, the treatment may be restarted, if the benefit/risk is favourable.
Grade 3 (interfering with activities of daily living)	Discontinue treatment
Grade 4 (neuropathy which is disabling)	Discontinue treatment

#### *Allergic reactions and severe skin reactions*

Thalidomide interruption or discontinuation should be considered for Grade 2-3 skin rash.

Thalidomide must be discontinued for angioedema, anaphylactic reaction, Grade 4 rash, exfoliative or bullous rash, or if Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) or drug reaction

with eosinophilia and systemic symptoms (DRESS) is suspected and should not be resumed following discontinuation for these reactions.

#### *Elderly population*

No specific dose adjustments are recommended for the elderly  $\leq 75$  years of age. For patients  $> 75$  years of age, the thalidomide recommended starting dose is 100 mg per day. The initial dose of melphalan is reduced for elderly  $> 75$  years of age considering baseline bone marrow reserve and renal function. The melphalan recommended starting dose is 0.1 to 0.2 mg/kg daily according to bone marrow reserve along with a further 50 % dose reduction for moderate (creatinine clearance:  $\geq 30$  but  $< 50$  mL/minute) or severe (CrCl:  $< 30$  mL/minute) renal insufficiency. The maximum daily melphalan dose is 20 mg in patients  $> 75$  years of age (see Table 1).

#### *Patients with renal or hepatic impairment*

Thalidomide BMS has not formally been studied in patients with impaired renal or hepatic function. No specific dose recommendations for these patient populations are available. Patients with severe organ impairment should be carefully monitored for adverse reactions.

#### *Paediatric population*

There is no relevant use of Thalidomide BMS in the paediatric population in the indication of multiple myeloma.

#### Method of administration

Thalidomide BMS should be taken as a single dose at bedtime, to reduce the impact of somnolence. Capsules should not be opened or crushed (see section 6.6).

It is recommended to press only on one end of the capsule to remove it from the blister, thereby reducing the risk of capsule deformation or breakage.

### **4.3 Contraindications**

- Hypersensitivity to thalidomide or to any of the excipients listed in section 6.1.
- Women who are pregnant (see section 4.6).
- Women of childbearing potential unless all the conditions of the Pregnancy Prevention Programme are met (see sections 4.4 and 4.6).
- Male patients unable to follow or comply with the required contraceptive measures (see section 4.4).

### **4.4 Special warnings and precautions for use**

#### Teratogenic effects

Thalidomide is a powerful human teratogen, inducing a high frequency of severe and life-threatening birth defects. Thalidomide must never be used by women who are pregnant or by women who could become pregnant unless all the conditions of the Pregnancy Prevention Programme are met. The conditions of the Pregnancy Prevention Programme must be fulfilled for all male and female patients.

#### Criteria for women of non-childbearing potential

A female patient or a female partner of a male patient is considered to have childbearing potential unless she meets at least one of the following criteria:

- Age  $\geq 50$  years and naturally amenorrhoeic for  $\geq 1$  year (Amenorrhoea following cancer therapy or during breast-feeding does not rule out childbearing potential).
- Premature ovarian failure confirmed by a specialist gynaecologist.
- Previous bilateral salpingo-oophorectomy, or hysterectomy.
- XY genotype, Turner's syndrome, uterine agenesis.

### Counselling

For women of childbearing potential, thalidomide is contraindicated unless all of the following conditions are met:

- She understands the teratogenic risk to the unborn child
- She understands the need for effective contraception, without interruption, at least 4 weeks before starting treatment, throughout the entire duration of treatment, and at least 4 weeks after the end of treatment
- Even if a woman of childbearing potential has amenorrhoea she must follow all the advice on effective contraception
- She should be capable of complying with effective contraceptive measures
- She is informed and understands the potential consequences of pregnancy and the need to rapidly consult her doctor if there is a risk of pregnancy
- She understands the need to commence the treatment as soon as thalidomide is dispensed following a negative pregnancy test
- She understands the need and accepts to undergo pregnancy testing every 4 weeks except in case of confirmed tubal sterilisation
- She acknowledges that she understands the hazards and necessary precautions associated with the use of thalidomide.

As thalidomide is found in semen, as a precaution all male patients taking thalidomide must meet the following conditions:

- He understands the teratogenic risk if engaged in sexual activity with a pregnant woman or a woman of childbearing potential.
- He understands the need for the use of a condom if engaged in sexual activity with a pregnant woman or a woman of childbearing potential not using effective contraception (even if the man has had a vasectomy), during treatment, during dose interruption and for at least 7 days following discontinuation of treatment.
- He understands that if his female partner becomes pregnant whilst he is taking thalidomide or 7 days after he has stopped taking thalidomide, he should inform his treating physician immediately and that it is recommended to refer the female partner to a physician specialised or experienced in teratology for evaluation and advice.

The prescriber must ensure that:

- The patient complies with the conditions of the Pregnancy Prevention Programme including confirmation that she has an adequate level of understanding
- The patient has acknowledged the aforementioned conditions.

### Contraception

Women of childbearing potential must use one effective method of contraception for at least 4 weeks before start of treatment, during treatment, and until at least 4 weeks after thalidomide treatment and even in case of dose interruption unless the patient commits to absolute and continuous abstinence confirmed on a monthly basis. If not established on effective contraception, the patient must be referred preferably to an appropriately trained healthcare professional for contraceptive advice in order that contraception can be initiated.

The following can be considered to be examples of effective methods of contraception:

- Implant
- Levonorgestrel-releasing intrauterine system (IUS)
- Medroxyprogesterone acetate depot
- Tubal sterilisation
- Sexual intercourse with a vasectomised male partner only; vasectomy must be confirmed by two negative semen analyses
- Ovulation inhibitory progesterone-only pills (i.e. desogestrel)

Because of the increased risk of venous thromboembolism in patients with multiple myeloma (MM), combined oral contraceptive pills are not recommended (see section 4.5). If a patient is currently using

combined oral contraception, she should switch to one of the effective methods listed above. The risk of venous thromboembolism continues for 4-6 weeks after discontinuing combined oral contraception.

#### Pregnancy testing

Medically supervised pregnancy tests with a minimum sensitivity of 25 mIU/ml must be performed for women of childbearing potential as outlined below. This requirement includes women of childbearing potential who practice absolute and continuous abstinence.

#### *Prior to starting treatment*

A medically supervised pregnancy test should be performed during the consultation, when thalidomide is prescribed or in the 3 days prior to the visit to the prescriber once the patient had been using effective contraception for at least 4 weeks. The test should ensure the patient is not pregnant when she starts treatment with thalidomide.

#### *Follow-up and end of treatment*

A medically supervised pregnancy test should be repeated every 4 weeks, including 4 weeks after the end of treatment, except in the case of confirmed tubal sterilisation. These pregnancy tests should be performed on the day of the prescribing visit or in the 3 days prior to the visit to the prescriber.

#### Men

As thalidomide is found in semen, as a precaution all male patients must use condoms during treatment, during dose interruption and for at least 7 days following discontinuation of treatment if their partner is pregnant or is of childbearing potential not using effective contraception. Male patients should not donate semen or sperm during treatment (including during dose interruptions) and for at least 7 days following discontinuation of thalidomide.

#### Additional precautions

Patients should be instructed never to give this medicinal product to another person and to return any unused capsules to their pharmacist at the end of treatment.

Patients should not donate blood during treatment (including during dose interruptions) and for at least 7 days following discontinuation of thalidomide.

Healthcare professionals and caregivers should wear disposable gloves when handling the blister or capsule. Women who are pregnant or suspect they may be pregnant should not handle the blister or capsule (see section 6.6).

#### Educational materials, prescribing and dispensing restrictions

In order to assist patients in avoiding foetal exposure to thalidomide, the Marketing Authorisation Holder will provide educational material to healthcare professionals to reinforce the warnings about the teratogenicity of thalidomide, to provide advice on contraception before treatment is started and to provide guidance on the need for pregnancy testing. The prescriber must inform the patient about the expected teratogenic risk and the strict pregnancy prevention measures as specified in the Pregnancy Prevention Programme and provide patients with appropriate patient educational brochure, patient card and/or equivalent tool as agreed with each National Competent Authority. In collaboration with each National Competent Authority, a controlled access programme has been implemented which includes the use of a patient card and/or equivalent tool for prescribing and/or dispensing controls, and the collection of information relating to the indication in order to monitor the off-label use within the national territory. Ideally, pregnancy testing, issuing a prescription and dispensing should occur on the same day. Dispensing of thalidomide to women of childbearing potential should occur within 7 days of the prescription and following a medically supervised negative pregnancy test result. Prescriptions for women of childbearing potential can be for a maximum duration of treatment of 4 weeks according to the approved indications dosing regimens (see section 4.2), and prescriptions for all other patients can be for a maximum duration of 12 weeks.

### Amenorrhoea

The use of thalidomide could be associated with menstrual disorders including amenorrhoea. Amenorrhoea during thalidomide therapy should be assumed to result from pregnancy, until it is medically confirmed that the patient is not pregnant. A clear mechanism by which thalidomide can induce amenorrhoea is not elucidated. The reported events occurred in young (premenopausal) women (median age 36 years) receiving thalidomide for non-multiple myeloma indications, had an onset within 6 months of initiating treatment and reversed upon discontinuation of thalidomide. In documented case reports with hormone evaluation, the event of amenorrhoea was associated with decreased estradiol levels and elevated FSH/LH levels. When provided, antiovary antibodies were negative and prolactin level was within the normal range.

### Cardiovascular disorders

#### Myocardial infarction

Myocardial infarction (MI) has been reported in patients receiving thalidomide, particularly in those with known risk factors. Patients with known risk factors for MI, including prior thrombosis, should be closely monitored and action should be taken to try to minimise all modifiable risk factors (e.g. smoking, hypertension, and hyperlipidaemia).

#### Venous and arterial thromboembolic events

Patients treated with thalidomide have an increased risk of venous thromboembolism (such as deep vein thrombosis and pulmonary embolism) and arterial thromboembolism (such as myocardial infarction and cerebrovascular event) (see section 4.8). The risk appears to be greatest during the first 5 months of therapy. Thromboprophylaxis and dosing/anticoagulation therapy recommendations are provided in section 4.2.

Previous history of thromboembolic events or concomitant administration of erythropoietic agents or other agents such as hormone replacement therapy, may also increase thromboembolic risk in these patients. Therefore, these agents should be used with caution in multiple myeloma patients receiving thalidomide with prednisone and melphalan. Particularly, a haemoglobin concentration above 12g/dl should lead to discontinuation of erythropoietic agents. Action should be taken to try to minimize all modifiable risk factors (e.g. smoking, hypertension and hyperlipidaemia).

Patients and physicians are advised to be observant for the signs and symptoms of thromboembolism. Patients should be instructed to seek medical care if they develop symptoms such as shortness of breath, chest pain, arm or leg swelling.

### Thyroid disorders

Cases of hypothyroidism have been reported. Optimal control of co-morbid conditions influencing thyroid function is recommended before start of treatment. Baseline and ongoing monitoring of thyroid function is recommended.

### Peripheral neuropathy

Peripheral neuropathy is a very common, potentially severe, adverse reaction to treatment with thalidomide that may result in irreversible damage (see section 4.8). In a phase 3 study, the median time to first neuropathy event was 42.3 weeks.

If the patient experiences peripheral neuropathy, follow the dose and schedule modification instruction provided in section 4.2.

Careful monitoring of patients for symptoms of neuropathy is recommended. Symptoms include paraesthesia, dysaesthesia, discomfort, abnormal co-ordination or weakness.

It is recommended that clinical and neurological examinations are performed in patients prior to starting thalidomide therapy, and that routine monitoring is carried out regularly during treatment. Medicinal products known to be associated with neuropathy should be used with caution in patients receiving thalidomide (see section 4.5).

Thalidomide may also potentially aggravate existing neuropathy and should therefore not be used in patients with clinical signs or symptoms of peripheral neuropathy unless the clinical benefits outweigh the risks.

#### Syncope, bradycardia and atrioventricular block

Patients should be monitored for syncope, bradycardia and atrioventricular block; dose reduction or discontinuation may be required.

#### Pulmonary hypertension

Cases of pulmonary hypertension, some fatal, have been reported in patients treated with thalidomide. Patients should be evaluated for signs and symptoms of underlying cardiopulmonary disease prior to initiating and during thalidomide therapy.

#### Haematological disorders

##### Neutropenia

The incidence of neutropenia grade 3 or 4 reported as adverse reactions was higher in multiple myeloma patients receiving MPT (Melphalan, Prednisone, Thalidomide) than in those receiving MP (Melphalan, Prednisone): 42.7 % versus 29.5 % respectively (study IFM 99-06). Adverse reactions from post-marketing experience such as febrile neutropenia and pancytopenia were reported with thalidomide. Patients should be monitored and dose delay, reduction or discontinuation may be required (see section 4.2).

##### Thrombocytopenia

Thrombocytopenia, including grade 3 or 4 adverse reactions, has been reported in multiple myeloma patients receiving MPT. Patients should be monitored and dose delay, reduction or discontinuation may be required (see section 4.2). Patients and physicians are advised to be observant for signs and symptoms of bleeding including petechiae, epistaxis and gastrointestinal haemorrhage, especially in case of concomitant medicinal product prone to inducing bleeding (see sections 4.5 and 4.8).

##### Hepatic disorders

Hepatic disorders, mainly abnormal liver test results, were reported. No specific pattern was identified between hepatocellular and cholestatic abnormalities, with some cases having a mixed presentation. The majority of the reactions occurred within the first 2 months of therapy and resolved spontaneously without treatment after thalidomide discontinuation. Patients should be monitored for liver function, particularly in case of pre-existing liver disorder or concomitant use of medicinal product susceptible to induce liver dysfunction (see section 4.8).

#### Allergic reactions and severe skin reactions

Cases of allergic reactions including angioedema, anaphylactic reaction and severe cutaneous reactions including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported with the use of Thalidomide. Patients should be advised of the signs and symptoms of these reactions by their prescribers and should be told to seek medical attention immediately if they develop these symptoms. Thalidomide interruption or discontinuation should be considered for Grade 2-3 skin rash. Thalidomide must be discontinued for angioedema, anaphylactic reaction, Grade 4 rash, exfoliative or bullous rash, or if SJS, TEN or DRESS is suspected, and should not be resumed following discontinuation for these reactions. (see sections 4.2 and 4.8).

#### Somnolence

It is very common that thalidomide causes somnolence. Patients should be instructed to avoid situations where somnolence may be a problem and to seek medical advice before taking other medicinal products known to cause somnolence. Patients should be monitored and dose reduction may be required.

Patients should be advised as to the possible impairment of mental and/or physical abilities required for the performance of hazardous tasks (see section 4.7).



### Tumour lysis syndrome

The patients at risk of tumour lysis syndrome are those with high tumour burden prior to treatment. These patients should be monitored closely and appropriate precautions taken.

### Infections

Patients should be monitored for severe infections including sepsis and septic shock.

Cases of viral reactivation have been reported in patients receiving thalidomide, including serious cases of herpes zoster or hepatitis B virus (HBV) reactivation.

Some of the cases of herpes zoster reactivation resulted in disseminated herpes zoster, requiring a temporary hold of the treatment with thalidomide and adequate antiviral treatment.

Some of the cases of HBV reactivation progressed to acute hepatic failure and resulted in discontinuation of thalidomide. Hepatitis B virus status should be established before initiating treatment with thalidomide. For patients who test positive for HBV infection, consultation with a physician with expertise in the treatment of hepatitis B is recommended.

Previously infected patients should be closely monitored for signs and symptoms of viral reactivation, including active HBV infection, throughout therapy.

### Progressive multifocal leukoencephalopathy (PML)

Cases of progressive multifocal leukoencephalopathy, including fatal cases, have been reported with thalidomide. PML was reported several months to several years after starting the treatment with thalidomide. Cases have generally been reported in patients taking concomitant dexamethasone or prior treatment with other immunosuppressive chemotherapy. Physicians should monitor patients at regular intervals and should consider PML in the differential diagnosis in patients with new or worsening neurological symptoms, cognitive or behavioural signs or symptoms. Patients should also be advised to inform their partner or caregivers about their treatment, since they may notice symptoms that the patient is not aware of.

The evaluation for PML should be based on neurological examination, magnetic resonance imaging of the brain, and cerebrospinal fluid analysis for JC virus (JCV) DNA by polymerase chain reaction (PCR) or a brain biopsy with testing for JCV. A negative JCV PCR does not exclude PML. Additional follow-up and evaluation may be warranted if no alternative diagnosis can be established.

If PML is suspected, further dosing must be suspended until PML has been excluded. If PML is confirmed, thalidomide must be permanently discontinued.

### Acute myeloid leukaemia (AML) and myelodysplastic syndromes (MDS)

A statistically significant increase of AML and MDS was observed in one clinical study in patients with previously untreated MM receiving the combination of melphalan, prednisone, and thalidomide (MPT). The risk increased over time and was about 2 % after two years and about 4 % after three years. An increased incidence of second primary malignancies (SPM) has also been observed in patients with newly diagnosed MM receiving lenalidomide. Among invasive SPMs, cases of MDS/AML were observed in patients receiving lenalidomide in combination with melphalan or immediately following high dose melphalan and autologous stem cell transplantation.

The benefit achieved with thalidomide and the risk of AML and MDS must be taken into account before initiating treatment with thalidomide in combination with melphalan and prednisone. Physicians should carefully evaluate patients before and during treatment using standard cancer screening and institute treatment as indicated.

### Patients with renal or hepatic impairment

Studies conducted in healthy subjects and patients with multiple myeloma suggest that thalidomide is not influenced to any significant extent by renal or hepatic function (see section 5.2). However, this

has not formally been studied in patients with impaired renal or hepatic function; therefore patients with severe renal or hepatic impairment should be carefully monitored for any adverse events.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Thalidomide is a poor substrate for cytochrome P450 isoenzymes and therefore clinically important interactions with medicinal products that are inhibitors and/or inducers of this enzyme system are unlikely. Non-enzymatic hydrolysis of thalidomide, being the primary clearance mechanism, suggests that the potential for drug-drug interactions with thalidomide is low.

##### Increase of sedative effects of other medicinal products

Thalidomide has sedative properties, thus may enhance the sedation induced by anxiolytics, hypnotics, antipsychotics, H<sub>1</sub> antihistamines, opiate derivatives, barbiturates and alcohol. Caution should be used when thalidomide is given in combination with medicinal products that cause drowsiness.

##### Bradycardic effect

Due to thalidomide's potential to induce bradycardia, caution should be exercised with medicinal products having the same pharmacodynamic effect such as active substances known to induce torsade de pointes, beta blockers or anticholinesterase agents.

##### Medicinal products known to cause peripheral neuropathy

Medicinal products known to be associated with peripheral neuropathy (e.g. vincristine and bortezomib) should be used with caution in patients receiving thalidomide.

##### Hormonal contraceptives

Thalidomide does not interact with hormonal contraceptives. In 10 healthy women, the pharmacokinetic profiles of norethindrone and ethinyl estradiol following administration of a single dose containing 1.0 mg of norethindrone acetate and 0.75 mg of ethinyl estradiol were studied. The results were similar with and without co-administration of thalidomide 200 mg/day to steady-state levels. However, combined hormonal contraceptives are not recommended due to the increased risk of venous thromboembolic disease.

##### Warfarin

Multiple dose administration of 200 mg thalidomide q.d. for 4 days had no effect on the international normalized ratio (INR) in healthy volunteers. However, due to the increased risk of thrombosis in cancer patients, and a potentially accelerated metabolism of warfarin with corticosteroids, close monitoring of INR values is advised during thalidomide-prednisone combination treatment as well as during the first weeks after ending these treatments.

##### Digoxin

Thalidomide does not interact with digoxin. In 18 healthy male volunteers, multiple dose administration of 200 mg thalidomide had no apparent effect on the single dose pharmacokinetics of digoxin. In addition, single dose administration of 0.5 mg digoxin had no apparent effect on thalidomide pharmacokinetics. It is not known whether the effect will be different in multiple myeloma patients.

#### **4.6 Fertility, pregnancy and lactation**

##### Women of childbearing potential/Contraception in males and females

Women of childbearing potential must use one effective method of contraception for at least 4 weeks before start of treatment, during treatment including during dose interruptions, and until at least 4 weeks after thalidomide treatment (see section 4.4). If pregnancy occurs in a woman treated with thalidomide, treatment must be stopped immediately and the patient should be referred to a physician specialised or experienced in teratology for evaluation and advice.

As thalidomide is found in semen, as a precaution all male patients must use condoms during treatment, during dose interruption and for at least 7 days following discontinuation of treatment when

having sexual intercourse with a pregnant woman or with a woman of childbearing potential who is not using effective contraception. This applies even if the man has had a vasectomy.

If pregnancy occurs in a partner of a male patient taking thalidomide, the female partner should be referred to a physician specialised or experienced in teratology for evaluation and advice.

#### Pregnancy

Thalidomide is contraindicated during pregnancy and in women of childbearing potential unless all the conditions of the Pregnancy Prevention Programme are met (see section 4.3)

Thalidomide is a powerful human teratogen, inducing a high frequency (about 30 %) of severe and live-threatening birth defects such as: ectromelia (amelia, phocomelia, hemimelia) of the upper and/or lower extremities, microtia with abnormality of the external acoustic meatus (blind or absent), middle and internal ear lesions (less frequent), ocular lesions (anophthalmia, microphthalmia), congenital heart disease, renal abnormalities. Other less frequent abnormalities have also been described.

#### Breast-feeding

It is unknown whether thalidomide is excreted in human breast milk. Animal studies have shown excretion of thalidomide in breast milk. Therefore breast-feeding should be discontinued during treatment with thalidomide.

#### Fertility

A study in rabbits demonstrated no effect on fertility indices in males or females although testicular degeneration was observed in males.

### **4.7 Effects on ability to drive and use machines**

Thalidomide BMS as per the recommended posology has minor or moderate influence on the ability to drive and use machines.

Thalidomide may cause fatigue (very common), dizziness (very common), somnolence (very common) and blurred vision (common) (see section 4.8). Patients should be instructed not to drive cars, use machines or perform hazardous tasks while being treated with thalidomide if they feel tired, dizzy, sleepy or have blurred vision.

### **4.8 Undesirable effects**

#### Summary of the safety profile

Most patients taking thalidomide can be expected to experience adverse reactions.

The most commonly observed adverse reactions associated with the use of thalidomide in combination with melphalan and prednisone are: neutropenia, leukopenia, constipation, somnolence, paraesthesia, peripheral neuropathy, anaemia, lymphopenia, thrombocytopenia, dizziness, dysaesthesia, tremor and peripheral oedema.

In addition to the adverse reactions outlined above, thalidomide in combination with dexamethasone in other clinical studies led to the very common adverse reaction of fatigue; common adverse reactions of transient ischaemic event, syncope, vertigo, hypotension, mood altered, anxiety, blurred vision, nausea and dyspepsia; and uncommon adverse reactions of cerebrovascular accident, diverticular perforation, peritonitis, orthostatic hypotension and bronchitis.

The most clinically important adverse reactions associated with the use of thalidomide in combination with melphalan and prednisone or dexamethasone include: deep vein thrombosis and pulmonary embolism, peripheral neuropathy, severe skin reactions including Stevens-Johnson syndrome, toxic epidermal necrolysis and drug reaction with eosinophilia and systemic symptoms, syncope, bradycardia, and dizziness (see sections 4.2, 4.4 and 4.5).

#### Tabulated list of adverse reactions

Table 3 contains only the adverse reactions for which a causal relationship with medicinal product treatment could reasonably be established observed in the pivotal study and from post-marketing

experience. Frequencies given are based on the observations during a pivotal comparative clinical study investigating the effect of thalidomide in combination with melphalan and prednisone in previously untreated multiple myeloma patients.

Frequencies are defined as: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1000$ ); very rare ( $< 1/10,000$ ) and not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

**Table 3: Adverse drug reactions (ADRs) reported in pivotal clinical study with thalidomide in combination with melphalan and prednisone and from post marketing use**

System Organ Class	Frequency	Adverse reaction
<b>Infections and infestations</b>	<u>Common</u>	Pneumonia
	<u>Not Known</u>	Severe infections (e.g. fatal sepsis including septic shock) <sup>†</sup> , Viral infections, including herpes zoster and hepatitis B virus reactivation <sup>†</sup>
<b>Neoplasms benign, malignant and unspecified (incl cysts and polyps)</b>	<u>Common</u>	Acute myeloid leukaemia* <sup>^</sup>
	<u>Uncommon</u>	Myelodysplastic syndrome* <sup>^</sup>
	<u>Not Known</u>	Tumour lysis syndrome <sup>†</sup>
<b>Blood and lymphatic system disorders</b>	<u>Very Common</u>	Neutropenia, Leukopenia, Anaemia, Lymphopenia, Thrombocytopenia
	<u>Common</u>	Febrile neutropenia <sup>†</sup> , Pancytopenia <sup>†</sup>
<b>Immune System Disorders</b>	<u>Not Known</u>	Allergic reactions (hypersensitivity, angioedema, anaphylactic reaction, urticaria) <sup>†</sup>
<b>Endocrine Disorders</b>	<u>Not Known</u>	Hypothyroidism <sup>†</sup>
<b>Psychiatric disorders</b>	<u>Common</u>	Confusional state, Depression
<b>Nervous system disorders</b>	<u>Very Common</u>	Peripheral neuropathy*, Tremor, Dizziness, Paraesthesia, Dysaesthesia, Somnolence
	<u>Common</u>	Convulsions <sup>†</sup> , Abnormal coordination
	<u>Not Known</u>	Posterior reversible encephalopathy syndrome (PRES)* <sup>†</sup> , Worsening of Parkinson's disease symptoms <sup>†</sup>
<b>Ear and labyrinth disorders</b>	<u>Common</u>	Hearing impaired or deafness <sup>†</sup>
<b>Cardiac disorders</b>	<u>Common</u>	Cardiac failure, Bradycardia
	<u>Uncommon</u>	Myocardial infarction <sup>†</sup> , Atrial fibrillation <sup>†</sup> , Atrioventricular block <sup>†</sup>
<b>Vascular disorders</b>	<u>Common</u>	Deep vein thrombosis*
<b>Respiratory, thoracic and mediastinal disorders</b>	<u>Common</u>	Pulmonary embolism*, Interstitial lung disease, Bronchopneumopathy, Dyspnea
	<u>Not Known</u>	Pulmonary hypertension <sup>†</sup>
<b>Gastrointestinal disorders</b>	<u>Very Common</u>	Constipation
	<u>Common</u>	Vomiting, Dry mouth
	<u>Uncommon</u>	Intestinal obstruction <sup>†</sup>
	<u>Not Known</u>	Gastrointestinal perforation <sup>†</sup> , Pancreatitis <sup>†</sup> , Gastrointestinal haemorrhage <sup>†</sup>
<b>Hepatobiliary disorders</b>	<u>Not Known</u>	Hepatic disorders <sup>†</sup>
	<u>Common</u>	Toxic skin eruption, Rash, Dry skin

System Organ Class	Frequency	Adverse reaction
<b>Skin and subcutaneous tissue disorders</b>	<u>Not Known</u>	Stevens-Johnson syndrome <sup>*,†</sup> , Toxic epidermal necrolysis <sup>*,†</sup> , Drug reaction with eosinophilia and systemic symptoms <sup>*,†</sup> , Leukocytoclastic vasculitis <sup>†</sup>
<b>Renal and urinary disorders</b>	<u>Common</u>	Renal failure <sup>†</sup>
<b>Reproductive System and Breast Disorders</b>	<u>Not Known</u>	Sexual dysfunction <sup>†</sup> , Menstrual disorders including amenorrhea <sup>†</sup>
<b>General disorders and administration site conditions</b>	<u>Very Common</u>	Peripheral oedema
	<u>Common</u>	Pyrexia, Asthenia, Malaise

\* see section 4.8 description of selected adverse reactions

† identified from post marketing data

^ Acute myeloid leukaemia and Myelodysplastic syndrome were reported in one clinical study in patients with previously untreated MM receiving the combination of melphalan, prednisone and thalidomide (MPT)

### Description of selected adverse reactions

#### *Blood and lymphatic system disorders*

Adverse reactions for haematological disorders are provided compared to the comparator arm, as the comparator has a significant effect on these disorders (Table 4).

**Table 4: Comparison of haematological disorders for the melphalan, prednisone (MP) and melphalan, prednisone, thalidomide (MPT) combinations in study IFM 99-06 (see section 5.1)**

	n (% of patients)	
	MP (n=193)	MPT (n=124)
	Grades 3 and 4*	
<b>Neutropenia</b>	57 (29.5)	53 (42.7)
<b>Leukopenia</b>	32 (16.6)	32 (25.8)
<b>Anaemia</b>	28 (14.5)	17 (13.7)
<b>Lymphopenia</b>	14 (7.3)	15 (12.1)
<b>Thrombocytopenia</b>	19 (9.8)	14 (11.3)

\* WHO Criteria

Additional adverse reactions from post-marketing experience with thalidomide and not seen in the pivotal study include febrile neutropenia and pancytopenia.

#### *Teratogenicity*

The risk of intra-uterine death or severe birth defects, primarily phocomelia, is extremely high. Thalidomide must not be used at any time during pregnancy (see sections 4.4 and 4.6).

#### *Venous and arterial thromboembolic events*

An increased risk of venous thromboembolism (such as deep vein thrombosis and pulmonary embolism) and arterial thromboembolism (such as myocardial infarction and cerebrovascular event) has been reported in patients treated with thalidomide (see section 4.4).

#### *Peripheral neuropathy*

Peripheral neuropathy is a very common, potentially severe, adverse reaction of treatment with thalidomide that may result in irreversible damage (see section 4.4). Peripheral neuropathy generally occurs following chronic use over a period of months. However, reports following relatively short-term use also exist. Incidence of neuropathy events leading to discontinuation, dose reduction or interruption increases with cumulative dose and duration of therapy. Symptoms may occur some time after thalidomide treatment has been stopped and may resolve slowly or not at all.

### *Posterior reversible encephalopathy syndrome (PRES)/ Reversible posterior leukoencephalopathy syndrome (RPLS)*

Cases of PRES/RPLS have been reported. Signs and symptoms included visual disturbance, headache, seizures and altered mental status, with or without associated hypertension. A diagnosis of PRES/RPLS requires confirmation by brain imaging. The majority of the reported cases had recognized risk factors for PRES/RPLS, including hypertension, renal impairment and concomitant use of high dose corticosteroids and/or chemotherapy.

### *Acute myeloid leukaemia (AML) and myelodysplastic syndromes (MDS)*

AML and MDS were reported in one clinical study in patients with previously untreated multiple myeloma receiving the combination of melphalan, prednisone, and thalidomide (see section 4.4).

### *Allergic reactions and severe skin reactions*

Cases of allergic reactions including angioedema, anaphylactic reaction and severe cutaneous reactions including Stevens-Johnson syndrome, TEN and DRESS have been reported with the use of thalidomide therapy. If angioedema, anaphylactic reaction, Stevens-Johnson syndrome, TEN or DRESS is suspected, use of thalidomide should not be resumed (see section 4.2 and 4.4).

### Elderly population

The adverse reaction profile reported in patients > 75 years of age treated with thalidomide 100 mg once daily was similar to the adverse reaction profile observed in patients ≤ 75 years of age treated with thalidomide 200 mg once daily (see Table 3). However, patients with age > 75 years are potentially at risk for a higher frequency of serious adverse reactions.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via [the national reporting system listed in Appendix V](#).

## **4.9 Overdose**

Eighteen cases of overdose have been reported in the literature concerning doses up to 14.4 grams. In thirteen of these cases, patients took thalidomide alone; amounts ranged from 350 mg to 4000 mg. These patients either exhibited no symptoms or exhibited symptoms of drowsiness, irritability, "sickness," and/or headache. In one 2-year-old child who took 700 mg, there was an abnormal plantar response in addition to drowsiness and irritability. No fatalities have been reported and all overdose patients recovered without sequelae. There is no specific antidote for a thalidomide overdose. In the event of an overdose, the patient's vital signs should be monitored and appropriate supportive care given to maintain blood pressure and respiratory status.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: immunosuppressants, other immunosuppressants, ATC code: L04AX02.

Thalidomide has a chiral centre and is used clinically as a racemate of (+)-(R)- and (-)-(S)-thalidomide. The spectrum of activity of thalidomide is not fully characterised.

### Mechanism of action

Thalidomide shows immunomodulatory, anti-inflammatory and potential anti-neoplastic activities. Data from *in vitro* studies and clinical trials suggest that the immunomodulatory, anti-inflammatory and anti-neoplastic effects of thalidomide may be related to suppression of excessive tumour necrosis factor-alpha (TNF- $\alpha$ ) production, down-modulation of selected cell surface adhesion molecules


involved in leukocyte migration and anti-angiogenic activity. Thalidomide is also a non-barbiturate centrally active hypnotic sedative. It has no antibacterial effects.

### Clinical efficacy and safety

Results from IFM 99-06, a Phase 3, randomised, open label, parallel group, multicentre study have demonstrated a survival advantage when thalidomide is used in combination with melphalan and prednisone for 12 cycles of 6 weeks in the treatment of newly diagnosed multiple myeloma patients. In this study the age range of patients was 65-75 years, with 41 % (183/447) of patients 70 years old or older. The median dose of thalidomide was 217 mg and > 40 % of patients received 9 cycles. Melphalan and prednisone were dosed at 0.25 mg/kg/day and 2 mg/kg/day respectively on days 1 to 4 of each 6 weeks cycle.

Further to the per protocol analysis, an update was conducted for the IFM 99-06 study providing an additional 15 months follow-up data. The median overall survival (OS) was  $51.6 \pm 4.5$  and  $33.2 \pm 3.2$  months in the MPT and MP groups, respectively (97.5 % CI 0.42 to 0.84). This 18 month difference was statistically significant with a hazard ratio of reduction of risk of death in the MPT arm of 0.59, 97.5 % confidence interval of 0.42-0.84 and p-value of < 0.001 (see Figure 1).

**Figure 1: Overall survival according to treatment**



### Paediatric Population

The European Medicines Agency has waived the obligation to submit the results of studies with thalidomide in all subsets of the paediatric population in multiple myeloma (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

### Absorption

Absorption of thalidomide is slow after oral administration. The maximum plasma concentrations are reached 1-5 hours after administration. Co-administration of food delayed absorption but did not alter the overall extent of absorption.

### Distribution

The plasma protein binding of the (+)-(R) and (-)-(S) enantiomers was found to be 55 % and 65 % respectively. Thalidomide is present in the semen of male patients at levels similar to plasma concentrations (see section 4.4). The distribution of thalidomide is not influenced by age, gender, renal function and blood chemistry variables, to any significant level.

### Biotransformation

Thalidomide is metabolised almost exclusively by non-enzymatic hydrolysis. In plasma, unchanged thalidomide represents 80 % of the circulatory components. Unchanged thalidomide was a minor component (< 3 % of the dose) in urine. In addition to thalidomide, hydrolytic products N-(o-carboxybenzoyl) glutarimide and phthaloyl isoglutamine formed via non-enzymatic processes are also present in plasma and in majority in urine. Oxidative metabolism does not contribute significantly to the overall metabolism of thalidomide. There is minimal cytochrome P450 catalysed hepatic metabolism of thalidomide. There are *in vitro* data indicating that prednisone may give rise to enzyme induction which could reduce the systemic exposure of concomitantly used medicinal products. The *in vivo* relevance of these findings is unknown.

### Elimination

The mean elimination half-life of thalidomide in plasma following single oral doses between 50 mg and 400 mg was 5.5 to 7.3 hours. Following a single oral dose of 400 mg of radio-labelled thalidomide, the total mean recovery was 93.6 % of the administered dose by day 8. The majority of the radioactive dose was excreted within 48 hour following dose administration. The major route of excretion was via the urine (> 90 %) while faecal excretion was minor.

There is a linear relationship between body weight and estimated thalidomide clearance; in multiple myeloma patients with body weight from 47-133 kg, thalidomide clearance ranged from approximately 6-12 L/h, representing an increase in thalidomide clearance of 0.621 L/h per 10 kg body weight increase.

### Linearity/non-linearity

Total systemic exposure (AUC) is proportional to dose at single-dose conditions. No time dependency of the pharmacokinetics has been observed.

### Hepatic and renal impairment

The extent of thalidomide metabolism by the liver cytochrome P450 system is minimal and intact thalidomide is not excreted by the kidney. Measures of renal function (CrCl) and liver function (blood chemistry) indicate minimal effect of kidney and liver function on the pharmacokinetics of thalidomide. As such the metabolism of thalidomide is not expected to be affected by hepatic or renal dysfunction. Data from patients with end-stage renal disease suggest no impact of kidney function on thalidomide pharmacokinetics.

## **5.3 Preclinical safety data**

In the male dog, after one year of dosing, reversible bile plugs in canaliculi were observed at exposures greater than 1.9-fold the human exposure.

Decreased platelet counts were noted in the mouse and rat studies. The latter appears to be related to thalidomide and occurred at exposures greater than 2.4-fold the human exposure. This decrease did not result in clinical signs.

In a one-year dog study, enlarged and/or blue discoloration of mammary glands and prolonged estrus were observed in females at exposures equal to 1.8 or greater than 3.6-fold the human exposure, respectively. The relevance to humans is unknown.

The effect of thalidomide on thyroid function was assessed in both rats and dogs. No effects were observed in dogs; however in rats, there was an apparent dose-dependent decrease in total and free T4 that was more consistent in the female.



No mutagenic or genotoxic effect has been revealed when thalidomide was assayed in a standard battery of genotoxicity tests. No evidence of carcinogenicity was observed at exposures approximately 15, 13 and 39 times the estimated clinical AUC at the recommended starting dose in mice, male rats and female rats respectively.

Animal studies have demonstrated differences in species susceptibility to the teratogenic effects of thalidomide. In humans, thalidomide is a proven teratogen.

A study in rabbits demonstrated no effect on fertility indices in males or females although testicular degeneration was observed in males.

A peri- and postnatal toxicity study performed in rabbits with thalidomide administered at doses up to 500 mg/kg/day resulted in abortions, increased stillbirths and decreased pup viability during lactation. Pups from mothers treated with thalidomide had increased abortions, reduced body weight gain, alterations in learning and memory, decreased fertility, and reduced pregnancy index.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Capsule contents

Starch, pregelatinised  
Magnesium stearate

#### Capsule shell

Gelatin  
Titanium dioxide (E171)

#### Printing ink

Shellac  
Black iron oxide (E172)  
Propylene glycol

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

5 years

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

PVC/PCTFE/aluminium blister containing 14 capsules.

Pack sizes: 28 capsules (two blisters) in a wallet card.

### **6.6 Special precautions for disposal and other handling**

Capsules should not be opened or crushed. If powder from thalidomide makes contact with the skin, the skin should be washed immediately and thoroughly with soap and water. If thalidomide makes

contact with the mucous membranes, they should be thoroughly flushed with water.

Healthcare professionals and caregivers should wear disposable gloves when handling the blister or capsule. Gloves should then be removed carefully to prevent skin exposure, placed in a sealable plastic polyethylene bag and disposed of in accordance with local requirements. Hands should then be washed thoroughly with soap and water. Women who are pregnant or suspect they may be pregnant should not handle the blister or capsule (see section 4.4).

All unused capsules should be returned to the pharmacist at the end of treatment.

## **7. MARKETING AUTHORISATION HOLDER**

Bristol-Myers Squibb Pharma EEIG  
Plaza 254  
Blanchardstown Corporate Park 2  
Dublin 15, D15 T867  
Ireland

## **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/08/443/001

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 16 April 2008

Date of latest renewal: 08 February 2018

## **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURER (S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer(s) responsible for batch release

Celgene Distribution B.V.  
Orteliuslaan 1000  
3528 BD Utrecht  
Netherlands

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

### **• Periodic safety update reports (PSURs)**

The requirements for submission of periodic safety update reports for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

### **• Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency.
  - Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.
- **Additional risk minimisation measures**
1. The MAH shall agree the details of a controlled access programme with the National Competent Authorities and must implement such programme nationally to ensure that:
    - Prior to launch, all doctors who intend to prescribe Thalidomide BMS and all pharmacists who may dispense Thalidomide BMS receive a Direct Healthcare Professional Communication as described below.
    - Prior to prescribing (where appropriate, and in agreement with the National Competent Authority, dispensing) all healthcare professionals who intend to prescribe (and dispense) Thalidomide BMS are provided with an Educational Healthcare Professional's Kit containing the following:
      - Educational Healthcare Professional brochure
      - Educational brochures for patients
      - Patient card
      - Risk awareness forms
      - Information on where to find latest Summary of Product Characteristics (SmPC)

2. The MAH shall implement a pregnancy prevention programme (PPP) in each Member State. Details of the PPP should be agreed with the National Competent Authorities in each Member State and put in place prior to the launch of the medicinal product.
3. The MAH should agree the final text of the Direct Healthcare Professional Communication and the contents of the Educational Healthcare Professional's Kit with the National Competent Authority in each Member State prior to launch of the medicinal product and ensure that the materials contain the key elements as described below.
4. The MAH should agree on the implementation of the controlled access programme in each Member State.
5. Prior to approval by the National Competent Authority and prior to launch of the medicinal product, the MAH should ensure that the educational materials are provided to and reviewed by the national patients' organisations or if such an organisation does not exist or can not be involved, by a relevant patients group. Patients involved should be preferably naïve to the history of thalidomide. Results of the user testing will have to be provided to the National Competent Authority and final materials validated at a national level.
6. The MAH should also agree with each Member State prior to launch of the medicinal product:
  - The most appropriate strategies to monitor the off-label use within national territories
  - The collection of detailed data to understand demographics of target population, indication and number of women of child bearing potential in order to monitor the off-label use within national territory.
7. The MAH shall notify the EMA and the appropriate national patients and victims representatives of the proposed launch date before launch in each Member State.

### **Key elements to be included**

#### **Direct Healthcare Professional Communication (prior to launch)**

The Direct Healthcare Professional Communication shall consist of two parts:

- A core text as agreed by the CHMP
- National specific requirements agreed with the National Competent Authority regarding:
  - Distribution of the medicinal product
  - Procedures to ensure that all appropriate measures have been performed prior to Thalidomide BMS being dispensed

#### **Educational Healthcare Professional's Kit**

The Educational Healthcare Professional's Kit shall contain the following elements:

#### **Educational Healthcare Professional brochure**

- History and background of thalidomide
- Maximum duration of treatment prescribed
  - 4 weeks for women with childbearing potential
  - 12 weeks for men and women without childbearing potential
- Teratogenicity and the need to avoid foetal exposure
- Guidance on handling the blister or capsule of Thalidomide BMS for healthcare professionals and caregivers
- Obligations of the healthcare professionals who intend to prescribe or dispense Thalidomide BMS
  - Need to provide comprehensive advice and counselling to patients
  - That patients should be capable of complying with the requirements for the safe use of Thalidomide BMS
  - Need to provide patients with the appropriate patient educational brochure, patient card and/or equivalent tool
- Safety advice relevant to all patients
  - Description and management of ischaemic heart disease (including myocardial infarction)

- Local country specific arrangements for a prescription for thalidomide to be dispensed
- That any unused capsules should be returned to the pharmacist at the end of the treatment
- That the patient should not donate blood during treatment (including during dose interruptions) and for at least 7 days following discontinuation of Thalidomide BMS
- Description of the PPP and categorisation of patients based on sex and childbearing potential
  - Algorithm for implementation of PPP
  - Definition of women of childbearing potential (WCBP) and actions the prescriber should take if unsure
- Safety advice for women of childbearing potential
  - The need to avoid foetal exposure
  - Description of the PPP
  - Need for effective contraception (even if the woman has amenorrhoea) and definition of effective contraception
  - That if she needs to change or stop using her method of contraception she should inform:
    - The physician prescribing her contraception that she is on thalidomide
    - The physician prescribing thalidomide that she has stopped or changed her method of contraception
  - Pregnancy test regime
    - Advice on suitable tests
    - Before commencing treatment
    - During treatment based on method of contraception
    - After finishing treatment
  - Need to stop Thalidomide BMS immediately upon suspicion of pregnancy
  - Need to tell treating doctor immediately upon suspicion of pregnancy
- Safety advice for men
  - The need to avoid foetal exposure
  - The need to use condoms if sexual partner is pregnant or a WCBP not using effective contraception (even if the man has had a vasectomy)
    - During Thalidomide BMS treatment
    - For at least 7 days following final dose
  - That he should not donate semen or sperm during treatment (including during dose interruptions) and for at least 7 days following discontinuation of Thalidomide BMS treatment
  - That if his partner becomes pregnant whilst he is taking Thalidomide BMS or shortly after he has stopped taking Thalidomide BMS he should inform his treating doctor immediately
- Requirements in the event of pregnancy
  - Instructions to stop Thalidomide BMS immediately upon suspicion of pregnancy, if female patient
  - Need to refer patient to physician specialised or experienced in dealing with teratology and its diagnosis for evaluation and advice
  - Local contact details for reporting of any suspected pregnancy immediately
- Local contact details for reporting adverse reactions

### **Educational Brochures for patients**

The Educational brochures for patients should be of 3 types:

- Brochure for women of childbearing potential and their partner
- Brochure for women patients who are not of childbearing potential
- Brochure for male patients

All educational brochures for patients should contain the following elements:

- That thalidomide is teratogenic
- That thalidomide may cause ischaemic heart disease, (including myocardial infarction)

- Description of the patient card and its necessity
- Guidance on handling Thalidomide BMS for patients, caregivers and family members
- National or other applicable specific arrangements for a prescription for Thalidomide BMS to be dispensed
- That the patient must not give Thalidomide BMS to any other person
- That the patient should not donate blood during treatment (including during dose interruptions) and for at least 7 days after discontinuation of Thalidomide BMS treatment
- That the patient should tell their doctor about any adverse events
- That any unused capsules should be returned to the pharmacist at the end of the treatment

The following information should also be provided in the appropriate brochure:

#### Brochure for women patients with childbearing potential

- The need to avoid foetal exposure
- Description of the PPP
- The need for effective contraception and definition of effective contraception
- That if she needs to change or stop using her method of contraception she should inform:
  - The physician prescribing her contraception that she is on thalidomide
  - The physician prescribing thalidomide that she has stopped or changed her method of contraception
- Pregnancy test regime
  - Before commencing treatment
  - During treatment (including dose interruptions), at least every 4 weeks except in case of confirmed tubal sterilisation
  - After finishing treatment
- The need to stop Thalidomide BMS immediately upon suspicion of pregnancy
- The need to contact their doctor immediately upon suspicion of pregnancy

#### Brochure for male patients

- The need to avoid foetal exposure
- The need to use condoms if sexual partner is pregnant or a WCBP not using effective contraception (even if the man has had vasectomy)
  - During Thalidomide BMS treatment (including dose interruptions)
  - For at least 7 days following final dose
- That if his partner becomes pregnant, he should inform his treating doctor immediately
- That he should not donate semen or sperm during treatment (including during dose interruptions) and for at least 7 days following discontinuation of Thalidomide BMS treatment

#### Patient Card or equivalent tool

The patient card shall contain the following elements:

- Verification that appropriate counselling has taken place
- Documentation of childbearing potential status
- Check box (or similar) which physician ticks to confirm that patient is using effective contraception (if woman of childbearing potential)
- Pregnancy test dates and results

#### Risk Awareness Forms

There should be 3 types of risk awareness forms:

- Women of childbearing potential
- Women of non-childbearing potential
- Male patient

All risk awareness forms should contain the following elements:

- teratogenicity warning

- patients receive the appropriate counselling prior to treatment initiation
- affirmation of patient understanding of the risk of thalidomide and the PPP measures
- date of counselling
- patient details, signature and date
- prescriber name, signature and date
- aim of this document i.e. as stated in the PPP: “The aim of the risk awareness form is to protect patients and any possible foetuses by ensuring that patients are fully informed of and understand the risk of teratogenicity and other adverse reactions associated with the use of thalidomide. It is not a contract and does not absolve anybody from his/her responsibilities with regard to the safe use of the product and prevention of foetal exposure.”

Risk awareness forms for women of childbearing potential should also include:

- Confirmation that the physician has discussed the following:
  - the need to avoid foetal exposure
  - that if she is pregnant or plans to be, she must not take thalidomide
  - that she understands the need to avoid thalidomide during pregnancy and to apply effective contraceptive measures without interruption, at least 4 weeks before starting treatment, throughout the entire duration of treatment, and at least 4 weeks after the end of treatment
  - that if she needs to change or stop using her method of contraception she should inform:
    - the physician prescribing her contraception that she is taking Thalidomide BMS
    - the physician prescribing Thalidomide BMS that she has stopped or changed her method of contraception
  - of the need for pregnancy tests i.e. before treatment, at least every 4 weeks during treatment and after treatment
  - of the need to stop Thalidomide BMS immediately upon suspicion of pregnancy
  - of the need to contact their doctor immediately upon suspicion of pregnancy
  - that she should not share the medicinal product with any other person
  - that she should not donate blood during treatment (including during dose interruptions) and for at least 7 days following discontinuation of Thalidomide BMS
  - that she should return the unused capsules to the pharmacist at the end of treatment

Risk awareness forms for women with no childbearing potential should also include:

- Confirmation that the physician has discussed the following:
  - that she should not share the medicinal product with any other person
  - that she should not donate blood during treatment (including during dose interruptions) and for at least 7 days following discontinuation of Thalidomide BMS
  - that she should return the unused capsules to the pharmacist at the end of treatment

Risk awareness forms for male patients should also include:

- Confirmation that the physician has discussed the following:
  - the need to avoid foetal exposure
  - that thalidomide is found in semen and the need to use condoms if sexual partner is pregnant or is a WCBP not on effective contraception (even if the man has had vasectomy)
  - that if his partner becomes pregnant, he should inform his treating doctor immediately and always use a condom
  - that he should not share the medicinal product with any other person



- that he should not donate blood or semen during treatment (including during dose interruptions) and for at least 7 days following discontinuation of Thalidomide BMS
- that he should return the unused capsules to the pharmacist at the end of treatment

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**WALLET CARD**

**1. NAME OF THE MEDICINAL PRODUCT**

Thalidomide BMS 50 mg hard capsules  
thalidomide

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each capsule contains 50 mg of thalidomide.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

28 hard capsules

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For oral use.  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

WARNING: Thalidomide causes birth defects and foetal death. Do not use while pregnant or breastfeeding.

You must follow the Thalidomide BMS Pregnancy Prevention Programme.

Keep the package intact.

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

Unused medicinal product should be returned to your pharmacist.

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Bristol-Myers Squibb Pharma EEIG  
Plaza 254  
Blanchardstown Corporate Park 2  
Dublin 15, D15 T867  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/08/443/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Thalidomide BMS 50 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2 D bar code carrying the unique identifier

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

Thalidomide BMS 50 mg  
thalidomide

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

Bristol-Myers Squibb Pharma EEIG

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the patient

### Thalidomide BMS 50 mg hard capsules thalidomide

#### WARNING

**Thalidomide causes birth defects and foetal death. Do not take thalidomide if you are pregnant or could become pregnant. You must follow the contraception advice given to you by your doctor.**

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Thalidomide BMS is and what it is used for
2. What you need to know before you take Thalidomide BMS
3. How to take Thalidomide BMS
4. Possible side effects
5. How to store Thalidomide BMS
6. Contents of the pack and other information

#### 1. What Thalidomide BMS is and what it is used for

##### What Thalidomide BMS is

Thalidomide BMS contains an active substance called thalidomide. This belongs to a group of medicines which affect how your immune system works.

##### What Thalidomide BMS is used for

Thalidomide BMS is used with two other medicines called ‘melphalan’ and ‘prednisone’ to treat adults with a type of cancer called multiple myeloma. It is used in people who have recently been diagnosed and who have not been prescribed another medicine for their multiple myeloma before who are aged 65 years and over, or aged less than 65 years who cannot be treated with high dose chemotherapy, which can be very difficult for the body to handle.

##### What is multiple myeloma

Multiple myeloma is a type of cancer which affects a certain type of white blood cell, called the plasma cell. These cells collect in the bone marrow and divide out of control. This can damage the bone and kidneys. Multiple myeloma generally cannot be cured. However, the signs and symptoms can be greatly reduced or disappear for a period of time. This is called a ‘remission’.

##### How Thalidomide BMS works

Thalidomide BMS works by helping the body’s immune system and directly attacking the cancer. It works in a number of different ways:

- by stopping the cancer cells developing
- by stopping blood vessels growing in the cancer
- by stimulating part of the immune system to attack the cancer cells.



## 2. What you need to know before you take Thalidomide BMS

You will have been given specific instructions by your doctor, particularly on the effects of thalidomide on unborn babies (outlined in the Thalidomide BMS Pregnancy Prevention Programme).

You will have been given an educational brochure for patient by your doctor. Read it carefully and follow the related instructions.

If you do not fully understand these instructions, please ask your doctor to explain them again before you take thalidomide. See also further information in this section under “Warnings and precautions” and “Pregnancy and breast-feeding”.

### Do not take Thalidomide BMS

- if you are pregnant or think you may be pregnant or are planning to become pregnant, **as Thalidomide BMS causes birth defects and foetal death.**
- if you are able to become pregnant, unless you are able to follow or comply with the required contraceptive measures to prevent you from becoming pregnant (see section 2 “Warnings and precautions” and “Pregnancy and breast-feeding”).
- if you are able to become pregnant, your doctor will record with each prescription that the necessary measures have been taken and will provide you with this confirmation.
- if you are allergic to thalidomide or any of the other ingredients of this medicine listed in section 6 “Contents of the pack and other information”.

Do not take Thalidomide BMS if any of the above applies to you. If you are not sure, talk to your doctor or pharmacist before taking Thalidomide BMS.

### Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking this medicine in the following situations:

#### For women taking Thalidomide BMS

Before starting the treatment, you should ask your doctor if you are able to become pregnant, even if you think this is unlikely. Even if you do not have a menstrual bleeding following cancer therapy, you may become pregnant.

If you are able to become pregnant:

- Your doctor will make sure that you have pregnancy tests
  - before treatment
  - every 4 weeks during treatment
  - 4 weeks after stopping treatment
- You must use one effective method of contraception:
  - for at least 4 weeks before starting treatment
  - during treatment
  - until at least 4 weeks after stopping treatment

Your doctor will tell you what method of contraception to use.

If you are able to become pregnant, your doctor will record with each prescription that the necessary measures, as outlined above, have been taken

#### For men taking Thalidomide BMS

Thalidomide passes into semen. Therefore, do not have unprotected intercourse, even if you had a vasectomy.

- Pregnancy and any exposure during pregnancy must be avoided. Always use a condom:
  - during treatment
  - for at least 7 days after stopping treatment
- You must not donate semen:
  - during treatment
  - for at least 7 days after stopping treatment

## **For all patients**

Talk to your doctor before taking Thalidomide BMS if

- you do not understand the contraception advice given to you by your doctor or if you do not feel able to follow this advice.
- you have had a heart attack, have ever had a blood clot in the past, or if you smoke, have high blood pressure or high cholesterol levels. During the treatment with Thalidomide BMS you have an increased risk of developing blood clots in the veins and arteries (see also section 4 “Possible side effects”).
- you have experienced or have existing neuropathy i.e. nerve damage causing tingling, abnormal co-ordination or pain in your hands or feet (see also section 4 “Possible side effects”).
- you experienced or have existing slow heart rate (this may be a symptom of bradycardia).
- you have high blood pressure in the arteries of the lungs (see also section 4 “Possible side effects”).
- you have a fall in the number of white blood cells (neutropenia) accompanied by fever and infection.
- you have a fall in the number of platelets. You will be more prone to bleeding and bruising.
- you have or have had injury to the liver (hepatic disorders) including abnormal liver test results.
- you experience or have experienced in the past severe skin reactions called Stevens-Johnson syndrome, toxic epidermal necrolysis or drug reaction with eosinophilia and systemic symptoms (which is also known as DRESS or drug hypersensitivity syndrome). (For description of symptoms see section 4 “Possible side effects”).
- you have had an allergic reaction whilst taking Thalidomide BMS such as rash, itching, swelling, dizziness or trouble breathing.
- you have experienced sleepiness.
- you have experienced fever, chills and severe shaking, and possibly complicated by low blood pressure and confusion (these may be symptoms of severe infections).
- you have or have ever had previous viral infection, particularly varicella zoster, hepatitis B infection, or HIV. If you are in doubt, talk to your doctor. Treatment with Thalidomide BMS may cause a virus to become active again in patients who carry it, resulting in a recurrence of the infection. Your doctor should check whether you have ever had hepatitis B infection.
- you have kidney or liver problems (see also section 4 “Possible side effects”).

Your thyroid function may be checked before you take thalidomide and monitored during treatment.

At any time during or after your treatment, tell your doctor or nurse immediately if you experience: blurred, loss of or double vision, difficulty speaking, weakness in an arm or a leg, a change in the way you walk or problems with your balance, persistent numbness, decreased sensation or loss of sensation, memory loss or confusion. These may all be symptoms of a serious and potentially fatal brain condition known as progressive multifocal leukoencephalopathy (PML). If you had these symptoms prior to treatment with Thalidomide BMS, tell your doctor about any change in these symptoms.

Your doctor may check if you have a high total amount of tumour throughout the body, including your bone marrow. This could lead to a condition where the tumours break down and cause unusual levels of chemicals in the body which can lead to kidneys failure (this condition is called Tumour Lysis Syndrome) (see also section 4 “Possible side effects”).

Your doctor should evaluate if you develop additional types of haematological malignancies (called acute myeloid leukaemia and myelodysplastic syndromes) during your treatment with Thalidomide BMS (see also section 4 “Possible side effects”).

You must not donate blood during Thalidomide BMS treatment and for at least 7 days after stopping treatment.

If you are not sure if any of the above apply to you, talk to your doctor before taking Thalidomide BMS.

### **Children and adolescents**

Thalidomide BMS is not recommended for use in children and young people under 18 years.

### **Other medicines and Thalidomide BMS**

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines obtained without a prescription, including herbal medicines.

Make sure you tell your doctor if you are taking any medicines which:

- cause sleepiness as thalidomide may increase their effects. This includes sedatives (such as anxiolytics, hypnotics, antipsychotics, H<sub>1</sub> antihistamines, opiate derivatives and barbiturates).
- slow the heart rate (induce bradycardia, such as anticholinesterases and beta blockers).
- are used for heart problems and complications (such as digoxin), or for thinning the blood (such as warfarin).
- are associated with neuropathy such as other treatments for cancer.
- are used for contraception.

### **Thalidomide BMS with food, drink and alcohol**

Do not drink alcohol while you are taking Thalidomide BMS. This is because alcohol can make you sleepy and Thalidomide BMS can make you even sleepier.

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

#### Pregnancy

Thalidomide causes severe birth defects or death to an unborn baby.

- As little as one capsule taken by a pregnant woman can cause a baby to have serious birth defects.
- These defects can include shortened arms or legs, malformed hands or feet, eye or ear defects, and problems with internal organs.

If you are pregnant, you must not take Thalidomide BMS. In addition, you must not become pregnant while taking Thalidomide BMS.

You must use one effective method of contraception if you are a woman who is able to become pregnant (see section 2, “What you need to know before you take Thalidomide BMS”).

#### **You must stop treatment and inform your doctor straight away if:**

- You miss or think you have missed a period, or you have unusual menstrual bleeding, or suspect you are pregnant.
- You have heterosexual intercourse without using an effective method of contraception.

If you do become pregnant during the treatment with thalidomide, you must stop the treatment and inform your doctor immediately.

For men taking Thalidomide BMS who have a female partner who is able to become pregnant, please see section 2 “What you need to know before you take Thalidomide BMS”. If your partner becomes pregnant whilst you are taking thalidomide, you should inform your doctor immediately.

#### Breast-feeding

Do not breastfeed when taking Thalidomide BMS as it is not known if thalidomide is passed into human breast milk.

### **Driving and using machines**

Do not drive or use any tools or machines if you experience side effects, such as dizziness, tiredness, sleepiness or blurred vision.

### 3. How to take Thalidomide BMS

Always take Thalidomide BMS exactly as your doctor or pharmacist has told you to. Check with your doctor or pharmacist if you are not sure.

#### How much to take


The recommended dose is 200 mg (4 x 50 mg capsules) a day for adults aged 75 years and under or 100 mg (2 x 50 mg capsules) a day for adults aged over 75 years. However your doctor will choose the dose for you, monitor your progress and may adjust your dose. Your doctor will tell you how to take Thalidomide BMS and for how long you will need to take it (see section 2, “What you need to know before you take Thalidomide BMS”).

Thalidomide BMS is taken daily in treatment cycles, each cycle lasting 6 weeks, in combination with melphalan and prednisone which are taken on days 1 to 4 of each 6 week cycle.

#### Taking this medicine

- Do not break, open or chew the capsules. If powder from a broken Thalidomide BMS capsule makes contact with the skin, wash the skin immediately and thoroughly with soap and water.
- Healthcare professionals, caregivers and family members should wear disposable gloves when handling the blister or capsule. Gloves should then be removed carefully to prevent skin exposure, placed in a sealable plastic polyethylene bag and disposed of in accordance with local requirements. Hands should then be washed thoroughly with soap and water. Women who are pregnant or suspect they may be pregnant should not handle the blister or capsule.
- Take this medicine by mouth.
- Swallow the capsules whole with a full glass of water.
- Do not crush or chew.
- Take the capsules as a single dose before going to bed. This will make you less likely to feel sleepy at other times.

To remove the capsule from the blister, press only one end of the capsule out to push it through the foil. Do not apply pressure on the centre of the capsule as this can cause it to break.



#### If you take more Thalidomide BMS than you should

If you take more Thalidomide BMS than you should, talk to a doctor or go to a hospital straightaway. If possible, take the medicine pack and this leaflet with you.

#### If you forget to take Thalidomide BMS

If you forget to take Thalidomide BMS at your regular time and

- less than 12 hours have passed: take your capsules immediately.
- more than 12 hours have passed: do not take your capsules. Take your next capsules at the usual time the next day.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following side effects may happen with this medicine:

**Stop taking Thalidomide BMS and see a doctor straight away if you notice the following serious side effects – you may need urgent medical treatment:**

- Extremely intense and serious skin reactions. The adverse reaction of the skin may appear as rashes with or without blisters. Skin irritation, sores or swelling in the mouth, throat, eyes, nose and around the genitals, oedema and fever and flulike symptoms may occur. These symptoms may be signs of the rare and serious skin reactions Stevens-Johnson syndrome, toxic epidermal necrolysis or DRESS syndrome.
- Allergic reactions such as a localised or generalised pruritic rash, angioedema and anaphylactic reaction (serious types of allergic reaction that may be manifested as hives, rashes, swelling of eyes, mouth or face, difficulty of breathing, or itching).

**Tell your doctor straight away if you notice any of the following serious side effects:**

- **Numbness, tingling, abnormal coordination or pain in your hands and feet.**  
This may be due to nerve damage (called ‘peripheral neuropathy’), which is a very common side effect. It may become very severe, painful and disabling. If you experience such symptoms, speak to your doctor straight away, who may reduce the dose or discontinue the treatment. This side effect usually happens after you have been taking this medicine for several months but can happen sooner than this. It can also happen sometime after treatment has stopped. It may not go away, or may go away slowly.
- **Sudden pain in your chest or difficulty in breathing.**  
This may be due to blood clots in the arteries leading to your lungs (called ‘pulmonary embolism’), which is a common side effect. These can happen during treatment, or after treatment has stopped.
- **Pain or swelling in your legs, especially in your lower leg or calves.**  
This may be due to blood clots in the veins of your leg (deep vein thrombosis), which is a common side effect. These can happen during treatment, or after treatment has stopped.
- **Chest pain spreading to the arms, neck, jaw, back or stomach, feeling sweaty and breathless, feeling sick or vomiting.**  
These may be symptoms of a heart attack/myocardial infarction (which may be due to blood clots in the arteries of your heart).
- **Having difficulty in seeing or speaking, which is temporary.**  
These may be symptoms of a stroke (which may be due to a clot in an artery in your brain).
- **Fever, chills, sore throat, cough, mouth ulcers or any other symptoms of infection.**
- **Bleeding or bruising in the absence of injury.**

**Other side effects include:**

It is important to note that a small number of patients with multiple myeloma may develop additional types of cancer, especially haematological malignancies, and it is possible that this risk may be increased with Thalidomide BMS treatment; therefore your doctor should carefully evaluate the benefit and risk when you are prescribed Thalidomide BMS.

**Very common** (may affect more than 1 in 10 people)

- Constipation.
- Feeling dizzy.
- Sleepiness, feeling tired.
- Shaking (tremor).
- Decreased or abnormal sensation (dysaesthesia).
- Swelling of hands and feet.

- Low blood cell counts. This may mean that you are more likely to develop infections. Your doctor may monitor your blood cell counts during treatment with Thalidomide BMS.

**Common** (may affect up to 1 in 10 people)

- Indigestion, feeling sick (nausea), being sick (vomiting), dry mouth.
- Rash, dryness of the skin.
- A fall in the number of white blood cells (neutropenia) accompanied by fever and infection.
- A fall in the number of red and white blood cells and platelets at the same time (pancytopenia).
- Feeling weak, faint or unsteady, lack of energy or strength, low blood pressure.
- Fever, feeling generally unwell.
- Convulsions.
- A spinning feeling in your head, making it difficult to stand up and move normally.
- Blurred vision.
- Chest infection (pneumonia), lung disease.
- A slow heart rate, heart failure.
- Depression, confusion, mood changes, anxiety.
- Hearing decreased or deafness.
- Kidney disease (renal failure).

**Uncommon** (may affect up to 1 in 100 people)

- Inflammation and swelling of the tubes in your lungs (bronchitis).
- Inflammation of the cells lining your stomach wall.
- A hole in part of your large bowel (colon) which can cause infection.
- Bowel obstruction.
- Fall of blood pressure on standing which may lead to fainting.
- Irregularities of the heartbeat (heart block or atrial fibrillation), feeling faint or fainting.

**Not known** (frequency cannot be estimated from the available data) :

- Underactive thyroid (hypothyroidism).
- Sexual dysfunction, for example impotence.
- Severe blood infection (sepsis) accompanied by fever, chills and severe shaking, and possibly complicated by low blood pressure and confusion (septic shock).
- Tumour Lysis Syndrome - metabolic complications that can occur during the treatment of cancer and sometimes even without treatment. These complications are caused by the break-down products of dying cancer cells and may include the following: changes to blood chemistry; high potassium, phosphorus, uric acid, and low calcium consequently leading to changes in kidney function, heart beat, seizures, and sometimes death.
- Injury to the liver (hepatic disorder) including abnormal liver test results.
- Bleeding from the stomach or bowels (gastrointestinal haemorrhage).
- Worsening of Parkinson's disease symptoms (such as tremor, depression or confusion).
- Pain in the upper abdomen and/or back, which may be severe and which remains for a few days, possibly accompanied by nausea, vomiting, fever and a rapid pulse – these symptoms may be due to the inflammation of the pancreas (pancreatitis).
- Increase in blood pressure within blood vessels that supply the lungs which can lead to shortness of breath, tiredness, dizziness, pain in the chest, a faster heartbeat, or swelling in the legs or ankles (pulmonary hypertension).
- Viral infections, including herpes zoster (also known as 'shingles', a viral disease that causes a painful skin rash with blisters) and recurrence of hepatitis B infection (which can cause yellowing of the skin and eyes, dark brown-coloured urine, right-sided stomach pain, fever and feeling nauseous or being sick).
- A brain condition with symptoms including vision changes, headache, seizures, and confusion, with or without high blood pressure (Posterior Reversible Encephalopathy Syndrome or PRES).
- A condition affecting the skin caused by inflammation of small blood vessels, along with pain in the joints and fever (leukocytoclastic vasculitis).

## Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Thalidomide BMS

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the wallet card and the blister after EXP. The expiry date refers to the last day of that month.

Do not use if you notice any damage or signs of tampering.

This medicine does not require any special storage conditions.

At the end of your treatment you should return all unused capsules to the pharmacist or doctor. These measures will prevent misuse.

## 6. Contents of the pack and other information

### What Thalidomide BMS contains

- The active substance is thalidomide. Each capsule contains 50 mg of thalidomide.
- The other excipients are:
  - The capsule content contains pregelatinised starch and magnesium stearate.
  - The capsule shell contains gelatin and titanium dioxide (E171).
  - The printing ink is composed of shellac, black iron oxide (E172) and propylene glycol.

### What Thalidomide BMS looks like and contents of the pack

Thalidomide BMS are white hard capsules marked “Thalidomide BMS 50 mg”. The capsules are supplied in a wallet card containing 28 capsules (2 blisters of 14 capsules each).

### Marketing Authorisation Holder

Bristol-Myers Squibb Pharma EEIG  
Plaza 254  
Blanchardstown Corporate Park 2  
Dublin 15, D15 T867  
Ireland

### Manufacturer

Celgene Distribution B.V.  
Orteliuslaan 1000  
3528 BD Utrecht  
Netherlands

### This leaflet was last revised in

### Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site: <http://www.ema.europa.eu>. There are also links to other websites about rare diseases and treatments.