

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

## 1. NAME OF THE MEDICINAL PRODUCT

Zydelig 100 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 100 mg of idelalisib.

### Excipient with known effect

Each tablet contains 0.1 mg sunset yellow FCF (E110) (see section 4.4).

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet.

Orange, oval-shaped, film-coated tablet of dimensions 9.7 mm by 6.0 mm, debossed on one side with “GSI” and “100” on the other side.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Zydelig is indicated in combination with rituximab for the treatment of adult patients with chronic lymphocytic leukaemia (CLL):

- who have received at least one prior therapy (see section 4.4), or
- as first line treatment in the presence of 17p deletion or *TP53* mutation in patients who are not eligible for any other therapies (see section 4.4).

Zydelig is indicated as monotherapy for the treatment of adult patients with follicular lymphoma (FL) that is refractory to two prior lines of treatment (see section 4.4).

### 4.2 Posology and method of administration

Treatment with Zydelig should be conducted by a physician experienced in the use of anti-cancer therapies.

#### Posology

The recommended dose is 150 mg idelalisib twice daily. Treatment should be continued until disease progression or unacceptable toxicity.

If the patient misses a dose of Zydelig within 6 hours of the time it is usually taken, the patient should take the missed dose as soon as possible and resume the normal dosing schedule. If a patient misses a dose by more than 6 hours, the patient should not take the missed dose and simply resume the usual dosing schedule.

#### Dose modification

##### *Elevated liver transaminases*

Treatment with Zydelig must be withheld in the event of a Grade 3 or 4 aminotransferase elevation (alanine aminotransferase [ALT]/aspartate aminotransferase [AST] > 5 x upper limit of normal [ULN]). Once values have returned to Grade 1 or below (ALT/AST ≤ 3 x ULN), treatment can be resumed at 100 mg twice daily.

If the event does not recur, the dose can be re-escalated to 150 mg twice daily at the discretion of the treating physician.

If the event recurs, treatment with Zydelig must be withheld until the values return to Grade 1 or less, after which re-initiation at 100 mg twice daily may be considered at the discretion of the physician (see sections 4.4 and 4.8).

*Diarrhoea/colitis*

Treatment with Zydelig must be withheld in the event of Grade 3 or 4 diarrhoea/colitis. Once diarrhoea/colitis has returned to Grade 1 or below, treatment can be resumed at 100 mg twice daily. If diarrhoea/colitis does not recur, the dose can be re-escalated to 150 mg twice daily at the discretion of the treating physician (see section 4.8).

*Pneumonitis*

Treatment with Zydelig must be withheld in the event of suspected pneumonitis. Once pneumonitis has resolved and if re-treatment is appropriate, resumption of treatment at 100 mg twice daily can be considered. Treatment with Zydelig must be permanently discontinued in the event of moderate or severe symptomatic pneumonitis or organising pneumonia (see sections 4.4 and 4.8).

*Rash*

Treatment with Zydelig must be withheld in the event of Grade 3 or 4 rash. Once rash has returned to Grade 1 or below, treatment can be resumed at 100 mg twice daily. If rash does not recur, the dose can be re-escalated to 150 mg twice daily at the discretion of the treating physician (see section 4.8).

*Neutropenia*

Treatment with Zydelig should be withheld in patients while absolute neutrophil count (ANC) is below 500 per mm<sup>3</sup>. ANC should be monitored at least weekly until ANC is ≥ 500 per mm<sup>3</sup> when treatment can be resumed at 100 mg twice daily (see section 4.4).

| ANC 1,000 to < 1,500/mm <sup>3</sup> | ANC 500 to < 1,000/mm <sup>3</sup>                           | ANC < 500/mm <sup>3</sup>  |
|--------------------------------------|--|--|
| Maintain Zydelig dosing.             | Maintain Zydelig dosing.<br><br>Monitor ANC at least weekly. | Interrupt Zydelig dosing.<br><br>Monitor ANC at least weekly until ANC ≥ 500/mm <sup>3</sup> , then may resume Zydelig dosing at 100 mg twice daily. |

*Special populations*

*Elderly*

No specific dose adjustment is required for elderly patients (aged ≥ 65 years) (see section 5.2).

*Renal impairment*

No dose adjustment is required for patients with mild, moderate, or severe renal impairment (see section 5.2).

*Hepatic impairment*

No dose adjustment is required when initiating treatment with Zydelig in patients with mild or moderate hepatic impairment, but an intensified monitoring of adverse reactions is recommended (see sections 4.4 and 5.2).

There is insufficient data to make dose recommendations for patients with severe hepatic impairment. Therefore, caution is recommended when administering Zydelig in this population and an intensified monitoring of adverse reactions is recommended (see sections 4.4 and 5.2).

### Paediatric population

The safety and efficacy of Zydelig in children under the age of 18 years have not been established. No data are available.

### Method of administration

Zydelig is for oral use. Patients should be instructed to swallow the tablet whole. The film-coated tablet should not be chewed or crushed. The film-coated tablet can be taken with or without food (see section 5.2).

## **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

## **4.4 Special warnings and precautions for use**

### Serious infections

Treatment with Zydelig should not be initiated in patients with any evidence of ongoing systemic bacterial, fungal, or viral infection.

Serious and fatal infections have occurred with idelalisib, including opportunistic infections such as *Pneumocystis jirovecii* pneumonia (PJP) and cytomegalovirus (CMV). Prophylaxis for PJP should therefore be administered to all patients throughout idelalisib treatment, and for a period of 2 to 6 months after discontinuation. The duration of post-treatment prophylaxis should be based on clinical judgment and may take into account a patient's risk factors such as concomitant corticosteroid treatment and prolonged neutropenia (see section 4.8).

Patients should be monitored for respiratory signs and symptoms throughout treatment. Patients should be advised to report new respiratory symptoms promptly.

Regular clinical and laboratory monitoring for CMV infection is recommended in patients with positive CMV serology at the start of treatment with idelalisib or with other evidence of a history of CMV infection. Patients with CMV viraemia without associated clinical signs of CMV infection should be carefully monitored. For patients with evidence of CMV viraemia and clinical signs of CMV infection, consideration should be given to interrupting idelalisib until the infection has resolved. If the benefits of resuming idelalisib are judged to outweigh the risks, consideration should be given to administering pre-emptive CMV therapy.

Cases of progressive multifocal leukoencephalopathy (PML) have been reported following the use of idelalisib within the context of prior or concomitant immunosuppressive therapies that have been associated with PML. Physicians should consider PML in the differential diagnosis in patients with new or worsening neurological, cognitive or behavioural signs or symptoms. If PML is suspected then appropriate diagnostic evaluations should be undertaken and treatment suspended until PML is excluded. If any doubt exists, referral to a neurologist and appropriate diagnostic measures for PML including MRI scan preferably with contrast, cerebrospinal fluid (CSF) testing for JC viral DNA and repeat neurological assessments should be considered.

### Neutropenia

Treatment-emergent Grade 3 or 4 neutropenia, including febrile neutropenia, have occurred in patients treated with idelalisib. Blood counts should be monitored in all patients at least every 2 weeks for the first 6 months of treatment with idelalisib, and at least weekly in patients while ANC is less than 1,000 per mm<sup>3</sup> (see section 4.2).

### Hepatotoxicity

Elevations in ALT and AST of Grade 3 and 4 (> 5 x ULN) have been observed in clinical studies of idelalisib. There have also been reports of hepatocellular injury including hepatic failure. Increases in liver transaminases were generally observed within the first 12 weeks of treatment, and were reversible with dose interruption (see section 4.2). In patients who resumed idelalisib at a lower dose,

26% had recurrence of ALT/AST elevation. Treatment with Zydelig must be withheld in the event of Grade 3 or 4 ALT/AST elevation and liver function monitored. Treatment may be resumed at a lower dose once values have returned to Grade 1 or below (ALT/AST  $\leq 3 \times$  ULN).

ALT, AST, and total bilirubin must be monitored in all patients every 2 weeks for the first 3 months of treatment, then as clinically indicated. If Grade 2 or higher elevations in ALT and/or AST are observed, patients' ALT, AST, and total bilirubin must be monitored weekly until the values return to Grade 1 or below.

#### Diarrhoea/colitis

Cases of severe drug-related colitis occurred relatively late (months) after the start of therapy, sometimes with rapid aggravation, but resolved within a few weeks with dose interruption and additional symptomatic treatment (e.g., anti-inflammatory agents such as enteric budesonide).

There is very limited experience from the treatment of patients with a history of inflammatory bowel disease.

#### Pneumonitis and organising pneumonia

Cases of pneumonitis and organising pneumonia (some with fatal outcome) have been reported with idelalisib. In patients presenting with serious lung events, idelalisib should be interrupted and the patient assessed for an explanatory aetiology. If either moderate or severe symptomatic pneumonitis or organising pneumonia is diagnosed, appropriate treatment should be initiated and idelalisib must be permanently discontinued.

#### Severe Cutaneous Reactions

Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS) have occurred with idelalisib. Cases of SJS and TEN with fatal outcomes have been reported when idelalisib was administered concomitantly with other medicinal products associated with these syndromes. If SJS, TEN or DRESS is suspected, idelalisib should be interrupted and the patient assessed and treated accordingly. If a diagnosis of SJS, TEN, or DRESS is confirmed, idelalisib should be permanently discontinued.

#### CYP3A inducers

Idelalisib exposure may be reduced when co-administered with CYP3A inducers such as rifampicin, phenytoin, St. John's wort (*Hypericum perforatum*), or carbamazepine. Since a reduction in idelalisib plasma concentrations may result in decreased efficacy, co-administration of Zydelig with moderate or strong CYP3A inducers should be avoided (see section 4.5).

#### CYP3A substrates

The primary metabolite of idelalisib, GS-563117, is a strong CYP3A4 inhibitor. Thus, idelalisib has the potential to interact with medicinal products that are metabolised by CYP3A, which may lead to increased serum concentrations of the other product (see section 4.5). When idelalisib is co-administered with other medicinal products, the Summary of Product Characteristics (SmPC) for the other product must be consulted for the recommendations regarding co-administration with CYP3A4 inhibitors. Concomitant treatment of idelalisib with CYP3A substrates with serious and/or life-threatening adverse reactions (e.g., alfuzosin, amiodarone, cisapride, pimozone, quinidine, ergotamine, dihydroergotamine, quetiapine, lovastatin, simvastatin, sildenafil, midazolam, triazolam) should be avoided and alternative medicinal products that are less sensitive to CYP3A4 inhibition should be used if possible.

#### Hepatic impairment

Intensified monitoring of adverse reactions is recommended in patients with impaired hepatic function as exposure is expected to be increased in this population, in particular in patients with severe hepatic impairment. No patients with severe hepatic impairment were included in clinical studies of idelalisib. Caution is recommended when administering Zydelig in this population.

### Chronic hepatitis

Idelalisib has not been studied in patients with chronic active hepatitis including viral hepatitis. Caution should be exercised when administering Zydelig in patients with active hepatitis.

### Women of childbearing potential

Women of childbearing potential must use highly effective contraception while taking idelalisib and for 1 month after stopping treatment (see section 4.6). Women using hormonal contraceptives should add a barrier method as a second form of contraception since it is currently unknown whether idelalisib may reduce the effectiveness of hormonal contraceptives.

### Excipients

Zydelig contains the azo colouring agent sunset yellow FCF (E110), which may cause allergic reactions.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Idelalisib is metabolised primarily via aldehyde oxidase, and to a lesser extent via CYP3A and glucuronidation (UGT1A4). Its primary metabolite is GS-563117, which is not pharmacologically active. Idelalisib and GS-563117 are substrates of P-gp and BCRP.

### Effect of other medicinal products on idelalisib pharmacokinetics

#### *CYP3A inducers*

A clinical drug interaction study found that co-administration of a single dose of 150 mg idelalisib with rifampicin (a strong CYP3A inducer) resulted in a ~75% reduction in idelalisib AUC<sub>inf</sub>. Co-administration of Zydelig with moderate or strong CYP3A inducers such as rifampicin, phenytoin, St. John's wort, or carbamazepine should be avoided as this may result in decreased efficacy (see section 4.4).

#### *CYP3A/P-gp inhibitors*

A clinical drug interaction study found that co-administration of a single dose of 400 mg idelalisib with 400 mg once daily ketoconazole (a strong CYP3A, P-gp and BCRP inhibitor) resulted in a 26% increase in C<sub>max</sub> and a 79% increase in AUC<sub>inf</sub> of idelalisib. No initial dose adjustment of idelalisib is considered necessary when administered with CYP3A/P-gp inhibitors, but an intensified monitoring of adverse reactions is recommended.

### Effect of idelalisib on the pharmacokinetics of other medicinal products

#### *CYP3A substrates*

The primary metabolite of idelalisib, GS-563117, is a strong CYP3A inhibitor. A clinical drug interaction study found that co-administration of idelalisib with midazolam (a sensitive CYP3A substrate) resulted in a ~140% increase in C<sub>max</sub> and a ~440% increase in AUC<sub>inf</sub> of midazolam due to the CYP3A inhibition by GS-563117. Co-administration of idelalisib with CYP3A substrates may increase their systemic exposures and increase or prolong their therapeutic activity and adverse reactions. *In vitro*, the CYP3A4 inhibition was irreversible, and return to normal enzyme activity is therefore expected to take several days after stopping idelalisib administration.

Potential interactions between idelalisib and co-administered medicinal products that are CYP3A substrates are listed in Table 1 (increase is indicated as “↑”). This list is not exhaustive and is intended to serve as guidance only. In general, the SmPC for the other product must be consulted for the recommendations regarding co-administration with CYP3A4 inhibitors (see section 4.4).

**Table 1: Interactions between idelalisib and other medicinal products that are CYP3A substrates**

| Medicinal product  | Expected effect of idelalisib on drug levels | Clinical recommendation upon co-administration with idelalisib  |
|--|--|---|
| <b>ALPHA-1 ADRENORECEPTOR ANTAGONISTS</b>  |  |   |
| Alfuzosin  | ↑ serum concentrations                       | Idelalisib should not be co-administered with alfuzosin.  |
| <b>ANALGESICS</b>  |  |   |
| Fentanyl, alfentanil, methadone, buprenorphine/naloxone                                      | ↑ serum concentrations                       | Careful monitoring of adverse reactions (e.g., respiratory depression, sedation) is recommended.  |
| <b>ANTIARRHYTHMICS</b>   |  |   |
| Amiodarone, quinidine  | ↑ serum concentrations                       | Idelalisib should not be co-administered with amiodarone or quinidine.  |
| Bepidil, disopyramide, lidocaine   | ↑ serum concentrations                       | Clinical monitoring is recommended.   |
| <b>ANTI-CANCER AGENTS</b>  |  |   |
| Tyrosine kinase inhibitors such as dasatinib and nilotinib, also vincristine and vinblastine | ↑ serum concentrations                       | Careful monitoring of the tolerance to these anti-cancer agents is recommended.   |
| <b>ANTICOAGULANTS</b>  |  |   |
| Warfarin   | ↑ serum concentrations                       | It is recommended that the international normalised ratio (INR) be monitored upon co-administration and following ceasing treatment with idelalisib.  |
| <b>ANTICONVULSANTS</b>   |  |   |
| Carbamazepine  | ↑ serum concentrations                       | Anticonvulsant drug levels should be monitored.   |
| <b>ANTIDEPRESSANTS</b>   |  |   |
| Trazodone  | ↑ serum concentrations                       | Careful dose titration of the antidepressant and monitoring for antidepressant response is recommended.   |
| <b>ANTI-GOUT</b>   |  |   |
| Colchicine   | ↑ serum concentrations                       | Dose reductions of colchicine may be required. Idelalisib should not be co-administered with colchicine to patients with renal or hepatic impairment. |
| <b>ANTI-HYPERTENSIVES</b>  |  |   |
| Amlodipine, diltiazem, felodipine, nifedipine, nicardipine                                   | ↑ serum concentrations                       | Clinical monitoring of therapeutic effect and adverse reactions is recommended.   |
| <b>ANTI-INFECTIVES</b>   |  |   |
| <b>Antifungals</b>   |  |   |
| Ketoconazole, itraconazole, posaconazole, voriconazole                                       | ↑ serum concentrations                       | Clinical monitoring is recommended.   |
| <b>Antimycobacterials</b>  |  |   |
| Rifabutin  | ↑ serum concentrations                       | Increased monitoring for rifabutin-associated adverse reactions including neutropenia and uveitis is recommended.                                     |

| <b>Medicinal product</b>                                  | <b>Expected effect of idelalisib on drug levels</b> | <b>Clinical recommendation upon co-administration with idelalisib</b>  |
|---|---|--|
| <b>HCV protease inhibitors</b>                            |   |  |
| Boceprevir, telaprevir                                    | ↑ serum concentrations                              | Clinical monitoring is recommended.  |
| <b>Macrolide antibiotics</b>                              |   |  |
| Clarithromycin, telithromycin                             | ↑ serum concentrations                              | No dose adjustment of clarithromycin is required for patients with normal renal function or mild renal impairment (creatinine clearance [CrCl] 60-90 mL/min). Clinical monitoring is recommended for patients with CrCl < 90 mL/min. For patients with CrCl < 60 mL/min, alternative antibacterials should be considered.<br><br>Clinical monitoring is recommended for telithromycin. |
| <b>ANTI-PSYCHOTICS/NEUROLEPTICS</b>                       |   |  |
| Quetiapine, pimozone                                      | ↑ serum concentrations                              | Idelalisib should not be co-administered with quetiapine or pimozone.<br><br>Alternative medicinal products, such as olanzapine, may be considered.  |
| <b>ENDOTHELIN RECEPTOR ANTAGONISTS</b>                    |   |  |
| Bosentan  | ↑ serum concentrations                              | Caution should be exercised and patients closely observed for bosentan-related toxicity.   |
| <b>ERGOT ALKALOIDS</b>                                    |   |  |
| Ergotamine, dihydroergotamine                             | ↑ serum concentrations                              | Idelalisib should not be co-administered with ergotamine or dihydroergotamine.   |
| <b>GASTROINTESTINAL MOTILITY AGENTS</b>                   |   |  |
| Cisapride   | ↑ serum concentrations                              | Idelalisib should not be co-administered with cisapride.   |
| <b>GLUCOCORTICOIDS</b>                                    |   |  |
| Inhaled/nasal corticosteroids:<br>Budesonide, fluticasone | ↑ serum concentrations                              | Clinical monitoring is recommended.  |
| Oral budesonide   | ↑ serum concentrations                              | Clinical monitoring is recommended for increased signs/symptoms of corticosteroid effects.   |
| <b>HMG CO-A REDUCTASE INHIBITORS</b>                      |   |  |
| Lovastatin, simvastatin                                   | ↑ serum concentrations                              | Idelalisib should not be co-administered with lovastatin or simvastatin.   |
| Atorvastatin  | ↑ serum concentrations                              | Clinical monitoring is recommended and a lower starting dose of atorvastatin may be considered. Alternatively, switching to pravastatin, rosuvastatin, or pitavastatin may be considered.  |



| Medicinal product   | Expected effect of idelalisib on drug levels | Clinical recommendation upon co-administration with idelalisib   |
|---|--|--|
| <b>IMMUNOSUPPRESSANTS</b>   |  |  |
| Ciclosporin, sirolimus, tacrolimus                                | ↑ serum concentrations                       | Therapeutic monitoring is recommended.   |
| <b>INHALED BETA AGONIST</b>                                       |  |  |
| Salmeterol  | ↑ serum concentrations                       | Concurrent administration of salmeterol and idelalisib is not recommended. The combination may result in increased risk of cardiovascular adverse events associated with salmeterol, including QT prolongation, palpitations, and sinus tachycardia. |
| <b>PHOSPHODIESTERASE INHIBITORS</b>                               |  |  |
| Sildenafil  | ↑ serum concentrations                       | For pulmonary arterial hypertension:<br>Idelalisib should not be co-administered with sildenafil.  |
| Tadalafil   | ↑ serum concentrations                       | Caution should be exercised, including consideration of dose reduction, when co-administering tadalafil with idelalisib.   |
| Sildenafil, tadalafil   | ↑ serum concentrations                       | For erectile dysfunction:<br>Particular caution must be used and dose reduction may be considered when prescribing sildenafil or tadalafil with idelalisib with increased monitoring for adverse events.   |
| <b>SEDATIVES/HYPNOTICS</b>  |  |  |
| Midazolam (oral), triazolam                                       | ↑ serum concentrations                       | Idelalisib should not be co-administered with midazolam (oral) or triazolam.   |
| Buspirone, clorazepate, diazepam, estazolam, flurazepam, zolpidem | ↑ serum concentrations                       | Concentration monitoring of sedatives/hypnotics is recommended and dose reduction may be considered.   |

#### *CYP2C8 substrates*

*In vitro*, idelalisib both inhibited and induced CYP2C8, but it is not known whether this translates to an *in vivo* effect on CYP2C8 substrates. Caution is advised if Zydelig is used together with narrow therapeutic index drugs that are substrates of CYP2C8 (paclitaxel).

#### *Substrates of inducible enzymes (e.g., CYP2C9, CYP2C19, CYP2B6 and UGT)*

*In vitro*, idelalisib was an inducer of several enzymes, and a risk for decreased exposure and thereby decreased efficacy of substrates of inducible enzymes such as CYP2C9, CYP2C19, CYP2B6 and UGT cannot be excluded. Caution is advised if Zydelig is used together with narrow therapeutic index drugs that are substrates of these enzymes (warfarin, phenytoin, S-mephenytoin).

#### *BCRP, OATP1B1, OATP1B3 and P-gp substrates*

Co-administration of multiple doses of idelalisib 150 mg twice daily to healthy subjects resulted in comparable exposures for rosuvastatin (AUC 90% CI: 87, 121) and digoxin (AUC 90% CI: 98, 111), suggesting no clinically relevant inhibition of BCRP, OATP1B1/1B3 or systemic P-gp by idelalisib. A

risk for P-gp inhibition in the gastrointestinal tract, that could result in increased exposure of sensitive substrates for intestinal P-gp such as dabigatran etexilate, cannot be excluded.

#### Paediatric population

Interaction studies have only been performed in adults.

### **4.6 Fertility, pregnancy and lactation**

#### Women of childbearing potential / contraception

Based on findings in animals, idelalisib may cause foetal harm. Women should avoid becoming pregnant while taking Zydelig, and for up to 1 month after ending treatment. Therefore, women of childbearing potential must use highly effective contraception while taking Zydelig and for 1 month after stopping treatment. It is currently unknown whether idelalisib may reduce the effectiveness of hormonal contraceptives, and therefore women using hormonal contraceptives should add a barrier method as a second form of contraception.

#### Pregnancy

There are no or limited amount of data from the use of idelalisib in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3).

Zydelig is not recommended during pregnancy and in women of childbearing potential not using contraception.

#### Breast-feeding

It is not known whether idelalisib and its metabolites are excreted in human milk.

A risk to the newborns/infants cannot be excluded.

Breast-feeding should be discontinued during treatment with Zydelig.

#### Fertility

No human data on the effect of idelalisib on fertility are available. Animal studies indicate the potential for harmful effects of idelalisib on fertility and foetal development (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Zydelig has no or negligible influence on the ability to drive and use machines.

### **4.8 Undesirable effects**

#### Summary of the safety profile

Assessment of adverse reactions is based on two Phase 3 studies (study 312-0116 and study 312-0119) and six Phase 1 and 2 studies. Study 312-0116 was a randomised, double-blind, placebo-controlled study in which 110 subjects with previously treated CLL received idelalisib + rituximab. In addition, 86 subjects from this study who were randomised to receive placebo + rituximab went on to receive idelalisib as a single agent in an extension study (study 312-0117). Study 312-0119 was a randomised, controlled, open-label study in which 173 subjects with previously treated CLL received idelalisib + ofatumumab. The Phase 1 and 2 studies assessed the safety of idelalisib in a total of 536 subjects with haematologic malignancies, including 400 subjects who received idelalisib (any dose) as a single agent and 136 subjects who received idelalisib in combination with an anti-CD20 monoclonal antibody (rituximab or ofatumumab).

#### Tabulated list of adverse reactions

The adverse drug reactions reported with idelalisib alone or in combination with anti-CD20 monoclonal antibodies (rituximab or ofatumumab) are provided in Table 2. Adverse reactions are listed by system organ class and frequency. Frequencies are defined as follows: very common

(≥ 1/10), common (≥ 1/100 to < 1/10), uncommon (≥ 1/1,000 to < 1/100), rare (≥ 1/10,000 to < 1/1,000), very rare (< 1/10,000), and not known (cannot be estimated from available data).

**Table 2: Adverse drug reactions reported in clinical studies in subjects with haematologic malignancies receiving idelalisib and post-marketing.**

| Reaction  | Any grade   | Grade ≥ 3   |
|---|-------------|-------------|
| <i>Infections and infestations</i>                                      |             |             |
| Infections (including <i>Pneumocystis jirovecii</i> pneumonia and CMV)* | Very common | Very common |
| <i>Blood and lymphatic system disorders</i>                             |             |             |
| Neutropenia   | Very common | Very common |
| Lymphocytosis**   | Very common | Very common |
| <i>Respiratory, thoracic and mediastinal disorders</i>                  |             |             |
| Pneumonitis   | Common      | Common      |
| Organising pneumonia****  | Uncommon    | Uncommon    |
| <i>Gastrointestinal disorders</i>                                       |             |             |
| Diarrhoea/colitis   | Very common | Very common |
| <i>Hepatobiliary disorders</i>  |             |             |
| Transaminase increased  | Very common | Very common |
| Hepatocellular injury   | Common      | Common      |
| <i>Skin and subcutaneous tissue disorders</i>                           |             |             |
| Rash***   | Very common | Common      |
| Stevens-Johnson syndrome/<br>toxic epidermal necrolysis****             | Rare        | Rare        |
| Drug reaction with<br>eosinophilia and systemic<br>symptoms (DRESS)**** | Not known   | Not known   |
| <i>General disorders and administration site conditions</i>             |             |             |
| Pyrexia   | Very common | Common      |
| <i>Investigations</i>   |             |             |
| Increased triglycerides   | Very common | Common      |

\* Comprised of opportunistic infections as well as bacterial and viral infections such as pneumonia, bronchitis, and sepsis.

\*\* Idelalisib-induced lymphocytosis should not be considered progressive disease in the absence of other clinical findings (see section 5.1).

\*\*\* Includes the preferred terms dermatitis exfoliative generalised, drug eruption, rash, rash erythematous, rash generalised, rash macular, rash maculo-papular, rash papular, rash pruritic, rash pustular, rash vesicular, papule, skin plaque, and exfoliative rash.

\*\*\*\* Observed in post-marketing data

### Description of selected adverse reactions

#### *Infections (see section 4.4)*

Higher frequencies of infections overall, including Grade 3 and 4 infections, were observed in the idelalisib arms compared to the control arms of idelalisib clinical studies. Most frequently observed were infections in the respiratory system and septic events. In many instances the pathogen was not identified; however, both conventional and opportunistic pathogens, including PJP and CMV, were among those identified. Nearly all PJP infections, including fatal cases, occurred in the absence of PJP prophylaxis. There have been cases of PJP after stopping idelalisib treatment.

#### *Rash*

Rash was generally mild to moderate and resulted in discontinuation of treatment in 2.1% of subjects. In studies 312-0116/0117 and 312-0119, rash (reported as dermatitis exfoliative generalised, drug eruption, rash, rash erythematous, rash generalised, rash macular, rash maculo-papular, rash papular, rash pruritic, rash pustular, rash vesicular, papule and skin plaque) occurred in 31.1% of subjects who

received idelalisib + an anti-CD20 monoclonal antibody (rituximab or ofatumumab) and 8.2% who received an anti-CD20 monoclonal antibody only (rituximab or ofatumumab). Of these, 5.7% who received idelalisib + an anti-CD20 monoclonal antibody (rituximab or ofatumumab) and 1.5% who received an anti-CD20 monoclonal antibody only (rituximab or ofatumumab) had rash of Grade 3, and no subjects had an adverse reaction of Grade 4. Rash typically resolved with treatment (e.g., topical and/or oral steroids, diphenhydramine) and dose interruption for severe cases (see section 5.3, phototoxicity).

#### *Severe Cutaneous Reactions (see section 4.4)*

Cases of SJS, TEN and DRESS have occurred when idelalisib was administered concomitantly with other medicinal products associated with these syndromes (bendamustine, rituximab, allopurinol, amoxicillin, and sulfamethoxazole / trimethoprim). SJS or TEN occurred within one month of the medicinal combination and fatal outcomes have resulted.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

### **4.9 Overdose**

If overdose occurs the patient must be monitored for evidence of toxicity (see section 4.8). Treatment of overdose with Zydelig consists of general supportive measures including monitoring of vital signs as well as observation of the clinical status of the patient.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: antineoplastic agents, other antineoplastic agents, ATC code: L01EM01

#### Mechanism of action

Idelalisib inhibits phosphatidylinositol 3-kinase p110 $\delta$  (PI3K $\delta$ ), which is hyperactive in B-cell malignancies and is central to multiple signalling pathways that drive proliferation, survival, homing, and retention of malignant cells in lymphoid tissues and bone marrow. Idelalisib is a selective inhibitor of adenosine-5'-triphosphate (ATP) binding to the catalytic domain of PI3K $\delta$ , resulting in inhibition of the phosphorylation of the key lipid second messenger phosphatidylinositol and prevention of Akt (protein kinase B) phosphorylation.

Idelalisib induces apoptosis and inhibits proliferation in cell lines derived from malignant B-cells and in primary tumour cells. Through inhibition of chemokine receptors CXCR4 and CXCR5 signalling induced by the chemokines CXCL12 and CXCL13, respectively, idelalisib inhibits homing and retention of malignant B-cells in the tumour microenvironment including lymphoid tissues and the bone marrow.

No mechanistic explanations for the development of resistance to treatment with idelalisib have been identified from clinical studies. Further investigation of this topic in current B-cell malignancy studies is not planned.

## Pharmacodynamic effects

### *Electrocardiographic*

The effect of idelalisib (150 mg and 400 mg) on the QT/QTc interval was evaluated in a placebo- and positive-controlled (moxifloxacin 400 mg) crossover study in 40 healthy subjects. At a dose 2.7 times the maximum recommended dose, idelalisib did not prolong the QT/QTc interval (i.e., < 10 ms).

### *Lymphocytosis*

Upon initiation of idelalisib, a temporary increase in lymphocyte counts (i.e.,  $\geq 50\%$  increase from baseline and above absolute lymphocyte count of 5,000/mcL) has been observed. This occurs in approximately two-thirds of patients with CLL treated with idelalisib monotherapy and one-fourth of patients with CLL treated with idelalisib combination therapy. The onset of isolated lymphocytosis typically occurs during the first 2 weeks of idelalisib therapy and is often associated with reduction of lymphadenopathy. This observed lymphocytosis is a pharmacodynamic effect and should not be considered progressive disease in the absence of other clinical findings.

## Clinical efficacy in chronic lymphocytic leukaemia

### *Idelalisib in combination with rituximab*

Study 312-0116 was a Phase 3, randomised, double-blind, placebo-controlled study in 220 subjects with previously treated CLL who required treatment but were not considered suitable for cytotoxic chemotherapy. Subjects were randomised 1:1 to receive 8 cycles of rituximab (first cycle at 375 mg/m<sup>2</sup> body surface area [BSA], subsequent cycles at 500 mg/m<sup>2</sup> BSA) in combination with either an oral placebo twice daily or with idelalisib 150 mg taken twice daily until disease progression or unacceptable toxicity.

The median age was 71 years (range: 47 to 92) with 78.2% of subjects over 65 years; 65.5% were male, and 90.0% were white; 64.1% had a Rai stage of III or IV, and 55.9% had Binet Stage C. Most subjects had adverse cytogenetic prognostic factors: 43.2% had a 17p chromosomal deletion and/or tumour protein 53 (*TP53*) mutation, and 83.6% had unmutated genes for the immunoglobulin heavy chain variable region (*IGHV*). The median time from diagnosis of CLL to randomisation was 8.5 years. Subjects had a median Cumulative Illness Rating Scale (CIRS) score of 8. The median number of prior therapies was 3.0. Nearly all (95.9%) subjects had received prior anti-CD20 monoclonal antibodies. The primary endpoint was progression free survival (PFS). Efficacy results are summarised in Tables 3 and 4. The Kaplan-Meier curve for PFS is provided in Figure 1.

Compared with rituximab + placebo, treatment with idelalisib + rituximab resulted in statistically significant and clinically meaningful improvements in physical well-being, social well-being, functional well-being, as well as in the leukaemia-specific subscales of the Functional Assessment of Cancer Therapy: Leukaemia (FACT-LEU) instruments, and in statistically significant and clinically meaningful improvements in anxiety, depression and usual activities as measured by the EuroQoL Five-Dimensions (EQ-5D) instrument.

**Table 3: Efficacy results from study 312-0116**

|  | <b>Idelalisib + R</b><br><b>N = 110</b> | <b>Placebo + R</b><br><b>N = 110</b> |
|--|---|--------------------------------------|
| <b>PFS</b> Median (months) (95% CI)            | 19.4 (12.3, NR)                         | 6.5 (4.0, 7.3)                       |
| Hazard ratio (95% CI)                          | 0.15 (0.09, 0.24)                       |                                      |
| P-value  | < 0.0001                                |                                      |
| <b>ORR*</b> n (%) (95% CI)                     | 92 (83.6%) (75.4, 90.0)                 | 17 (15.5%) (9.3, 23.6)               |
| Odds ratio (95% CI)                            | 27.76 (13.40, 57.49)                    |                                      |
| P-value  | < 0.0001                                |                                      |
| <b>LNR**</b> n/N (%) (95% CI)                  | 102/106 (96.2%) (90.6, 99.0)            | 7/104 (6.7%) (2.7, 13.4)             |
| Odds ratio (95% CI)                            | 225.83 (65.56, 777.94)                  |                                      |
| P-value  | < 0.0001                                |                                      |
| <b>OS<sup>^</sup></b> Median (months) (95% CI) | NR (NR, NR)                             | 20.8 (14.8, NR)                      |
| Hazard ratio (95% CI)                          | 0.34 (0.19, 0.60)                       |                                      |
| P-value  | 0.0001                                  |                                      |

CI: confidence interval; R: rituximab; n: number of responding subjects; N: number of subjects per group; NR: not reached. The analyses of PFS, overall response rate (ORR) and lymph node response rate (LNR) were based on evaluation by an independent review committee (IRC).

\* ORR defined as the proportion of subjects who achieved a complete response (CR) or partial response (PR) based on the 2013 National Comprehensive Cancer Network (NCCN) response criteria and Cheson (2012).

\*\* LNR defined as the proportion of subjects who achieved a  $\geq 50\%$  decrease in the sum of products of the greatest perpendicular diameters of index lesions. Only subjects that had both baseline and  $\geq 1$  evaluable post-baseline assessments were included in this analysis.


<sup>^</sup> Overall survival (OS) analysis includes data from subjects who received placebo + R on study 312-0116 and subsequently received idelalisib in an extension study, based on intent-to-treat analysis.

**Table 4: Summary of PFS and response rates in pre-specified subgroups from study 312-0116**

|                                       | <b>Idelalisib + R</b><br><b>N = 46</b> | <b>Placebo + R</b><br><b>N = 49</b> |
|---------------------------------------|--|-------------------------------------|
| <b>17p deletion/TP53 mutation</b>     |  |                                     |
| PFS median (months) (95% CI)          | NR (12.3, NR)                          | 4.0 (3.7, 5.7)                      |
| Hazard ratio (95% CI)                 | 0.13 (0.07, 0.27)                      |                                     |
| ORR (95% CI)                          | 84.8% (71.1, 93.7)                     | 12.2% (4.6, 24.8)                   |
| <b>Unmutated IGHV</b>                 | <b>N = 91</b>                          | <b>N = 93</b>                       |
| PFS median (months) (95% CI)          | 19.4 (13.9, NR)                        | 5.6 (4.0, 7.2)                      |
| Hazard ratio (95% CI)                 | 0.14 (0.08, 0.23)                      |                                     |
| ORR (95% CI)                          | 82.4% (73.0, 89.6)                     | 15.1% (8.5, 24.0)                   |
| <b>Age <math>\geq 65</math> years</b> | <b>N = 89</b>                          | <b>N = 83</b>                       |
| PFS median (months) (95% CI)          | 19.4 (12.3, NR)                        | 5.7 (4.0, 7.3)                      |
| Hazard ratio (95% CI)                 | 0.14 (0.08, 0.25)                      |                                     |
| ORR (95% CI)                          | 84.3% (75.0, 91.1)                     | 16.9% (9.5, 26.7)                   |

CI: confidence interval; R: rituximab; N: number of subjects per group; NR: not reached

**Figure 1: Kaplan-Meier curve of PFS from study 312-0116 (intent-to-treat population)**



N at risk (Events)

|                |         |         |         |         |         |         |         |        |        |        |        |        |
|----------------|---------|---------|---------|---------|---------|---------|---------|--------|--------|--------|--------|--------|
| Idelalisib + R | 110 (0) | 101 (3) | 93 (7)  | 73 (9)  | 59 (14) | 31 (19) | 20 (21) | 9 (24) | 7 (24) | 4 (24) | 1 (25) | 0 (25) |
| Placebo + R    | 110 (0) | 84 (21) | 48 (38) | 29 (46) | 20 (53) | 9 (63)  | 4 (67)  | 1 (69) | 0 (70) | 0 (70) | 0 (70) | 0 (70) |

Solid line: idelalisib + R (N = 110), dashed line: placebo + R (N = 110)

R: rituximab; N: number of subjects per group

The analysis of PFS was based on evaluation by an IRC. For subjects in the placebo + R group, the summary includes data up to the first dosing of idelalisib in an extension study.

Study 101-08/99 enrolled 64 subjects with previously untreated CLL, including 5 subjects with small lymphocytic lymphoma (SLL). Subjects received idelalisib 150 mg twice daily and rituximab 375 mg/m<sup>2</sup> BSA weekly for 8 doses. The ORR was 96.9%, with 12 CRs (18.8%) and 50 PRs (78.1%), including 3 CRs and 6 PRs in subjects with a 17p deletion and/or *TP53* mutation and 2 CRs and 34 PRs in subjects with unmutated *IGHV*. The median duration of response (DOR) has not been reached.

#### *Idelalisib in combination with ofatumumab*

Study 312-0119 was a Phase 3, randomised, open-label, multicentre, parallel-group study in 261 subjects with previously treated CLL who had measurable lymphadenopathy, required treatment, and experienced CLL progression < 24 months since the completion of the last prior therapy. Subjects were randomised 2:1 to receive idelalisib 150 mg twice daily and 12 infusions of ofatumumab over 24 weeks, or 12 infusions of ofatumumab only over 24 weeks. The first infusion of ofatumumab was administered at a dose of 300 mg and was continued at a dose of either 1,000 mg in the idelalisib + ofatumumab group or a dose of 2,000 mg in the ofatumumab only group, weekly for 7 doses, and then every 4 weeks for 4 doses. Idelalisib was taken until disease progression or unacceptable toxicity.

The median age was 68 years (range: 61 to 74) with 64.0% of subjects over 65 years; 71.3% were male, and 84.3% were white; 63.6% had a Rai stage of III or IV, and 58.2% had Binet Stage C. Most subjects had adverse cytogenetic prognostic factors: 39.5% had a 17p chromosomal deletion and/or *TP53* mutation, and 78.5% had unmutated genes for *IGHV*. The median time since diagnosis was 7.7 years. Subjects had a median CIRS score of 4. The median number of prior therapies was 3.0. The

primary endpoint was PFS. Efficacy results are summarised in Tables 5 and 6. The Kaplan-Meier curve for PFS is provided in Figure 2.

**Table 5: Efficacy results from study 312-0119**

|                                     | <b>Idelalisib + O</b><br><b>N = 174</b> | <b>Ofatumumab</b><br><b>N = 87</b> |
|-------------------------------------|---|------------------------------------|
| <b>PFS</b> Median (months) (95% CI) | 16.3 (13.6, 17.8)                       | 8.0 (5.7, 8.2)                     |
| Hazard ratio (95% CI)               | 0.27 (0.19, 0.39)                       |                                    |
| P-value                             | < 0.0001                                |                                    |
| <b>ORR*</b> n (%) (95% CI)          | 131 (75.3%) (68.2, 81.5)                | 16 (18.4%) (10.9, 28.1)            |
| Odds ratio (95% CI)                 | 15.94 (7.8, 32.58)                      |                                    |
| P-value                             | < 0.0001                                |                                    |
| <b>LNR**</b> n/N (%) (95% CI)       | 153/164 (93.3%) (88.3, 96.6)            | 4/81 (4.9%) (1.4, 12.2)            |
| Odds ratio (95% CI)                 | 486.96 (97.91, 2,424.85)                |                                    |
| P-value                             | < 0.0001                                |                                    |
| <b>OS</b> Median (months) (95% CI)  | 20.9 (20.9, NR)                         | 19.4 (16.9, NR)                    |
| Hazard ratio (95% CI)               | 0.74 (0.44, 1.25)                       |                                    |
| P-value                             | 0.27                                    |                                    |

CI: confidence interval; O: ofatumumab; n: number of responding subjects; N: number of subjects per group; NR: not reached. The analyses of PFS, overall response rate (ORR) and lymph node response rate (LNR) were based on evaluation by an independent review committee (IRC).

\* ORR defined as the proportion of subjects who achieved a complete response (CR) or partial response (PR) and maintained their response for at least 8 weeks.

\*\* LNR defined as the proportion of subjects who achieved a  $\geq 50\%$  decrease in the sum of products of the greatest perpendicular diameters of index lesions. Only subjects that had both baseline and  $\geq 1$  evaluable post-baseline assessments were included in this analysis.


**Table 6: Summary of PFS and response rates in pre-specified subgroups from study 312-0119**

|                                       | <b>Idelalisib + O</b><br><b>N = 70</b> | <b>Ofatumumab</b><br><b>N = 33</b> |
|---------------------------------------|--|------------------------------------|
| <b>17p deletion/TP53 mutation</b>     |  |                                    |
| PFS median (months) (95% CI)          | 13.7 (11.0, 17.8)                      | 5.8 (4.5, 8.4)                     |
| Hazard ratio (95% CI)                 | 0.32 (0.18, 0.57)                      |                                    |
| ORR (95% CI)                          | 72.9% (60.9, 82.8)                     | 15.2% (5.1, 31.9)                  |
| <b>Unmutated IGHV</b>                 | <b>N = 137</b>                         | <b>N = 68</b>                      |
| PFS median (months) (95% CI)          | 14.9 (12.4, 17.8)                      | 7.3 (5.3, 8.1)                     |
| Hazard ratio (95% CI)                 | 0.25 (0.17, 0.38)                      |                                    |
| ORR (95% CI)                          | 74.5% (66.3, 81.5)                     | 13.2% (6.2, 23.6)                  |
| <b>Age <math>\geq 65</math> years</b> | <b>N = 107</b>                         | <b>N = 60</b>                      |
| PFS median (months) (95% CI)          | 16.4 (13.4, 17.8)                      | 8.0 (5.6, 8.4)                     |
| Hazard ratio (95% CI)                 | 0.30 (0.19, 0.47)                      |                                    |
| ORR (95% CI)                          | 72.0% (62.5, 80.2)                     | 18.3% (9.5, 30.4)                  |

CI: confidence interval; O: ofatumumab; N: number of subjects per group



**Figure 2: Kaplan-Meier curve of PFS from study 312-0119 (intent-to-treat population)**



N at risk (Events)

|                |         |         |          |          |          |          |         |         |         |        |        |        |
|----------------|---------|---------|----------|----------|----------|----------|---------|---------|---------|--------|--------|--------|
| Idelalisib + O | 174 (0) | 162 (6) | 151 (13) | 140 (22) | 129 (31) | 110 (45) | 82 (57) | 44 (67) | 37 (70) | 7 (76) | 1 (76) | 0 (76) |
| Ofatumumab     | 87 (0)  | 60 (14) | 47 (21)  | 34 (30)  | 26 (34)  | 11 (49)  | 8 (51)  | 6 (52)  | 6 (52)  | 2 (54) | 0 (54) | 0 (54) |

Solid line: idelalisib + O (N = 174), dashed line: ofatumumab (N = 87)

O: ofatumumab; N: number of subjects per group

### Clinical efficacy in follicular lymphoma

The safety and efficacy of idelalisib were assessed in a single-arm, multicentre clinical study (study 101-09) conducted in 125 subjects with indolent B-cell non-Hodgkin lymphoma (iNHL, including: FL, n = 72; SLL, n = 28; lymphoplasmacytic lymphoma/Waldenström macroglobulinaemia [LPL/WM], n = 10; and marginal zone lymphoma [MZL], n = 15). All subjects were refractory to rituximab and 124 of 125 subjects were refractory to at least one alkylating agent. One hundred and twelve (89.6%) subjects were refractory to their last regimen prior to study entry.

Of the 125 subjects enrolled, 80 (64%) were male, the median age was 64 years (range: 33 to 87), and 110 (89%) were white. Subjects received 150 mg of idelalisib orally twice daily until evidence of disease progression or unacceptable toxicity.

The primary endpoint was the ORR defined as the proportion of subjects who achieved a CR or PR (based on the Revised Response Criteria for Malignant Lymphoma [Cheson]), and, for subjects with Waldenström macroglobulinaemia, a minor response (MR) (based on the Response Assessment for Waldenström macroglobulinaemia [Owen]). DOR was a secondary endpoint and was defined as the time from the first documented response (CR, PR, or MR) to the first documentation of disease progression or death from any cause. Efficacy results are summarised in Table 7.

**Table 7: Summary of efficacy in Study 101-09 (IRC assessment)**

| Characteristic                  | Overall iNHL cohort (N=125)<br>n (%) | FL subset (N=72)<br>n (%) |
|---------------------------------|--------------------------------------|---------------------------|
| ORR *                           | 72 (57.6%)                           | 40 (55.6%)                |
| 95% CI                          | 48.4 – 66.4                          | 43.4 – 67.3               |
| Response category*†             |                                      |                           |
| CR                              | 13 (10.4%)                           | 12 (16.7%)                |
| PR                              | 58 (46.4%)                           | 28 (38.9%)                |
| DOR (months)<br>median (95% CI) | 12.5 (7.4, 22.4)                     | 11.8 (6.2, 26.9)          |
| PFS (months)<br>median (95% CI) | 11.1 (8.3, 14.0)                     | 11.0 (8.0, 14.0)          |
| OS (months)<br>median (95% CI)  | 48.6 (33.9, 71.7)                    | 61.2 (38.1, NR)           |

CI: confidence interval; n: number of responding subjects

NR: not reached

\* Response as determined by an independent review committee (IRC) where ORR = complete response (CR) + partial response (PR) + minor response (MR) in subjects with WM.

† In the overall iNHL cohort, 1 subject (0.6%) with WM had the best overall response of MR

The median DOR for all subjects was 12.5 months (12.5 months for SLL subjects, and 11.8 months for FL, 20.4 months for LPL/WM and 18.4 months for MZL subjects). Among the 122 subjects with measurable lymph nodes at both baseline and post-baseline, 71 subjects (58.2%) achieved a  $\geq 50\%$  decrease from baseline in the sum of the products of the diameters (SPD) of index lesions. Of the 53 subjects who did not respond, 41 (32.8%) had stable disease 10 (8.0%) had progressive disease, and 2 (1.6%) were not evaluable. The median OS, including long-term follow-up for all 125 subjects, was 48.6 months. The median OS, including long-term follow-up for all FL subjects was 61.2 months.

### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with idelalisib in all subsets of the paediatric population in the treatment of mature B-cell neoplasms (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

### Absorption

Following oral administration of a single dose of idelalisib, peak plasma concentrations were observed 2 to 4 hours post-dose under fed conditions and after 0.5 to 1.5 hours under fasted conditions.

Following 150 mg twice daily administration of idelalisib, average (range)  $C_{max}$  and AUC at steady-state were 1,953 (272; 3,905) ng/mL and 10,439 (2,349; 29,315) ng•h/mL for idelalisib and 4,039 (669; 10,897) ng/mL and 39,744 (6,002; 119,770) ng•h/mL for GS-563117, respectively. The plasma exposures ( $C_{max}$  and AUC) of idelalisib are approximately dose proportional between 50 mg and 100 mg and less than dose proportional above 100 mg.

### *Effects of food*

Relative to fasting conditions, administration of an early capsule formulation of idelalisib with a high-fat meal resulted in no change in  $C_{max}$  and a 36% increase in mean  $AUC_{inf}$ . Idelalisib can be administered without regard to food.

### Distribution

Idelalisib is 93% to 94% bound to human plasma proteins at concentrations observed clinically. The mean blood-to-plasma concentration ratio was approximately 0.5. The apparent volume of distribution for idelalisib (mean) was approximately 96 L.

### Biotransformation

Idelalisib is metabolised primarily via aldehyde oxidase, and to a lesser extent via CYP3A and UGT1A4. The primary and only circulating metabolite, GS-563117, is inactive against PI3K $\delta$ .

### Elimination

The terminal elimination half-life of idelalisib was 8.2 (range: 1.9; 37.2) hours and the apparent clearance of idelalisib was 14.9 (range: 5.1; 63.8) L/h following idelalisib 150 mg twice daily oral administration. Following a single 150 mg oral dose of [<sup>14</sup>C]-labelled idelalisib, approximately 78% and 15% was excreted in faeces and urine, respectively. Unchanged idelalisib accounted for 23% of total radioactivity recovered in urine over 48 hours and 12% of total radioactivity recovered in faeces over 144 hours.

### In vitro interaction data

*In vitro* data indicated that idelalisib is not an inhibitor of the metabolising enzymes CYP1A2, CYP2B6, CYP2C9, CYP2C19, CYP2D6, CYP3A, or UGT1A1, or of the transporters OAT1, OAT3, or OCT2.

GS-563117 is not an inhibitor of the metabolising enzymes CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6 or UGT1A1, or of the transporters P-gp, BCRP, OATP1B1, OATP1B3, OAT1, OAT3, or OCT2.

### Special populations

#### *Gender and race*

Population pharmacokinetic analyses indicated that gender and race had no clinically relevant effect on the exposures to idelalisib or GS-563117.

#### *Elderly*

Population pharmacokinetic analyses indicated that age had no clinically relevant effect on the exposures to idelalisib or GS-563117, including elderly subjects (65 years of age and older), compared to younger subjects.

#### *Renal impairment*

A study of pharmacokinetics and safety of idelalisib was performed in healthy subjects and subjects with severe renal impairment (estimated CrCl 15 to 29 mL/min). Following a single 150 mg dose, no clinically relevant changes in exposures to idelalisib or GS-563117 were observed in subjects with severe renal impairment compared to healthy subjects.

#### *Hepatic impairment*

A study of pharmacokinetics and safety of idelalisib was performed in healthy subjects and subjects with moderate (Child-Pugh Class B) or severe (Child-Pugh Class C) hepatic impairment. Following a single 150 mg dose, idelalisib AUC (total, i.e., bound plus unbound) was ~60% higher in moderate and severe impairment compared to matched controls. The idelalisib AUC (unbound), after accounting for differences in protein binding, was ~80% (1.8-fold) higher in moderate and ~152% (2.5-fold) higher in severe impairment compared to matched controls.

#### *Paediatric population*

The pharmacokinetics of idelalisib in paediatric subjects has not been established (see section 4.2).

## **5.3 Preclinical safety data**

### Repeated dose toxicity

Idelalisib induced lymphoid depletion in spleen, thymus, lymph nodes and gut-associated lymphoid tissue. In general, B-lymphocyte dependent areas were more affected than T-lymphocyte dependent areas. In rats, idelalisib has the potential to inhibit T-dependent antibody responses. However, idelalisib did not inhibit the normal host response to *Staphylococcus aureus* and did not exacerbate the myelosuppressive effect of cyclophosphamide. Idelalisib is not considered to have broad immunosuppressive activity.

Idelalisib induced inflammatory changes in both rats and dogs. In studies up to 4 weeks in rats and dogs, hepatic necrosis was observed at 7 and 5 times the human exposure based on AUC, respectively. Serum transaminase elevations correlated with hepatic necrosis in dogs, but were not observed in rats. No hepatic impairment or chronic transaminase elevations were observed in rats or dogs in studies of 13 weeks and longer duration.

#### Genotoxicity

Idelalisib did not induce mutations in the microbial mutagenesis (Ames) assay, was not clastogenic in the *in vitro* chromosome aberration assay using human peripheral blood lymphocytes, and was not genotoxic in the *in vivo* rat micronucleus study.

#### Carcinogenicity

The carcinogenicity potential of idelalisib was evaluated in a 26-week transgenic RasH2 mouse study and a 2-year rat study. Idelalisib was not carcinogenic at exposures up to 1.4/7.9-fold (male/female) in mice compared to the exposure in patients with haematologic malignancies administered the recommended dose of 150 mg twice daily. A dose-related increase in pancreatic islet cell tumors was observed at low incidence in male rats at exposures up to 0.4-fold compared to the human exposure at the recommended dose; a similar finding was not observed in female rats at 0.62-fold exposure margin.

#### Reproductive and developmental toxicity

In an embryo-foetal development study in rats, increased post-implantation loss, malformations (absence of caudal vertebrae and in some cases also of sacral vertebrae), skeletal variations and lower foetal body weights were observed. Malformations were observed at exposures from 12 times the human exposure based on AUC. Effects on embryo-foetal development were not investigated in a second species.

Degeneration of the seminiferous tubules in the testes was observed in 2- to 13-week repeated dose studies in dogs and rats, but not in studies of 26 weeks and longer duration. In a rat male fertility study, decreases in epididymides and testes weight were observed but no adverse effects on mating or fertility parameters, and no degeneration or loss in spermatogenesis were observed. Female fertility was not affected in rats.

#### Phototoxicity

Evaluation of the potential for phototoxicity in the embryonic murine fibroblast cell line BALB/c 3T3 was inconclusive for idelalisib due to cytotoxicity in the *in vitro* assay. The major metabolite, GS-563117, may enhance phototoxicity when cells are simultaneously exposed to UVA light. There is a potential risk that idelalisib, via its major metabolite, GS-563117, may cause photosensitivity in treated patients.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Microcrystalline cellulose  
Hydroxypropyl cellulose (E463)  
Croscarmellose sodium  
Sodium starch glycolate  
Magnesium stearate

#### Film-coating

Polyvinyl alcohol (E1203)

Macrogol 3350 (E1521)

Titanium dioxide (E171)

Talc (E553B)

Sunset yellow FCF (E110)

#### **6.2 Incompatibilities**

Not applicable.

#### **6.3 Shelf life**

5 years.

#### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

#### **6.5 Nature and contents of container**

High density polyethylene (HDPE) bottle, capped with a polypropylene child-resistant closure, containing 60 film-coated tablets and a polyester coil.

Each carton contains 1 bottle.

#### **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

### **7. MARKETING AUTHORISATION HOLDER**

Gilead Sciences Ireland UC  
Carrigtohill  
County Cork, T45 DP77  
Ireland

### **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/938/001

### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 18 September 2014

Date of latest renewal: 30 April 2019

### **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## 1. NAME OF THE MEDICINAL PRODUCT

Zydelig 150 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 150 mg of idelalisib.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet.

Pink, oval-shaped, film-coated tablet of dimensions 10.0 mm by 6.8 mm, debossed on one side with “GSP” and “150” on the other side.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Zydelig is indicated in combination with rituximab for the treatment of adult patients with chronic lymphocytic leukaemia (CLL):

- who have received at least one prior therapy (see section 4.4), or
- as first line treatment in the presence of 17p deletion or *TP53* mutation in patients who are not eligible for any other therapies (see section 4.4).

Zydelig is indicated as monotherapy for the treatment of adult patients with follicular lymphoma (FL) that is refractory to two prior lines of treatment (see section 4.4).

### 4.2 Posology and method of administration

Treatment with Zydelig should be conducted by a physician experienced in the use of anti-cancer therapies.

#### Posology

The recommended dose is 150 mg idelalisib twice daily. Treatment should be continued until disease progression or unacceptable toxicity.

If the patient misses a dose of Zydelig within 6 hours of the time it is usually taken, the patient should take the missed dose as soon as possible and resume the normal dosing schedule. If a patient misses a dose by more than 6 hours, the patient should not take the missed dose and simply resume the usual dosing schedule.

#### Dose modification

##### *Elevated liver transaminases*

Treatment with Zydelig must be withheld in the event of a Grade 3 or 4 aminotransferase elevation (alanine aminotransferase [ALT]/aspartate aminotransferase [AST] > 5 x upper limit of normal [ULN]). Once values have returned to Grade 1 or below (ALT/AST ≤ 3 x ULN), treatment can be resumed at 100 mg twice daily.

If the event does not recur, the dose can be re-escalated to 150 mg twice daily at the discretion of the treating physician.

If the event recurs, treatment with Zydelig must be withheld until the values return to Grade 1 or less, after which re-initiation at 100 mg twice daily may be considered at the discretion of the physician (see sections 4.4 and 4.8).

*Diarrhoea/colitis*

Treatment with Zydelig must be withheld in the event of Grade 3 or 4 diarrhoea/colitis. Once diarrhoea/colitis has returned to Grade 1 or below, treatment can be resumed at 100 mg twice daily. If diarrhoea/colitis does not recur, the dose can be re-escalated to 150 mg twice daily at the discretion of the treating physician (see section 4.8).

*Pneumonitis*

Treatment with Zydelig must be withheld in the event of suspected pneumonitis. Once pneumonitis has resolved and if re-treatment is appropriate, resumption of treatment at 100 mg twice daily can be considered. Treatment with Zydelig must be permanently discontinued in the event of moderate or severe symptomatic pneumonitis or organising pneumonia (see sections 4.4 and 4.8).

*Rash*

Treatment with Zydelig must be withheld in the event of Grade 3 or 4 rash. Once rash has returned to Grade 1 or below, treatment can be resumed at 100 mg twice daily. If rash does not recur, the dose can be re-escalated to 150 mg twice daily at the discretion of the treating physician (see section 4.8).

*Neutropenia*

Treatment with Zydelig should be withheld in patients while absolute neutrophil count (ANC) is below 500 per mm<sup>3</sup>. ANC should be monitored at least weekly until ANC is ≥ 500 per mm<sup>3</sup> when treatment can be resumed at 100 mg twice daily (see section 4.4).

| ANC 1,000 to < 1,500/mm <sup>3</sup> | ANC 500 to < 1,000/mm <sup>3</sup>                           | ANC < 500/mm <sup>3</sup>  |
|--------------------------------------|--|--|
| Maintain Zydelig dosing.             | Maintain Zydelig dosing.<br><br>Monitor ANC at least weekly. | Interrupt Zydelig dosing.<br><br>Monitor ANC at least weekly until ANC ≥ 500/mm <sup>3</sup> , then may resume Zydelig dosing at 100 mg twice daily. |

*Special populations*

*Elderly*

No specific dose adjustment is required for elderly patients (aged ≥ 65 years) (see section 5.2).

*Renal impairment*

No dose adjustment is required for patients with mild, moderate, or severe renal impairment (see section 5.2).

*Hepatic impairment*

No dose adjustment is required when initiating treatment with Zydelig in patients with mild or moderate hepatic impairment, but an intensified monitoring of adverse reactions is recommended (see sections 4.4 and 5.2).

There is insufficient data to make dose recommendations for patients with severe hepatic impairment. Therefore, caution is recommended when administering Zydelig in this population and an intensified monitoring of adverse reactions is recommended (see sections 4.4 and 5.2).

*Paediatric population*

The safety and efficacy of Zydelig in children under the age of 18 years have not been established. No data are available.

#### Method of administration

Zydelig is for oral use. Patients should be instructed to swallow the tablet whole. The film-coated tablet should not be chewed or crushed. The film-coated tablet can be taken with or without food (see section 5.2).

#### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

#### **4.4 Special warnings and precautions for use**

##### Serious infections

Treatment with Zydelig should not be initiated in patients with any evidence of ongoing systemic bacterial, fungal, or viral infection.

Serious and fatal infections have occurred with idelalisib, including opportunistic infections such as *Pneumocystis jirovecii* pneumonia (PJP) and cytomegalovirus (CMV). Prophylaxis for PJP should therefore be administered to all patients throughout idelalisib treatment, and for a period of 2 to 6 months after discontinuation. The duration of post-treatment prophylaxis should be based on clinical judgment and may take into account a patient's risk factors such as concomitant corticosteroid treatment and prolonged neutropenia (see section 4.8).

Patients should be monitored for respiratory signs and symptoms throughout treatment. Patients should be advised to report new respiratory symptoms promptly.

Regular clinical and laboratory monitoring for CMV infection is recommended in patients with positive CMV serology at the start of treatment with idelalisib or with other evidence of a history of CMV infection. Patients with CMV viraemia without associated clinical signs of CMV infection should be carefully monitored. For patients with evidence of CMV viraemia and clinical signs of CMV infection, consideration should be given to interrupting idelalisib until the infection has resolved. If the benefits of resuming idelalisib are judged to outweigh the risks, consideration should be given to administering pre-emptive CMV therapy.

Cases of progressive multifocal leukoencephalopathy (PML) have been reported following the use of idelalisib within the context of prior or concomitant immunosuppressive therapies that have been associated with PML. Physicians should consider PML in the differential diagnosis in patients with new or worsening neurological, cognitive or behavioural signs or symptoms. If PML is suspected then appropriate diagnostic evaluations should be undertaken and treatment suspended until PML is excluded. If any doubt exists, referral to a neurologist and appropriate diagnostic measures for PML including MRI scan preferably with contrast, cerebrospinal fluid (CSF) testing for JC viral DNA and repeat neurological assessments should be considered.

##### Neutropenia

Treatment-emergent Grade 3 or 4 neutropenia, including febrile neutropenia, have occurred in patients treated with idelalisib. Blood counts should be monitored in all patients at least every 2 weeks for the first 6 months of treatment with idelalisib, and at least weekly in patients while ANC is less than 1,000 per mm<sup>3</sup> (see section 4.2).

##### Hepatotoxicity

Elevations in ALT and AST of Grade 3 and 4 (> 5 x ULN) have been observed in clinical studies of idelalisib. There have also been reports of hepatocellular injury including hepatic failure. Increases in liver transaminases were generally observed within the first 12 weeks of treatment, and were reversible with dose interruption (see section 4.2). In patients who resumed idelalisib at a lower dose, 26% had recurrence of ALT/AST elevation. Treatment with Zydelig must be withheld in the event of Grade 3 or 4 ALT/AST elevation and liver function monitored. Treatment may be resumed at a lower dose once values have returned to Grade 1 or below (ALT/AST ≤ 3 x ULN).



ALT, AST, and total bilirubin must be monitored in all patients every 2 weeks for the first 3 months of treatment, then as clinically indicated. If Grade 2 or higher elevations in ALT and/or AST are observed, patients' ALT, AST, and total bilirubin must be monitored weekly until the values return to Grade 1 or below.

#### Diarrhoea/colitis

Cases of severe drug-related colitis occurred relatively late (months) after the start of therapy, sometimes with rapid aggravation, but resolved within a few weeks with dose interruption and additional symptomatic treatment (e.g., anti-inflammatory agents such as enteric budesonide).

There is very limited experience from the treatment of patients with a history of inflammatory bowel disease.

#### Pneumonitis and organising pneumonia

Cases of pneumonitis and organising pneumonia (some with fatal outcome) have been reported with idelalisib. In patients presenting with serious lung events, idelalisib should be interrupted and the patient assessed for an explanatory aetiology. If either moderate or severe symptomatic pneumonitis or organising pneumonia is diagnosed, appropriate treatment should be initiated and idelalisib must be permanently discontinued.

#### Severe Cutaneous Reactions

Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS) have occurred with idelalisib. Cases of SJS and TEN with fatal outcomes have been reported when idelalisib was administered concomitantly with other medicinal products associated with these syndromes. If SJS, TEN or DRESS is suspected, idelalisib should be interrupted and the patient assessed and treated accordingly. If a diagnosis of SJS, TEN, or DRESS is confirmed, idelalisib should be permanently discontinued.

#### CYP3A inducers

Idelalisib exposure may be reduced when co-administered with CYP3A inducers such as rifampicin, phenytoin, St. John's wort (*Hypericum perforatum*), or carbamazepine. Since a reduction in idelalisib plasma concentrations may result in decreased efficacy, co-administration of Zydelig with moderate or strong CYP3A inducers should be avoided (see section 4.5).

#### CYP3A substrates

The primary metabolite of idelalisib, GS-563117, is a strong CYP3A4 inhibitor. Thus, idelalisib has the potential to interact with medicinal products that are metabolised by CYP3A, which may lead to increased serum concentrations of the other product (see section 4.5). When idelalisib is co-administered with other medicinal products, the Summary of Product Characteristics (SmPC) for the other product must be consulted for the recommendations regarding co-administration with CYP3A4 inhibitors. Concomitant treatment of idelalisib with CYP3A substrates with serious and/or life-threatening adverse reactions (e.g., alfuzosin, amiodarone, cisapride, pimozide, quinidine, ergotamine, dihydroergotamine, quetiapine, lovastatin, simvastatin, sildenafil, midazolam, triazolam) should be avoided and alternative medicinal products that are less sensitive to CYP3A4 inhibition should be used if possible.

#### Hepatic impairment

Intensified monitoring of adverse reactions is recommended in patients with impaired hepatic function as exposure is expected to be increased in this population, in particular in patients with severe hepatic impairment. No patients with severe hepatic impairment were included in clinical studies of idelalisib. Caution is recommended when administering Zydelig in this population.

#### Chronic hepatitis

Idelalisib has not been studied in patients with chronic active hepatitis including viral hepatitis. Caution should be exercised when administering Zydelig in patients with active hepatitis.

### Women of childbearing potential

Women of childbearing potential must use highly effective contraception while taking idelalisib and for 1 month after stopping treatment (see section 4.6). Women using hormonal contraceptives should add a barrier method as a second form of contraception since it is currently unknown whether idelalisib may reduce the effectiveness of hormonal contraceptives.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Idelalisib is metabolised primarily via aldehyde oxidase, and to a lesser extent via CYP3A and glucuronidation (UGT1A4). Its primary metabolite is GS-563117, which is not pharmacologically active. Idelalisib and GS-563117 are substrates of P-gp and BCRP.

### Effect of other medicinal products on idelalisib pharmacokinetics

#### *CYP3A inducers*

A clinical drug interaction study found that co-administration of a single dose of 150 mg idelalisib with rifampicin (a strong CYP3A inducer) resulted in a ~75% reduction in idelalisib AUC<sub>inf</sub>. Co-administration of Zydelig with moderate or strong CYP3A inducers such as rifampicin, phenytoin, St. John's wort, or carbamazepine should be avoided as this may result in decreased efficacy (see section 4.4).

#### *CYP3A/P-gp inhibitors*

A clinical drug interaction study found that co-administration of a single dose of 400 mg idelalisib with 400 mg once daily ketoconazole (a strong CYP3A, P-gp and BCRP inhibitor) resulted in a 26% increase in C<sub>max</sub> and a 79% increase in AUC<sub>inf</sub> of idelalisib. No initial dose adjustment of idelalisib is considered necessary when administered with CYP3A/P-gp inhibitors, but an intensified monitoring of adverse reactions is recommended.

### Effect of idelalisib on the pharmacokinetics of other medicinal products

#### *CYP3A substrates*

The primary metabolite of idelalisib, GS-563117, is a strong CYP3A inhibitor. A clinical drug interaction study found that co-administration of idelalisib with midazolam (a sensitive CYP3A substrate) resulted in a ~140% increase in C<sub>max</sub> and a ~440% increase in AUC<sub>inf</sub> of midazolam due to the CYP3A inhibition by GS-563117. Co-administration of idelalisib with CYP3A substrates may increase their systemic exposures and increase or prolong their therapeutic activity and adverse reactions. *In vitro*, the CYP3A4 inhibition was irreversible, and return to normal enzyme activity is therefore expected to take several days after stopping idelalisib administration.

Potential interactions between idelalisib and co-administered medicinal products that are CYP3A substrates are listed in Table 1 (increase is indicated as “↑”). This list is not exhaustive and is intended to serve as guidance only. In general, the SmPC for the other product must be consulted for the recommendations regarding co-administration with CYP3A4 inhibitors (see section 4.4).

**Table 1: Interactions between idelalisib and other medicinal products that are CYP3A substrates**

| Medicinal product                                       | Expected effect of idelalisib on drug levels | Clinical recommendation upon co-administration with idelalisib                                   |
|---|--|--|
| <b>ALPHA-1 ADRENORECEPTOR ANTAGONISTS</b>               |  |  |
| Alfuzosin   | ↑ serum concentrations                       | Idelalisib should not be co-administered with alfuzosin.   |
| <b>ANALGESICS</b>                                       |  |  |
| Fentanyl, alfentanil, methadone, buprenorphine/naloxone | ↑ serum concentrations                       | Careful monitoring of adverse reactions (e.g., respiratory depression, sedation) is recommended. |

| <b>Medicinal product</b>   | <b>Expected effect of idelalisib on drug levels</b> | <b>Clinical recommendation upon co-administration with idelalisib</b>   |
|--|---|---|
| <b>ANTIARRHYTHMICS</b>   |   |   |
| Amiodarone, quinidine  | ↑ serum concentrations                              | Idelalisib should not be co-administered with amiodarone or quinidine.  |
| Bepidil, disopyramide, lidocaine   | ↑ serum concentrations                              | Clinical monitoring is recommended.   |
| <b>ANTI-CANCER AGENTS</b>  |   |   |
| Tyrosine kinase inhibitors such as dasatinib and nilotinib, also vincristine and vinblastine | ↑ serum concentrations                              | Careful monitoring of the tolerance to these anti-cancer agents is recommended.   |
| <b>ANTICOAGULANTS</b>  |   |   |
| Warfarin   | ↑ serum concentrations                              | It is recommended that the international normalised ratio (INR) be monitored upon co-administration and following ceasing treatment with idelalisib.  |
| <b>ANTICONVULSANTS</b>   |   |   |
| Carbamazepine  | ↑ serum concentrations                              | Anticonvulsant drug levels should be monitored.   |
| <b>ANTIDEPRESSANTS</b>   |   |   |
| Trazodone  | ↑ serum concentrations                              | Careful dose titration of the antidepressant and monitoring for antidepressant response is recommended.   |
| <b>ANTI-GOUT</b>   |   |   |
| Colchicine   | ↑ serum concentrations                              | Dose reductions of colchicine may be required. Idelalisib should not be co-administered with colchicine to patients with renal or hepatic impairment. |
| <b>ANTI-HYPERTENSIVES</b>  |   |   |
| Amlodipine, diltiazem, felodipine, nifedipine, nicardipine                                   | ↑ serum concentrations                              | Clinical monitoring of therapeutic effect and adverse reactions is recommended.   |
| <b>ANTI-INFECTIVES</b>   |   |   |
| <b>Antifungals</b>   |   |   |
| Ketoconazole, itraconazole, posaconazole, voriconazole                                       | ↑ serum concentrations                              | Clinical monitoring is recommended.   |
| <b>Antimycobacterials</b>  |   |   |
| Rifabutin  | ↑ serum concentrations                              | Increased monitoring for rifabutin-associated adverse reactions including neutropenia and uveitis is recommended.                                     |
| <b>HCV protease inhibitors</b>   |   |   |
| Boceprevir, telaprevir   | ↑ serum concentrations                              | Clinical monitoring is recommended.   |

| <b>Medicinal product</b>                                  | <b>Expected effect of idelalisib on drug levels</b> | <b>Clinical recommendation upon co-administration with idelalisib</b>  |
|---|---|--|
| <b>Macrolide antibiotics</b>                              |   |  |
| Clarithromycin, telithromycin                             | ↑ serum concentrations                              | No dose adjustment of clarithromycin is required for patients with normal renal function or mild renal impairment (creatinine clearance [CrCl] 60-90 mL/min). Clinical monitoring is recommended for patients with CrCl < 90 mL/min. For patients with CrCl < 60 mL/min, alternative antibacterials should be considered.<br><br>Clinical monitoring is recommended for telithromycin. |
| <b>ANTI-PSYCHOTICS/NEUROLEPTICS</b>                       |   |  |
| Quetiapine, pimozone                                      | ↑ serum concentrations                              | Idelalisib should not be co-administered with quetiapine or pimozone.<br><br>Alternative medicinal products, such as olanzapine, may be considered.  |
| <b>ENDOTHELIN RECEPTOR ANTAGONISTS</b>                    |   |  |
| Bosentan  | ↑ serum concentrations                              | Caution should be exercised and patients closely observed for bosentan-related toxicity.   |
| <b>ERGOT ALKALOIDS</b>                                    |   |  |
| Ergotamine, dihydroergotamine                             | ↑ serum concentrations                              | Idelalisib should not be co-administered with ergotamine or dihydroergotamine.   |
| <b>GASTROINTESTINAL MOTILITY AGENTS</b>                   |   |  |
| Cisapride   | ↑ serum concentrations                              | Idelalisib should not be co-administered with cisapride.   |
| <b>GLUCOCORTICOIDS</b>                                    |   |  |
| Inhaled/nasal corticosteroids:<br>Budesonide, fluticasone | ↑ serum concentrations                              | Clinical monitoring is recommended.  |
| Oral budesonide   | ↑ serum concentrations                              | Clinical monitoring is recommended for increased signs/symptoms of corticosteroid effects.   |
| <b>HMG CO-A REDUCTASE INHIBITORS</b>                      |   |  |
| Lovastatin, simvastatin                                   | ↑ serum concentrations                              | Idelalisib should not be co-administered with lovastatin or simvastatin.   |
| Atorvastatin  | ↑ serum concentrations                              | Clinical monitoring is recommended and a lower starting dose of atorvastatin may be considered. Alternatively, switching to pravastatin, rosuvastatin, or pitavastatin may be considered.  |
| <b>IMMUNOSUPPRESSANTS</b>                                 |   |  |
| Ciclosporin, sirolimus, tacrolimus                        | ↑ serum concentrations                              | Therapeutic monitoring is recommended.   |

| Medicinal product   | Expected effect of idelalisib on drug levels | Clinical recommendation upon co-administration with idelalisib   |
|---|--|--|
| <b>INHALED BETA AGONIST</b>                                       |  |  |
| Salmeterol  | ↑ serum concentrations                       | Concurrent administration of salmeterol and idelalisib is not recommended. The combination may result in increased risk of cardiovascular adverse events associated with salmeterol, including QT prolongation, palpitations, and sinus tachycardia. |
| <b>PHOSPHODIESTERASE INHIBITORS</b>                               |  |  |
| Sildenafil  | ↑ serum concentrations                       | For pulmonary arterial hypertension:<br><br>Idelalisib should not be co-administered with sildenafil.  |
| Tadalafil   | ↑ serum concentrations                       | Caution should be exercised, including consideration of dose reduction, when co-administering tadalafil with idelalisib.<br><br>For erectile dysfunction:  |
| Sildenafil, tadalafil   | ↑ serum concentrations                       | Particular caution must be used and dose reduction may be considered when prescribing sildenafil or tadalafil with idelalisib with increased monitoring for adverse events.  |
| <b>SEDATIVES/HYPNOTICS</b>  |  |  |
| Midazolam (oral), triazolam                                       | ↑ serum concentrations                       | Idelalisib should not be co-administered with midazolam (oral) or triazolam.   |
| Buspirone, clorazepate, diazepam, estazolam, flurazepam, zolpidem | ↑ serum concentrations                       | Concentration monitoring of sedatives/hypnotics is recommended and dose reduction may be considered.   |

#### *CYP2C8 substrates*

*In vitro*, idelalisib both inhibited and induced CYP2C8, but it is not known whether this translates to an *in vivo* effect on CYP2C8 substrates. Caution is advised if Zydelig is used together with narrow therapeutic index drugs that are substrates of CYP2C8 (paclitaxel).

#### *Substrates of inducible enzymes (e.g., CYP2C9, CYP2C19, CYP2B6 and UGT)*

*In vitro*, idelalisib was an inducer of several enzymes, and a risk for decreased exposure and thereby decreased efficacy of substrates of inducible enzymes such as CYP2C9, CYP2C19, CYP2B6 and UGT cannot be excluded. Caution is advised if Zydelig is used together with narrow therapeutic index drugs that are substrates of these enzymes (warfarin, phenytoin, S-mephenytoin).

#### *BCRP, OATP1B1, OATP1B3 and P-gp substrates*

Co-administration of multiple doses of idelalisib 150 mg twice daily to healthy subjects resulted in comparable exposures for rosuvastatin (AUC 90% CI: 87, 121) and digoxin (AUC 90% CI: 98, 111), suggesting no clinically relevant inhibition of BCRP, OATP1B1/1B3 or systemic P-gp by idelalisib. A risk for P-gp inhibition in the gastrointestinal tract, that could result in increased exposure of sensitive substrates for intestinal P-gp such as dabigatran etexilate, cannot be excluded.

#### Paediatric population

Interaction studies have only been performed in adults.

### **4.6 Fertility, pregnancy and lactation**

#### Women of childbearing potential / contraception

Based on findings in animals, idelalisib may cause foetal harm. Women should avoid becoming pregnant while taking Zydelig, and for up to 1 month after ending treatment. Therefore, women of childbearing potential must use highly effective contraception while taking Zydelig and for 1 month after stopping treatment. It is currently unknown whether idelalisib may reduce the effectiveness of hormonal contraceptives, and therefore women using hormonal contraceptives should add a barrier method as a second form of contraception.

#### Pregnancy

There are no or limited amount of data from the use of idelalisib in pregnant women. Studies in animals have shown reproductive toxicity (see section 5.3).

Zydelig is not recommended during pregnancy and in women of childbearing potential not using contraception.

#### Breast-feeding

It is not known whether idelalisib and its metabolites are excreted in human milk.

A risk to the newborns/infants cannot be excluded.

Breast-feeding should be discontinued during treatment with Zydelig.

#### Fertility

No human data on the effect of idelalisib on fertility are available. Animal studies indicate the potential for harmful effects of idelalisib on fertility and foetal development (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Zydelig has no or negligible influence on the ability to drive and use machines.

### **4.8 Undesirable effects**

#### Summary of the safety profile

Assessment of adverse reactions is based on two Phase 3 studies (study 312-0116 and study 312-0119) and six Phase 1 and 2 studies. Study 312-0116 was a randomised, double-blind, placebo-controlled study in which 110 subjects with previously treated CLL received idelalisib + rituximab. In addition, 86 subjects from this study who were randomised to receive placebo + rituximab went on to receive idelalisib as a single agent in an extension study (study 312-0117). Study 312-0119 was a randomised, controlled, open-label study in which 173 subjects with previously treated CLL received idelalisib + ofatumumab. The Phase 1 and 2 studies assessed the safety of idelalisib in a total of 536 subjects with haematologic malignancies, including 400 subjects who received idelalisib (any dose) as a single agent and 136 subjects who received idelalisib in combination with an anti-CD20 monoclonal antibody (rituximab or ofatumumab).

#### Tabulated list of adverse reactions

The adverse drug reactions reported with idelalisib alone or in combination with anti-CD20 monoclonal antibodies (rituximab or ofatumumab) are provided in Table 2. Adverse reactions are listed by system organ class and frequency. Frequencies are defined as follows: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ), very rare ( $< 1/10,000$ ), and not known (cannot be estimated from available data).

**Table 2: Adverse drug reactions reported in clinical studies in subjects with haematologic malignancies receiving idelalisib and post marketing**

| <b>Reaction</b>   | <b>Any grade</b> | <b>Grade ≥ 3</b> |
|---|------------------|------------------|
| <i>Infections and infestations</i>                                      |                  |                  |
| Infections (including <i>Pneumocystis jirovecii</i> pneumonia and CMV)* | Very common      | Very common      |
| <i>Blood and lymphatic system disorders</i>                             |                  |                  |
| Neutropenia   | Very common      | Very common      |
| Lymphocytosis**   | Very common      | Very common      |
| <i>Respiratory, thoracic and mediastinal disorders</i>                  |                  |                  |
| Pneumonitis   | Common           | Common           |
| Organising pneumonia****  | Uncommon         | Uncommon         |
| <i>Gastrointestinal disorders</i>                                       |                  |                  |
| Diarrhoea/colitis   | Very common      | Very common      |
| <i>Hepatobiliary disorders</i>  |                  |                  |
| Transaminase increased  | Very common      | Very common      |
| Hepatocellular injury   | Common           | Common           |
| <i>Skin and subcutaneous tissue disorders</i>                           |                  |                  |
| Rash***   | Very common      | Common           |
| Stevens-Johnson syndrome/<br>toxic epidermal necrolysis****             | Rare             | Rare             |
| Drug reaction with<br>eosinophilia and systemic<br>symptoms (DRESS)**** | Not known        | Not known        |
| <i>General disorders and administration site conditions</i>             |                  |                  |
| Pyrexia   | Very common      | Common           |
| <i>Investigations</i>   |                  |                  |
| Increased triglycerides   | Very common      | Common           |

\* Comprised of opportunistic infections as well as bacterial and viral infections such as pneumonia, bronchitis, and sepsis.

\*\* Idelalisib-induced lymphocytosis should not be considered progressive disease in the absence of other clinical findings (see section 5.1).

\*\*\* Includes the preferred terms dermatitis exfoliative generalised, drug eruption, rash, rash erythematous, rash generalised, rash macular, rash maculo-papular, rash papular, rash pruritic, rash pustular, rash vesicular, papule, skin plaque, and exfoliative rash.

\*\*\*\* Observed in post-marketing data

#### Description of selected adverse reactions

##### *Infections (see section 4.4)*

Higher frequencies of infections overall, including Grade 3 and 4 infections, were observed in the idelalisib arms compared to the control arms of idelalisib clinical studies. Most frequently observed were infections in the respiratory system and septic events. In many instances the pathogen was not identified; however, both conventional and opportunistic pathogens, including PJP and CMV, were among those identified. Nearly all PJP infections, including fatal cases, occurred in the absence of PJP prophylaxis. There have been cases of PJP after stopping idelalisib treatment.

##### *Rash*

Rash was generally mild to moderate and resulted in discontinuation of treatment in 2.1% of subjects. In studies 312-0116/0117 and 312-0119, rash (reported as dermatitis exfoliative generalised, drug eruption, rash, rash erythematous, rash generalised, rash macular, rash maculo-papular, rash papular, rash pruritic, rash pustular, rash vesicular, papule and skin plaque) occurred in 31.1% of subjects who received idelalisib + an anti-CD20 monoclonal antibody (rituximab or ofatumumab) and 8.2% who received an anti-CD20 monoclonal antibody only (rituximab or ofatumumab). Of these, 5.7% who received idelalisib + an anti-CD20 monoclonal antibody (rituximab or ofatumumab) and 1.5% who

received an anti-CD20 monoclonal antibody only (rituximab or ofatumumab) had rash of Grade 3, and no subjects had an adverse reaction of Grade 4. Rash typically resolved with treatment (e.g., topical and/or oral steroids, diphenhydramine) and dose interruption for severe cases (see section 5.3, phototoxicity).

#### *Severe Cutaneous Reactions (see section 4.4)*

Cases of SJS, TEN and DRESS have occurred when idelalisib was administered concomitantly with other medicinal products associated with these syndromes (bendamustine, rituximab, allopurinol, amoxicillin, and sulfamethoxazole / trimethoprim). SJS or TEN occurred within one month of the medicinal combination and fatal outcomes have resulted.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

### **4.9 Overdose**

If overdose occurs the patient must be monitored for evidence of toxicity (see section 4.8). Treatment of overdose with Zydelig consists of general supportive measures including monitoring of vital signs as well as observation of the clinical status of the patient.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: antineoplastic agents, other antineoplastic agents, ATC code: **L01EM01**

#### Mechanism of action

Idelalisib inhibits phosphatidylinositol 3-kinase p110 $\delta$  (PI3K $\delta$ ), which is hyperactive in B-cell malignancies and is central to multiple signalling pathways that drive proliferation, survival, homing, and retention of malignant cells in lymphoid tissues and bone marrow. Idelalisib is a selective inhibitor of adenosine-5'-triphosphate (ATP) binding to the catalytic domain of PI3K $\delta$ , resulting in inhibition of the phosphorylation of the key lipid second messenger phosphatidylinositol and prevention of Akt (protein kinase B) phosphorylation.

Idelalisib induces apoptosis and inhibits proliferation in cell lines derived from malignant B-cells and in primary tumour cells. Through inhibition of chemokine receptors CXCR4 and CXCR5 signalling induced by the chemokines CXCL12 and CXCL13, respectively, idelalisib inhibits homing and retention of malignant B-cells in the tumour microenvironment including lymphoid tissues and the bone marrow.

No mechanistic explanations for the development of resistance to treatment with idelalisib have been identified from clinical studies. Further investigation of this topic in current B-cell malignancy studies is not planned.

#### Pharmacodynamic effects

##### *Electrocardiographic*

The effect of idelalisib (150 mg and 400 mg) on the QT/QTc interval was evaluated in a placebo- and positive-controlled (moxifloxacin 400 mg) crossover study in 40 healthy subjects. At a dose 2.7 times the maximum recommended dose, idelalisib did not prolong the QT/QTc interval (i.e., < 10 ms).



## Lymphocytosis

Upon initiation of idelalisib, a temporary increase in lymphocyte counts (i.e.,  $\geq 50\%$  increase from baseline and above absolute lymphocyte count of 5,000/ $\mu\text{L}$ ) has been observed. This occurs in approximately two-thirds of patients with CLL treated with idelalisib monotherapy and one-fourth of patients with CLL treated with idelalisib combination therapy. The onset of isolated lymphocytosis typically occurs during the first 2 weeks of idelalisib therapy and is often associated with reduction of lymphadenopathy. This observed lymphocytosis is a pharmacodynamic effect and should not be considered progressive disease in the absence of other clinical findings.

### Clinical efficacy in chronic lymphocytic leukaemia

#### *Idelalisib in combination with rituximab*

Study 312-0116 was a Phase 3, randomised, double-blind, placebo-controlled study in 220 subjects with previously treated CLL who required treatment but were not considered suitable for cytotoxic chemotherapy. Subjects were randomised 1:1 to receive 8 cycles of rituximab (first cycle at 375  $\text{mg}/\text{m}^2$  body surface area [BSA], subsequent cycles at 500  $\text{mg}/\text{m}^2$  BSA) in combination with either an oral placebo twice daily or with idelalisib 150  $\text{mg}$  taken twice daily until disease progression or unacceptable toxicity.

The median age was 71 years (range: 47 to 92) with 78.2% of subjects over 65 years; 65.5% were male, and 90.0% were white; 64.1% had a Rai stage of III or IV, and 55.9% had Binet Stage C. Most subjects had adverse cytogenetic prognostic factors: 43.2% had a 17p chromosomal deletion and/or tumour protein 53 (*TP53*) mutation, and 83.6% had unmutated genes for the immunoglobulin heavy chain variable region (*IGHV*). The median time from diagnosis of CLL to randomisation was 8.5 years. Subjects had a median Cumulative Illness Rating Scale (CIRS) score of 8. The median number of prior therapies was 3.0. Nearly all (95.9%) subjects had received prior anti-CD20 monoclonal antibodies. The primary endpoint was progression free survival (PFS). Efficacy results are summarised in Tables 3 and 4. The Kaplan-Meier curve for PFS is provided in Figure 1.

Compared with rituximab + placebo, treatment with idelalisib + rituximab resulted in statistically significant and clinically meaningful improvements in physical well-being, social well-being, functional well-being, as well as in the leukaemia-specific subscales of the Functional Assessment of Cancer Therapy: Leukaemia (FACT-LEU) instruments, and in statistically significant and clinically meaningful improvements in anxiety, depression and usual activities as measured by the EuroQoL Five-Dimensions (EQ-5D) instrument.

**Table 3: Efficacy results from study 312-0116**

|  | <b>Idelalisib + R</b><br><b>N = 110</b> | <b>Placebo + R</b><br><b>N = 110</b> |
|--|---|--------------------------------------|
| <b>PFS</b> Median (months) (95% CI)            | 19.4 (12.3, NR)                         | 6.5 (4.0, 7.3)                       |
| Hazard ratio (95% CI)                          | 0.15 (0.09, 0.24)                       |                                      |
| P-value  | < 0.0001                                |                                      |
| <b>ORR*</b> n (%) (95% CI)                     | 92 (83.6%) (75.4, 90.0)                 | 17 (15.5%) (9.3, 23.6)               |
| Odds ratio (95% CI)                            | 27.76 (13.40, 57.49)                    |                                      |
| P-value  | < 0.0001                                |                                      |
| <b>LNR**</b> n/N (%) (95% CI)                  | 102/106 (96.2%) (90.6, 99.0)            | 7/104 (6.7%) (2.7, 13.4)             |
| Odds ratio (95% CI)                            | 225.83 (65.56, 777.94)                  |                                      |
| P-value  | < 0.0001                                |                                      |
| <b>OS<sup>^</sup></b> Median (months) (95% CI) | NR (NR, NR)                             | 20.8 (14.8, NR)                      |
| Hazard ratio (95% CI)                          | 0.34 (0.19, 0.60)                       |                                      |
| P-value  | 0.0001                                  |                                      |

CI: confidence interval; R: rituximab; n: number of responding subjects; N: number of subjects per group; NR: not reached. The analyses of PFS, overall response rate (ORR) and lymph node response rate (LNR) were based on evaluation by an independent review committee (IRC).

\* ORR defined as the proportion of subjects who achieved a complete response (CR) or partial response (PR) based on the 2013 National Comprehensive Cancer Network (NCCN) response criteria and Cheson (2012).

\*\* LNR defined as the proportion of subjects who achieved a  $\geq 50\%$  decrease in the sum of products of the greatest perpendicular diameters of index lesions. Only subjects that had both baseline and  $\geq 1$  evaluable post-baseline assessments were included in this analysis.


^ Overall survival (OS) analysis includes data from subjects who received placebo + R on study 312-0116 and subsequently received idelalisib in an extension study, based on intent-to-treat analysis.

**Table 4: Summary of PFS and response rates in pre-specified subgroups from study 312-0116**

|  | <b>Idelalisib + R</b> | <b>Placebo + R</b> |
|--|-----------------------|--------------------|
| <b>17p deletion/<i>TP53</i> mutation</b> | <b>N = 46</b>         | <b>N = 49</b>      |
| PFS median (months) (95% CI)             | NR (12.3, NR)         | 4.0 (3.7, 5.7)     |
| Hazard ratio (95% CI)                    | 0.13 (0.07, 0.27)     |                    |
| ORR (95% CI)                             | 84.8% (71.1, 93.7)    | 12.2% (4.6, 24.8)  |
| <b>Unmutated <i>IGHV</i></b>             | <b>N = 91</b>         | <b>N = 93</b>      |
| PFS median (months) (95% CI)             | 19.4 (13.9, NR)       | 5.6 (4.0, 7.2)     |
| Hazard ratio (95% CI)                    | 0.14 (0.08, 0.23)     |                    |
| ORR (95% CI)                             | 82.4% (73.0, 89.6)    | 15.1% (8.5, 24.0)  |
| <b>Age <math>\geq 65</math> years</b>    | <b>N = 89</b>         | <b>N = 83</b>      |
| PFS median (months) (95% CI)             | 19.4 (12.3, NR)       | 5.7 (4.0, 7.3)     |
| Hazard ratio (95% CI)                    | 0.14 (0.08, 0.25)     |                    |
| ORR (95% CI)                             | 84.3% (75.0, 91.1)    | 16.9% (9.5, 26.7)  |

CI: confidence interval; R: rituximab; N: number of subjects per group; NR: not reached

**Figure 1: Kaplan-Meier curve of PFS from study 312-0116 (intent-to-treat population)**



N at risk (Events)

|                |         |         |         |         |         |         |         |        |        |        |        |        |
|----------------|---------|---------|---------|---------|---------|---------|---------|--------|--------|--------|--------|--------|
| Idelalisib + R | 110 (0) | 101 (3) | 93 (7)  | 73 (9)  | 59 (14) | 31 (19) | 20 (21) | 9 (24) | 7 (24) | 4 (24) | 1 (25) | 0 (25) |
| Placebo + R    | 110 (0) | 84 (21) | 48 (38) | 29 (46) | 20 (53) | 9 (63)  | 4 (67)  | 1 (69) | 0 (70) | 0 (70) | 0 (70) | 0 (70) |

Solid line: idelalisib + R (N = 110), dashed line: placebo + R (N = 110)

R: rituximab; N: number of subjects per group

The analysis of PFS was based on evaluation by an IRC. For subjects in the placebo + R group, the summary includes data up to the first dosing of idelalisib in an extension study.

Study 101-08/99 enrolled 64 subjects with previously untreated CLL, including 5 subjects with small lymphocytic lymphoma (SLL). Subjects received idelalisib 150 mg twice daily and rituximab 375 mg/m<sup>2</sup> BSA weekly for 8 doses. The ORR was 96.9%, with 12 CRs (18.8%) and 50 PRs (78.1%), including 3 CRs and 6 PRs in subjects with a 17p deletion and/or *TP53* mutation and 2 CRs and 34 PRs in subjects with unmutated *IGHV*. The median duration of response (DOR) has not been reached.

#### *Idelalisib in combination with ofatumumab*

Study 312-0119 was a Phase 3, randomised, open-label, multicentre, parallel-group study in 261 subjects with previously treated CLL who had measurable lymphadenopathy, required treatment, and experienced CLL progression < 24 months since the completion of the last prior therapy. Subjects were randomised 2:1 to receive idelalisib 150 mg twice daily and 12 infusions of ofatumumab over 24 weeks, or 12 infusions of ofatumumab only over 24 weeks. The first infusion of ofatumumab was administered at a dose of 300 mg and was continued at a dose of either 1,000 mg in the idelalisib + ofatumumab group or a dose of 2,000 mg in the ofatumumab only group, weekly for 7 doses, and then every 4 weeks for 4 doses. Idelalisib was taken until disease progression or unacceptable toxicity.

The median age was 68 years (range: 61 to 74) with 64.0% of subjects over 65 years; 71.3% were male, and 84.3% were white; 63.6% had a Rai stage of III or IV, and 58.2% had Binet Stage C. Most subjects had adverse cytogenetic prognostic factors: 39.5% had a 17p chromosomal deletion and/or *TP53* mutation, and 78.5% had unmutated genes for *IGHV*. The median time since diagnosis was 7.7 years. Subjects had a median CIRS score of 4. The median number of prior therapies was 3.0. The

primary endpoint was PFS. Efficacy results are summarised in Tables 5 and 6. The Kaplan-Meier curve for PFS is provided in Figure 2.

**Table 5: Efficacy results from study 312-0119**

|                                     | <b>Idelalisib + O</b><br><b>N = 174</b> | <b>Ofatumumab</b><br><b>N = 87</b> |
|-------------------------------------|---|------------------------------------|
| <b>PFS</b> Median (months) (95% CI) | 16.3 (13.6, 17.8)                       | 8.0 (5.7, 8.2)                     |
| Hazard ratio (95% CI)               | 0.27 (0.19, 0.39)                       |                                    |
| P-value                             | < 0.0001                                |                                    |
| <b>ORR*</b> n (%) (95% CI)          | 131 (75.3%) (68.2, 81.5)                | 16 (18.4%) (10.9, 28.1)            |
| Odds ratio (95% CI)                 | 15.94 (7.8, 32.58)                      |                                    |
| P-value                             | < 0.0001                                |                                    |
| <b>LNR**</b> n/N (%) (95% CI)       | 153/164 (93.3%) (88.3, 96.6)            | 4/81 (4.9%) (1.4, 12.2)            |
| Odds ratio (95% CI)                 | 486.96 (97.91, 2,424.85)                |                                    |
| P-value                             | < 0.0001                                |                                    |
| <b>OS</b> Median (months) (95% CI)  | 20.9 (20.9, NR)                         | 19.4 (16.9, NR)                    |
| Hazard ratio (95% CI)               | 0.74 (0.44, 1.25)                       |                                    |
| P-value                             | 0.27                                    |                                    |

CI: confidence interval; O: ofatumumab; n: number of responding subjects; N: number of subjects per group; NR: not reached. The analyses of PFS, overall response rate (ORR) and lymph node response rate (LNR) were based on evaluation by an independent review committee (IRC).

\* ORR defined as the proportion of subjects who achieved a complete response (CR) or partial response (PR) and maintained their response for at least 8 weeks.


\*\* LNR defined as the proportion of subjects who achieved a  $\geq 50\%$  decrease in the sum of products of the greatest perpendicular diameters of index lesions. Only subjects that had both baseline and  $\geq 1$  evaluable post-baseline assessments were included in this analysis.

**Table 6: Summary of PFS and response rates in pre-specified subgroups from study 312-0119**

|                                       | <b>Idelalisib + O</b><br><b>N = 70</b> | <b>Ofatumumab</b><br><b>N = 33</b> |
|---------------------------------------|--|------------------------------------|
| <b>17p deletion/TP53 mutation</b>     |  |                                    |
| PFS median (months) (95% CI)          | 13.7 (11.0, 17.8)                      | 5.8 (4.5, 8.4)                     |
| Hazard ratio (95% CI)                 | 0.32 (0.18, 0.57)                      |                                    |
| ORR (95% CI)                          | 72.9% (60.9, 82.8)                     | 15.2% (5.1, 31.9)                  |
| <b>Unmutated IGHV</b>                 | <b>N = 137</b>                         | <b>N = 68</b>                      |
| PFS median (months) (95% CI)          | 14.9 (12.4, 17.8)                      | 7.3 (5.3, 8.1)                     |
| Hazard ratio (95% CI)                 | 0.25 (0.17, 0.38)                      |                                    |
| ORR (95% CI)                          | 74.5% (66.3, 81.5)                     | 13.2% (6.2, 23.6)                  |
| <b>Age <math>\geq 65</math> years</b> | <b>N = 107</b>                         | <b>N = 60</b>                      |
| PFS median (months) (95% CI)          | 16.4 (13.4, 17.8)                      | 8.0 (5.6, 8.4)                     |
| Hazard ratio (95% CI)                 | 0.30 (0.19, 0.47)                      |                                    |
| ORR (95% CI)                          | 72.0% (62.5, 80.2)                     | 18.3% (9.5, 30.4)                  |

CI: confidence interval; O: ofatumumab; N: number of subjects per group

**Figure 2: Kaplan-Meier curve of PFS from study 312-0119 (intent-to-treat population)**



N at risk (Events)

|                |         |         |          |          |          |          |         |         |         |        |        |        |
|----------------|---------|---------|----------|----------|----------|----------|---------|---------|---------|--------|--------|--------|
| Idelalisib + O | 174 (0) | 162 (6) | 151 (13) | 140 (22) | 129 (31) | 110 (45) | 82 (57) | 44 (67) | 37 (70) | 7 (76) | 1 (76) | 0 (76) |
| Ofatumumab     | 87 (0)  | 60 (14) | 47 (21)  | 34 (30)  | 26 (34)  | 11 (49)  | 8 (51)  | 6 (52)  | 6 (52)  | 2 (54) | 0 (54) | 0 (54) |

Solid line: idelalisib + O (N = 174), dashed line: ofatumumab (N = 87)

O: ofatumumab; N: number of subjects per group

### Clinical efficacy in follicular lymphoma

The safety and efficacy of idelalisib were assessed in a single-arm, multicentre clinical study (study 101-09) conducted in 125 subjects with indolent B-cell non-Hodgkin lymphoma (iNHL, including: FL, n = 72; SLL, n = 28; lymphoplasmacytic lymphoma/Waldenström macroglobulinaemia [LPL/WM], n = 10; and marginal zone lymphoma [MZL], n = 15). All subjects were refractory to rituximab and 124 of 125 subjects were refractory to at least one alkylating agent. One hundred and twelve (89.6%) subjects were refractory to their last regimen prior to study entry.

Of the 125 subjects enrolled, 80 (64%) were male, the median age was 64 years (range: 33 to 87), and 110 (89%) were white. Subjects received 150 mg of idelalisib orally twice daily until evidence of disease progression or unacceptable toxicity.

The primary endpoint was the ORR defined as the proportion of subjects who achieved a CR or PR (based on the Revised Response Criteria for Malignant Lymphoma [Cheson]), and, for subjects with Waldenström macroglobulinaemia, a minor response (MR) (based on the Response Assessment for Waldenström macroglobulinaemia [Owen]). DOR was a secondary endpoint and was defined as the time from the first documented response (CR, PR, or MR) to the first documentation of disease progression or death from any cause. Efficacy results are summarised in Table 7.

**Table 7: Summary of efficacy in Study 101-09 (IRC assessment)**

| Characteristic                  | Overall iNHL cohort (N=125)<br>n (%) | FL subset (N=72)<br>n (%) |
|---------------------------------|--------------------------------------|---------------------------|
| ORR *                           | 72 (57.6%)                           | 40 (55.6%)                |
| 95% CI                          | 48.4 – 66.4                          | 43.4 – 67.3               |
| Response category*†             |                                      |                           |
| CR                              | 13 (10.4%)                           | 12 (16.7%)                |
| PR                              | 58 (46.4%)                           | 28 (38.9%)                |
| DOR (months)<br>median (95% CI) | 12.5 (7.4, 22.4)                     | 11.8 (6.2, 26.9)          |
| PFS (months)<br>median (95% CI) | 11.1 (8.3, 14.0)                     | 11.0 (8.0, 14.0)          |
| OS (months)<br>median (95% CI)  | 48.6 (33.9, 71.7)                    | 61.2 (38.1, NR)           |

CI: confidence interval; n: number of responding subjects

NR: not reached

\* Response as determined by an independent review committee (IRC) where ORR = complete response (CR) + partial response (PR) + minor response (MR) in subjects with WM.

† In the overall iNHL cohort, 1 subject (0.6%) with WM had the best overall response of MR

The median DOR for all subjects was 12.5 months (12.5 months for SLL subjects, and 11.8 months for FL, 20.4 months for LPL/WM and 18.4 months for MZL subjects). Among the 122 subjects with measurable lymph nodes at both baseline and post-baseline, 71 subjects (58.2%) achieved a  $\geq 50\%$  decrease from baseline in the sum of the products of the diameters (SPD) of index lesions. Of the 53 subjects who did not respond, 41 (32.8%) had stable disease, 10 (8.0%) had progressive disease, and 2 (1.6%) were not evaluable. The median OS, including long-term follow-up for all 125 subjects, was 48.6 months. The median OS, including long-term follow-up for all FL subjects was 61.2 months.

### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with idelalisib in all subsets of the paediatric population in the treatment of mature B-cell neoplasms (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

### Absorption

Following oral administration of a single dose of idelalisib, peak plasma concentrations were observed 2 to 4 hours post-dose under fed conditions and after 0.5 to 1.5 hours under fasted conditions.

Following 150 mg twice daily administration of idelalisib, average (range)  $C_{max}$  and AUC at steady-state were 1,953 (272; 3,905) ng/mL and 10,439 (2,349; 29,315) ng•h/mL for idelalisib and 4,039 (669; 10,897) ng/mL and 39,744 (6,002; 119,770) ng•h/mL for GS-563117, respectively. The plasma exposures ( $C_{max}$  and AUC) of idelalisib are approximately dose proportional between 50 mg and 100 mg and less than dose proportional above 100 mg.

### *Effects of food*

Relative to fasting conditions, administration of an early capsule formulation of idelalisib with a high-fat meal resulted in no change in  $C_{max}$  and a 36% increase in mean  $AUC_{inf}$ . Idelalisib can be administered without regard to food.

### Distribution

Idelalisib is 93% to 94% bound to human plasma proteins at concentrations observed clinically. The mean blood-to-plasma concentration ratio was approximately 0.5. The apparent volume of distribution for idelalisib (mean) was approximately 96 L.

### Biotransformation

Idelalisib is metabolised primarily via aldehyde oxidase, and to a lesser extent via CYP3A and UGT1A4. The primary and only circulating metabolite, GS-563117, is inactive against PI3K $\delta$ .

### Elimination

The terminal elimination half-life of idelalisib was 8.2 (range: 1.9; 37.2) hours and the apparent clearance of idelalisib was 14.9 (range: 5.1; 63.8) L/h following idelalisib 150 mg twice daily oral administration. Following a single 150 mg oral dose of [<sup>14</sup>C]-labelled idelalisib, approximately 78% and 15% was excreted in faeces and urine, respectively. Unchanged idelalisib accounted for 23% of total radioactivity recovered in urine over 48 hours and 12% of total radioactivity recovered in faeces over 144 hours.

### In vitro interaction data

*In vitro* data indicated that idelalisib is not an inhibitor of the metabolising enzymes CYP1A2, CYP2B6, CYP2C9, CYP2C19, CYP2D6, CYP3A, or UGT1A1, or of the transporters OAT1, OAT3, or OCT2.

GS-563117 is not an inhibitor of the metabolising enzymes CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6 or UGT1A1, or of the transporters P-gp, BCRP, OATP1B1, OATP1B3, OAT1, OAT3, or OCT2.

### Special populations

#### *Gender and race*

Population pharmacokinetic analyses indicated that gender and race had no clinically relevant effect on the exposures to idelalisib or GS-563117.

#### *Elderly*

Population pharmacokinetic analyses indicated that age had no clinically relevant effect on the exposures to idelalisib or GS-563117, including elderly subjects (65 years of age and older), compared to younger subjects.

#### *Renal impairment*

A study of pharmacokinetics and safety of idelalisib was performed in healthy subjects and subjects with severe renal impairment (estimated CrCl 15 to 29 mL/min). Following a single 150 mg dose, no clinically relevant changes in exposures to idelalisib or GS-563117 were observed in subjects with severe renal impairment compared to healthy subjects.

#### *Hepatic impairment*

A study of pharmacokinetics and safety of idelalisib was performed in healthy subjects and subjects with moderate (Child-Pugh Class B) or severe (Child-Pugh Class C) hepatic impairment. Following a single 150 mg dose, idelalisib AUC (total, i.e., bound plus unbound) was ~60% higher in moderate and severe impairment compared to matched controls. The idelalisib AUC (unbound), after accounting for differences in protein binding, was ~80% (1.8-fold) higher in moderate and ~152% (2.5-fold) higher in severe impairment compared to matched controls.

#### *Paediatric population*

The pharmacokinetics of idelalisib in paediatric subjects has not been established (see section 4.2).

## **5.3 Preclinical safety data**

### Repeated dose toxicity

Idelalisib induced lymphoid depletion in spleen, thymus, lymph nodes and gut-associated lymphoid tissue. In general, B-lymphocyte dependent areas were more affected than T-lymphocyte dependent areas. In rats, idelalisib has the potential to inhibit T-dependent antibody responses. However, idelalisib did not inhibit the normal host response to *Staphylococcus aureus* and did not exacerbate the myelosuppressive effect of cyclophosphamide. Idelalisib is not considered to have broad immunosuppressive activity.

Idelalisib induced inflammatory changes in both rats and dogs. In studies up to 4 weeks in rats and dogs, hepatic necrosis was observed at 7 and 5 times the human exposure based on AUC, respectively. Serum transaminase elevations correlated with hepatic necrosis in dogs, but were not observed in rats. No hepatic impairment or chronic transaminase elevations were observed in rats or dogs in studies of 13 weeks and longer duration.

#### Genotoxicity

Idelalisib did not induce mutations in the microbial mutagenesis (Ames) assay, was not clastogenic in the *in vitro* chromosome aberration assay using human peripheral blood lymphocytes, and was not genotoxic in the *in vivo* rat micronucleus study.

#### Carcinogenicity

The carcinogenicity potential of idelalisib was evaluated in a 26-week transgenic RasH2 mouse study and a 2-year rat study. Idelalisib was not carcinogenic at exposures up to 1.4/7.9-fold (male/female) in mice compared to the exposure in patients with haematologic malignancies administered the recommended dose of 150 mg twice daily. A dose-related increase in pancreatic islet cell tumors was observed at low incidence in male rats at exposures up to 0.4-fold compared to the human exposure at the recommended dose; a similar finding was not observed in female rats at 0.62-fold exposure margin.

#### Reproductive and developmental toxicity

In an embryo-foetal development study in rats, increased post-implantation loss, malformations (absence of caudal vertebrae and in some cases also of sacral vertebrae), skeletal variations and lower foetal body weights were observed. Malformations were observed at exposures from 12 times the human exposure based on AUC. Effects on embryo-foetal development were not investigated in a second species.

Degeneration of the seminiferous tubules in the testes was observed in 2- to 13-week repeated dose studies in dogs and rats, but not in studies of 26 weeks and longer duration. In a rat male fertility study, decreases in epididymides and testes weight were observed but no adverse effects on mating or fertility parameters, and no degeneration or loss in spermatogenesis were observed. Female fertility was not affected in rats.

#### Phototoxicity

Evaluation of the potential for phototoxicity in the embryonic murine fibroblast cell line BALB/c 3T3 was inconclusive for idelalisib due to cytotoxicity in the *in vitro* assay. The major metabolite, GS-563117, may enhance phototoxicity when cells are simultaneously exposed to UVA light. There is a potential risk that idelalisib, via its major metabolite, GS-563117, may cause photosensitivity in treated patients.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Microcrystalline cellulose  
Hydroxypropyl cellulose (E463)  
Croscarmellose sodium  
Sodium starch glycolate  
Magnesium stearate



#### Film-coating

Polyvinyl alcohol (E1203)

Macrogol 3350 (E1521)

Titanium dioxide (E171)

Talc (E553B)

Iron oxide red (E172)

#### **6.2 Incompatibilities**

Not applicable.

#### **6.3 Shelf life**

5 years.

#### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

#### **6.5 Nature and contents of container**

High density polyethylene (HDPE) bottle, capped with a polypropylene child-resistant closure, containing 60 film-coated tablets and a polyester coil.

Each carton contains 1 bottle.

#### **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

### **7. MARKETING AUTHORISATION HOLDER**

Gilead Sciences Ireland UC  
Carrigtohill  
County Cork, T45 DP77  
Ireland

### **8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/938/002

### **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 18 September 2014

Date of latest renewal: 30 April 2019

### **10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

## **ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer(s) responsible for batch release

Gilead Sciences Ireland UC  
IDA Business & Technology Park  
Carrigtohill  
County Cork  
Ireland

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The marketing authorisation holder (MAH) shall submit the first PSUR for this product within 8 months following authorisation. Subsequently, the MAH shall submit PSURs for this product in accordance with the requirements set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

If the dates for submission of a PSUR and the update of a RMP coincide, they can be submitted at the same time.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Zydelig 100 mg film-coated tablets  
idelalisib

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 100 mg of idelalisib.

**3. LIST OF EXCIPIENTS**

Contains sunset yellow FCF (E110), see leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablets  
60 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Gilead Sciences Ireland UC  
Carrigtohill  
County Cork, T45 DP77  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/938/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Zydelig 100 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Zydelig 100 mg film-coated tablets  
idelalisib

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 100 mg of idelalisib.

**3. LIST OF EXCIPIENTS**

Contains sunset yellow FCF (E110), see leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablets  
60 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**



**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Gilead Sciences Ireland UC  
Carrigtohill  
County Cork, T45 DP77  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/938/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Zydelig 150 mg film-coated tablets  
idelalisib

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 150 mg of idelalisib.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablets  
60 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Gilead Sciences Ireland UC  
Carrigtohill  
County Cork, T45 DP77  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/938/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Zydelig 150 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Zydelig 150 mg film-coated tablets  
idelalisib

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 150 mg of idelalisib.

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablets  
60 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Read the package leaflet before use.

Oral use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Gilead Sciences Ireland UC  
Carrigtohill  
County Cork, T45 DP77  
Ireland

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/14/938/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

**B. PACKAGE LEAFLET**

## Package leaflet: Information for the patient

### Zydelig 100 mg film-coated tablets idelalisib

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Zydelig is and what it is used for
2. What you need to know before you take Zydelig
3. How to take Zydelig
4. Possible side effects
5. How to store Zydelig
6. Contents of the pack and other information

#### 1. What Zydelig is and what it is used for

Zydelig is a cancer medicine that contains the active substance idelalisib. It works by blocking the effects of an enzyme involved in multiplication and survival of certain white blood cells called lymphocytes. Because this enzyme is overactivated in certain cancerous white blood cells, by blocking it, Zydelig will kill and reduce the number of cancer cells.

Zydelig may be used for the treatment of two different cancers in adults:

##### Chronic lymphocytic leukaemia

*Chronic lymphocytic leukaemia* (CLL) is a cancer of a type of white blood cell called B-lymphocytes. In this disease, the lymphocytes multiply too quickly and live for too long, so that there are too many of them circulating in the blood.

In CLL Zydelig treatment is used in combination with another medicine (rituximab) in patients who have certain high-risk factors or in patients whose cancer has come back after at least one previous treatment.

##### Follicular lymphoma

*Follicular lymphoma* (FL) is a cancer of a type of white blood cell called B-lymphocytes. In follicular lymphoma, the B-lymphocytes multiply too quickly and live for too long, so there are too many of them in the lymph nodes. In FL Zydelig is used on its own in patients whose cancer has not responded to treatment with two previous cancer treatments.

## 2. What you need to know before you take Zydelig

### Do not take Zydelig

- if you are **allergic** to idelalisib or any of the other ingredients of this medicine (listed in section 6).  
→ **Talk to your doctor** if this applies to you.

### Warnings and precautions

Talk to your doctor before taking Zydelig. Tell your doctor:

- if you have liver problems
- if you have any other medical conditions or illness (especially an infection or fever)

Serious and fatal infections have occurred in patients taking Zydelig. You should take additional medicine provided by your doctor while you are taking Zydelig to prevent one type of infection. Your doctor will monitor you for evidence of infection. Tell your doctor right away if you become ill (especially with a fever, cough or breathing difficulties) while you are taking Zydelig.

Tell your doctor immediately if you notice or someone notices in you: memory loss, trouble thinking, difficulty walking or sight loss – these may be due to a very rare but serious brain infection which can be fatal (progressive multifocal leukoencephalopathy or PML).

**You will need regular blood tests** before and during treatment with Zydelig. This is to check that you do not have an infection, that your liver is working properly, and that your blood counts are normal. If necessary, your doctor may decide to stop treatment for a while, before starting treatment again at the same or a lower dose. Your doctor may also decide to permanently stop treatment with Zydelig.

Zydelig can cause severe diarrhoea. Tell your doctor right away at the first sign of diarrhoea.

Zydelig can cause lung inflammation. Tell your doctor right away:

- if you have a new or worsening cough
- if you have shortness of breath or difficulty breathing

Severe skin blistering conditions including Stevens-Johnson syndrome and toxic epidermal necrolysis, and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in association with idelalisib treatment. Stop using idelalisib and seek medical attention immediately if you notice any of the symptoms described in section 4.

Tell your doctor right away:

- if you have redness and blistering of the skin
- if you have swelling and blistering of the lining of the mouth, throat, nose, genitals, and/or eyes

Laboratory tests may show an increase in white blood cells (called “lymphocytes”) in your blood in the first few weeks of treatment. This is expected and may last for a few months. This generally does not mean that your blood cancer is getting worse. Your doctor will check your blood counts before or during treatment with Zydelig and in rare cases they may need to give you another medicine. Talk to your doctor about what your test results mean.

### Children and adolescents

Do not give this medicine to children and adolescents under 18 years of age because it has not been studied in this age group.



## Other medicines and Zydelig

**Zydelig should not be used with any other medicines unless your doctor has told you it is safe to do so.**

**Tell your doctor if you are taking, have recently taken or might take any other medicines.** This is extremely important, as using more than one medicine at the same time can strengthen or weaken their effect.

Taking Zydelig with certain medicines may stop them working properly, or may make side effects worse. In particular, tell your doctor if you are taking any of the following:

- **alfuzosin**, a medicine used to treat an enlarged prostate
- **dabigatran, warfarin**, medicines used to thin the blood
- **amiodarone, bepridil, disopyramide, lidocaine, quinidine**, medicines used to treat heart problems
- **dihydroergotamine, ergotamine**, medicines used to treat migraine headache
- **cisapride**, a medicine used to relieve certain stomach problems
- **pimozide**, a medicine used to treat abnormal thoughts or feelings
- **midazolam, triazolam**, when taken by mouth to help you sleep and/or relieve anxiety
- **quetiapine**, a medicine used to treat schizophrenia, bipolar disorder and major depressive disorder
- **amlodipine, diltiazem, felodipine, nicardipine, nifedipine**, medicines used to treat high blood pressure and heart problems
- **bosentan**, a medicine used to treat pulmonary arterial hypertension
- **sildenafil, tadalafil**, medicines used to treat impotence and pulmonary hypertension, a lung disease that makes breathing difficult
- **budesonide, fluticasone**, medicines used to treat hayfever and asthma, and **salmeterol**, used to treat asthma
- **rifabutin**, a medicine used to treat bacterial infections including tuberculosis
- **itraconazole, ketoconazole, posaconazole, voriconazole**, medicines used to treat fungal infections
- **boceprevir, telaprevir**, medicines used to treat hepatitis C
- **carbamazepine, S-mephenytoin, phenytoin**, medicines used to prevent seizures
- **rifampicin**, a medicine used to prevent and treat tuberculosis and other infections
- **St. John's wort (*Hypericum perforatum*)**, a herbal remedy used for depression and anxiety
- **alfentanil, fentanyl, methadone, buprenorphine/naloxone**, medicines used for pain relief
- **ciclosporin, sirolimus, tacrolimus**, medicines used to control your body's immune response after a transplant
- **colchicine**, a medicine used to treat gout
- **trazodone**, a medicine used to treat depression
- **buspirone, clorazepate, diazepam, estazolam, flurazepam, zolpidem**, medicines used to treat nervous system disorders
- **dasatinib, nilotinib, paclitaxel, vinblastine, vincristine**, medicines used to treat cancer
- **oral or implanted hormonal contraceptives**, used to prevent pregnancy
- **clarithromycin, telithromycin**, medicines used to treat bacterial infections
- **atorvastatin, lovastatin, simvastatin**, medicines used to lower cholesterol

Zydelig may be prescribed in combination with other medicines for the treatment of CLL. It is very important that you read the package leaflets that are provided with these medicines too.

Ask your doctor if you have any questions about any of your medicines.

## Pregnancy and breast-feeding

- **Zydelig should not be used during pregnancy.** There is no information about the safety of this medicine in pregnant women.
- **Use a reliable method of contraception** to avoid becoming pregnant while you are being treated with Zydelig, and for 1 month after your last treatment.
- **Zydelig may make the contraceptive “pill” and implanted hormonal contraceptives work less well.** You must also use a barrier method of contraception such as condoms or the “coil” while taking Zydelig and for 1 month after your last treatment.
- **Tell your doctor immediately if you become pregnant.**

**You should not breast-feed while taking Zydelig.** If you are currently breast-feeding, talk to your doctor before starting treatment. It is not known whether the active substance in Zydelig passes into human milk.

## Driving and using machines

Zydelig is unlikely to affect your ability to drive or use machines.

## Zydelig contains sunset yellow FCF (E110)

Tell your doctor if you have an allergy to sunset yellow FCF (E110). Zydelig contains sunset yellow FCF which may cause allergic reactions.

## 3. How to take Zydelig

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

**The recommended dose** is 150 mg by mouth twice a day. However, your doctor may reduce this dose to 100 mg twice a day if you experience particular side effects.

Zydelig can be taken with or without food.

**Swallow the tablet whole.** Do not chew or crush the tablet. Tell your doctor if you have problems swallowing tablets.

### If you take more Zydelig than you should

If you accidentally take more than the recommended dose of Zydelig, you may be at increased risk of side effects with this medicine (see section 4, *Possible side effects*).

Contact your doctor or nearest emergency department immediately for advice. Keep the bottle and this leaflet with you so that you can easily describe what you have taken.

### If you forget to take Zydelig

Take care to not miss a dose of Zydelig. If you miss a dose by less than 6 hours, take the missed dose right away. Then take your next dose as usual. If you miss a dose by more than 6 hours, wait and take the next dose at your usual time.

### Do not stop taking Zydelig

Do not stop taking this medicine unless your doctor tells you to.

If you have any further questions on the use of this medicine, ask your doctor.

#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Some side effects could be serious.**

**STOP taking Zydelig and seek medical help immediately if you experience any of the following:**

- reddish patches on the trunk, small circumscribed changes in the colour of the skin, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (Stevens-Johnson syndrome, toxic epidermal necrolysis).
- Widespread rash, high body temperature and enlarged lymph nodes (DRESS syndrome).

**Other side effects**

**Very common side effects**

*(may affect more than 1 in 10 people)*

- diarrhoea/inflammation of the large intestine
- rash
- changes in the number of white blood cells
- infections
- fever

*Blood tests may also show:*

- increased blood levels of liver enzymes

**Common side effects**

*(may affect up to 1 in 10 people)*

- inflammation of the lungs
- liver damage

*Blood tests may also show:*

- increased blood levels of fats

**Reporting of side effects**

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in Appendix V. By reporting side effects you can help provide more information on the safety of this medicine.

#### 5. How to store Zydelig

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the bottle and carton after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Zydelig contains

- **The active substance is** idelalisib. Each film-coated tablet contains 100 mg of idelalisib.
- **The other ingredients are:**  
*Tablet core:*  
Microcrystalline cellulose, hydroxypropyl cellulose (E463), croscarmellose sodium, sodium starch glycolate, magnesium stearate.  
  
*Film-coating:*  
Polyvinyl alcohol (E1203), macrogol 3350 (E1521), titanium dioxide (E171), talc (E553B), sunset yellow FCF (E110) (see Section 2, *What you need to know before you take Zydelig*).

### What Zydelig looks like and contents of the pack

The film-coated tablets are orange, oval-shaped tablets, debossed on one side with “GSP” and “100” on the other side.

The following pack size is available: outer carton containing 1 plastic bottle of 60 film-coated tablets.

### Marketing Authorisation Holder

Gilead Sciences Ireland UC  
Carrigtohill  
County Cork, T45 DP77  
Ireland

### Manufacturer

Gilead Sciences Ireland UC  
IDA Business & Technology Park  
Carrigtohill  
County Cork  
Ireland

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **België/Belgique/Belgien**

Gilead Sciences Belgium SRL-BV  
Tél/Tel: + 32 (0) 24 01 35 50

#### **Lietuva**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

#### **България**

Gilead Sciences Ireland UC  
Тел.: + 353 (0) 1 686 1888

#### **Luxembourg/Luxemburg**

Gilead Sciences Belgium SRL-BV  
Tél/Tel: + 32 (0) 24 01 35 50

#### **Česká republika**

Gilead Sciences s.r.o.  
Tel: + 420 910 871 986

#### **Magyarország**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

#### **Danmark**

Gilead Sciences Sweden AB  
Tlf: + 46 (0) 8 5057 1849

#### **Malta**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

**Deutschland**

Gilead Sciences GmbH  
Tel: + 49 (0) 89 899890-0

**Eesti**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

**Ελλάδα**

Gilead Sciences Ελλάς Μ.ΕΠΕ.  
Τηλ: + 30 210 8930 100

**España**

Gilead Sciences, S.L.  
Tel: + 34 91 378 98 30

**France**

Gilead Sciences  
Tél: + 33 (0) 1 46 09 41 00

**Hrvatska**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

**Ireland**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 214 825 999

**Ísland**

Gilead Sciences Sweden AB  
Sími: + 46 (0) 8 5057 1849

**Italia**

Gilead Sciences S.r.l.  
Tel: + 39 02 439201

**Κύπρος**

Gilead Sciences Ελλάς Μ.ΕΠΕ.  
Τηλ: + 30 210 8930 100

**Latvija**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

**Nederland**

Gilead Sciences Netherlands B.V.  
Tel: + 31 (0) 20 718 36 98

**Norge**

Gilead Sciences Sweden AB  
Tlf: + 46 (0) 8 5057 1849

**Österreich**

Gilead Sciences GesmbH  
Tel: + 43 1 260 830

**Polska**

Gilead Sciences Poland Sp. z o.o.  
Tel: + 48 22 262 8702

**Portugal**

Gilead Sciences, Lda.  
Tel: + 351 21 7928790

**România**

Gilead Sciences (GSR) S.R.L.  
Tel: + 40 31 631 18 00

**Slovenija**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

**Slovenská republika**

Gilead Sciences Slovakia s.r.o.  
Tel: + 421 232 121 210

**Suomi/Finland**

Gilead Sciences Sweden AB  
Puh/Tel: + 46 (0) 8 5057 1849

**Sverige**

Gilead Sciences Sweden AB  
Tel: + 46 (0) 8 5057 1849

**United Kingdom (Northern Ireland)**

Gilead Sciences Ireland UC  
Tel: + 44 (0) 8000 113700

**This leaflet was last revised in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.

## Package leaflet: Information for the patient

### Zydelig 150 mg film-coated tablets idelalisib

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Zydelig is and what it is used for
2. What you need to know before you take Zydelig
3. How to take Zydelig
4. Possible side effects
5. How to store Zydelig
6. Contents of the pack and other information

#### 1. What Zydelig is and what it is used for

Zydelig is a cancer medicine that contains the active substance idelalisib. It works by blocking the effects of an enzyme involved in multiplication and survival of certain white blood cells called lymphocytes. Because this enzyme is overactivated in certain cancerous white blood cells, by blocking it, Zydelig will kill and reduce the number of cancer cells.

Zydelig may be used for the treatment of two different cancers in adults:

##### Chronic lymphocytic leukaemia

*Chronic lymphocytic leukaemia* (CLL) is a cancer of a type of white blood cell called B-lymphocytes. In this disease, the lymphocytes multiply too quickly and live for too long, so that there are too many of them circulating in the blood.

In CLL Zydelig treatment is used in combination with another medicine (rituximab) in patients who have certain high-risk factors or in patients whose cancer has come back after at least one previous treatment.

##### Follicular lymphoma

*Follicular lymphoma* (FL) is a cancer of a type of white blood cell called B-lymphocytes. In follicular lymphoma, the B-lymphocytes multiply too quickly and live for too long, so there are too many of them in the lymph nodes. In FL Zydelig is used on its own in patients whose cancer has not responded to treatment with two previous cancer treatments.

## 2. What you need to know before you take Zydelig

### Do not take Zydelig

- if you are **allergic** to idelalisib or any of the other ingredients of this medicine (listed in section 6).  
→ **Talk to your doctor** if this applies to you.

### Warnings and precautions

Talk to your doctor before taking Zydelig. Tell your doctor:

- if you have liver problems
- if you have any other medical conditions or illness (especially an infection or fever)

Serious and fatal infections have occurred in patients taking Zydelig. You should take additional medicine provided by your doctor while you are taking Zydelig to prevent one type of infection. Your doctor will monitor you for evidence of infection. Tell your doctor right away if you become ill (especially with a fever, cough or breathing difficulties) while you are taking Zydelig.

Tell your doctor immediately if you notice or someone notices in you: memory loss, trouble thinking, difficulty walking or sight loss – these may be due to a very rare but serious brain infection which can be fatal (progressive multifocal leukoencephalopathy or PML).

**You will need regular blood tests** before and during treatment with Zydelig. This is to check that you do not have an infection, that your liver is working properly, and that your blood counts are normal. If necessary, your doctor may decide to stop treatment for a while, before starting treatment again at the same or a lower dose. Your doctor may also decide to permanently stop treatment with Zydelig.

Zydelig can cause severe diarrhoea. Tell your doctor right away at the first sign of diarrhoea.

Zydelig can cause lung inflammation. Tell your doctor right away:

- if you have a new or worsening cough
- if you have shortness of breath or difficulty breathing

Severe skin blistering conditions including Stevens-Johnson syndrome and toxic epidermal necrolysis, and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in association with idelalisib treatment. Stop using idelalisib and seek medical attention immediately if you notice any of the symptoms described in section 4.

Tell your doctor right away:

- if you have redness and blistering of the skin
- if you have swelling and blistering of the lining of the mouth, throat, nose, genitals, and/or eyes

Laboratory tests may show an increase in white blood cells (called “lymphocytes”) in your blood in the first few weeks of treatment. This is expected and may last for a few months. This generally does not mean that your blood cancer is getting worse. Your doctor will check your blood counts before or during treatment with Zydelig and in rare cases they may need to give you another medicine. Talk to your doctor about what your test results mean.

### Children and adolescents

Do not give this medicine to children and adolescents under 18 years of age because it has not been studied in this age group.

## Other medicines and Zydelig

**Zydelig should not be used with any other medicines unless your doctor has told you it is safe to do so.**

**Tell your doctor if you are taking, have recently taken or might take any other medicines.** This is extremely important, as using more than one medicine at the same time can strengthen or weaken their effect.

Taking Zydelig with certain medicines may stop them working properly, or may make side effects worse. In particular, tell your doctor if you are taking any of the following:

- **alfuzosin**, a medicine used to treat an enlarged prostate
- **dabigatran, warfarin**, medicines used to thin the blood
- **amiodarone, bepridil, disopyramide, lidocaine, quinidine**, medicines used to treat heart problems
- **dihydroergotamine, ergotamine**, medicines used to treat migraine headache
- **cisapride**, a medicine used to relieve certain stomach problems
- **pimozide**, a medicine used to treat abnormal thoughts or feelings
- **midazolam, triazolam**, when taken by mouth to help you sleep and/or relieve anxiety
- **quetiapine**, a medicine used to treat schizophrenia, bipolar disorder and major depressive disorder
- **amlodipine, diltiazem, felodipine, nicardipine, nifedipine**, medicines used to treat high blood pressure and heart problems
- **bosentan**, a medicine used to treat pulmonary arterial hypertension
- **sildenafil, tadalafil**, medicines used to treat impotence and pulmonary hypertension, a lung disease that makes breathing difficult
- **budesonide, fluticasone**, medicines used to treat hayfever and asthma, and **salmeterol**, used to treat asthma
- **rifabutin**, a medicine used to treat bacterial infections including tuberculosis
- **itraconazole, ketoconazole, posaconazole, voriconazole**, medicines used to treat fungal infections
- **boceprevir, telaprevir**, medicines used to treat hepatitis C
- **carbamazepine, S-mephenytoin, phenytoin**, medicines used to prevent seizures
- **rifampicin**, a medicine used to prevent and treat tuberculosis and other infections
- **St. John's wort (*Hypericum perforatum*)**, a herbal remedy used for depression and anxiety
- **alfentanil, fentanyl, methadone, buprenorphine/naloxone**, medicines used for pain relief
- **ciclosporin, sirolimus, tacrolimus**, medicines used to control your body's immune response after a transplant
- **colchicine**, a medicine used to treat gout
- **trazodone**, a medicine used to treat depression
- **buspirone, clorazepate, diazepam, estazolam, flurazepam, zolpidem**, medicines used to treat nervous system disorders
- **dasatinib, nilotinib, paclitaxel, vinblastine, vincristine**, medicines used to treat cancer
- **oral or implanted hormonal contraceptives**, used to prevent pregnancy
- **clarithromycin, telithromycin**, medicines used to treat bacterial infections
- **atorvastatin, lovastatin, simvastatin**, medicines used to lower cholesterol

Zydelig may be prescribed in combination with other medicines for the treatment of CLL. It is very important that you read the package leaflets that are provided with these medicines too.

Ask your doctor if you have any questions about any of your medicines.



## **Pregnancy and breast-feeding**

- **Zydelig should not be used during pregnancy.** There is no information about the safety of this medicine in pregnant women.
- **Use a reliable method of contraception** to avoid becoming pregnant while you are being treated with Zydelig, and for 1 month after your last treatment.
- **Zydelig may make the contraceptive “pill” and implanted hormonal contraceptives work less well.** You must also use a barrier method of contraception such as condoms or the “coil” while taking Zydelig and for 1 month after your last treatment.
- **Tell your doctor immediately if you become pregnant.**

**You should not breast-feed while taking Zydelig.** If you are currently breast-feeding, talk to your doctor before starting treatment. It is not known whether the active substance in Zydelig passes into human milk.

## **Driving and using machines**

Zydelig is unlikely to affect your ability to drive or use machines.

## **3. How to take Zydelig**

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

**The recommended dose** is 150 mg by mouth twice a day. However, your doctor may reduce this dose to 100 mg twice a day if you experience particular side effects.

Zydelig can be taken with or without food.

**Swallow the tablet whole.** Do not chew or crush the tablet. Tell your doctor if you have problems swallowing tablets.

### **If you take more Zydelig than you should**

If you accidentally take more than the recommended dose of Zydelig, you may be at increased risk of side effects with this medicine (see section 4, *Possible side effects*).

Contact your doctor or nearest emergency department immediately for advice. Keep the bottle and this leaflet with you so that you can easily describe what you have taken.

### **If you forget to take Zydelig**

Take care to not miss a dose of Zydelig. If you miss a dose by less than 6 hours, take the missed dose right away. Then take your next dose as usual. If you miss a dose by more than 6 hours, wait and take the next dose at your usual time.

### **Do not stop taking Zydelig**

Do not stop taking this medicine unless your doctor tells you to.

If you have any further questions on the use of this medicine, ask your doctor.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Some side effects could be serious.**

**STOP taking Zydelig and seek medical help immediately if you experience any of the following:**

- reddish patches on the trunk, small circumscribed changes in the colour of the skin, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (Stevens-Johnson syndrome, toxic epidermal necrolysis).
- Widespread rash, high body temperature and enlarged lymph nodes (DRESS syndrome).

**Other side effects**

**Very common side effects**

*(may affect more than 1 in 10 people)*

- diarrhoea/inflammation of the large intestine
- rash
- changes in the number of white blood cells
- infections
- fever

*Blood tests may also show:*

- increased blood levels of liver enzymes

**Common side effects**

*(may affect up to 1 in 10 people)*

- inflammation of the lungs
- liver damage

*Blood tests may also show:*

- increased blood levels of fats

**Reporting of side effects**

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Zydelig**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the bottle and carton after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

**What Zydelig contains**

- **The active substance is idelalisib.** Each film-coated tablet contains 150 mg of idelalisib.

- **The other ingredients are:**

*Tablet core:*

Microcrystalline cellulose, hydroxypropyl cellulose (E463), croscarmellose sodium, sodium starch glycolate, magnesium stearate.

*Film-coating:*

Polyvinyl alcohol (E1203), macrogol 3350 (E1521), titanium dioxide (E171), talc (E553B), iron oxide red (E172).

### **What Zydelig looks like and contents of the pack**

The film-coated tablets are pink, oval-shaped tablets, debossed on one side with “GSI” and “150” on the other side.

The following pack size is available: outer carton containing 1 plastic bottle of 60 film-coated tablets.

### **Marketing Authorisation Holder**

Gilead Sciences Ireland UC  
Carrigtohill  
County Cork, T45 DP77  
Ireland

### **Manufacturer**

Gilead Sciences Ireland UC  
IDA Business & Technology Park  
Carrigtohill  
County Cork  
Ireland

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **België/Belgique/Belgien**

Gilead Sciences Belgium SRL-BV  
Tél/Tel: + 32 (0) 24 01 35 50

#### **България**

Gilead Sciences Ireland UC  
Тел.: + 353 (0) 1 686 1888

#### **Česká republika**

Gilead Sciences s.r.o.  
Tel: + 420 910 871 986

#### **Danmark**

Gilead Sciences Sweden AB  
Tlf: + 46 (0) 8 5057 1849

#### **Deutschland**

Gilead Sciences GmbH  
Tel: + 49 (0) 89 899890-0

#### **Eesti**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

#### **Lietuva**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

#### **Luxembourg/Luxemburg**

Gilead Sciences Belgium SRL-BV  
Tél/Tel: + 32 (0) 24 01 35 50

#### **Magyarország**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

#### **Malta**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

#### **Nederland**

Gilead Sciences Netherlands B.V.  
Tel: + 31 (0) 20 718 36 98

#### **Norge**

Gilead Sciences Sweden AB  
Tlf: + 46 (0) 8 5057 1849

**Ελλάδα**

Gilead Sciences Ελλάς Μ.ΕΠΕ.  
Τηλ: + 30 210 8930 100

**España**

Gilead Sciences, S.L.  
Tel: + 34 91 378 98 30

**France**

Gilead Sciences  
Tél: + 33 (0) 1 46 09 41 00

**Hrvatska**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

**Ireland**

Gilead Sciences Ireland UC  
Tel: +353 (0) 214 825 999

**Ísland**

Gilead Sciences Sweden AB  
Sími: + 46 (0) 8 5057 1849

**Italia**

Gilead Sciences S.r.l.  
Tel: + 39 02 439201

**Κύπρος**

Gilead Sciences Ελλάς Μ.ΕΠΕ.  
Τηλ: + 30 210 8930 100

**Latvija**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

**Österreich**

Gilead Sciences GesmbH  
Tel: + 43 1 260 830

**Polska**

Gilead Sciences Poland Sp. z o.o.  
Tel: + 48 22 262 8702

**Portugal**

Gilead Sciences, Lda.  
Tel: + 351 21 7928790

**România**

Gilead Sciences (GSR) S.R.L.  
Tel: + 40 31 631 18 00

**Slovenija**

Gilead Sciences Ireland UC  
Tel: + 353 (0) 1 686 1888

**Slovenská republika**

Gilead Sciences Slovakia s.r.o.  
Tel: + 421 232 121 210

**Suomi/Finland**

Gilead Sciences Sweden AB  
Puh/Tel: + 46 (0) 8 5057 1849

**Sverige**

Gilead Sciences Sweden AB  
Tel: + 46 (0) 8 5057 1849

**United Kingdom (Northern Ireland)**

Gilead Sciences Ireland UC  
Tel: + 44 (0) 8000 113700

**This leaflet was last revised in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>.