



EUROPEAN MEDICINES AGENCY
SCIENCE MEDICINES HEALTH

21 July 2011
EMA/CHMP/557933/2011
Committee for medicinal products for human use (CHMP)

Summary of opinion¹ (post authorisation)

Enbrel etanercept

On 21 July 2011 the Committee for Medicinal Products for Human Use (CHMP) adopted a positive opinion recommending a variation to the terms of the marketing authorisation for the medicinal product Enbrel. The marketing authorisation holder for this medicinal product is Wyeth Europa Ltd. They may request a re-examination of the CHMP opinion, provided that they notify the European Medicines Agency in writing of their intention within 15 days of receipt of the opinion.

The CHMP adopted a change to an indication as follows²:

Polyarticular juvenile idiopathic arthritis

Treatment of active polyarticular juvenile idiopathic arthritis in children and adolescents from the age of **2** years who have had an inadequate response to, or who have proved intolerant of, methotrexate. Enbrel has not been studied in children aged less than **2** years.

Detailed conditions for the use of this product will be described in the updated summary of product characteristics (SmPC), which will be published in the revised European public assessment report (EPAR), and will be available in all official European Union languages after the variation to the marketing authorisation has been granted by the European Commission.

For information, the full indications for Enbrel will be as follows:

Rheumatoid arthritis

Enbrel in combination with methotrexate is indicated for the treatment of moderate to severe active rheumatoid arthritis in adults when the response to disease-modifying antirheumatic drugs, including methotrexate (unless contraindicated), has been inadequate.

Enbrel can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate.

¹ Summaries of positive opinion are published without prejudice to the commission decision, which will normally be issued within 44 days (Type II variations) and 67 days (Annex II applications) from adoption of the opinion.

² The text in bold represents the new or the amended indication.



Enbrel is also indicated in the treatment of severe, active and progressive rheumatoid arthritis in adults not previously treated with methotrexate.

Enbrel, alone or in combination with methotrexate, has been shown to reduce the rate of progression of joint damage as measured by X-ray and to improve physical function.

Polyarticular juvenile idiopathic arthritis

Treatment of active polyarticular juvenile idiopathic arthritis in children and adolescents from the age of 2 years who have had an inadequate response to, or who have proved intolerant of, methotrexate. Enbrel has not been studied in children aged less than 2 years.

Psoriatic arthritis

Treatment of active and progressive psoriatic arthritis in adults when the response to previous disease-modifying antirheumatic drug therapy has been inadequate. Enbrel has been shown to improve physical function in patients with psoriatic arthritis, and to reduce the rate of progression of peripheral joint damage as measured by X-ray in patients with polyarticular symmetrical subtypes of the disease.

Ankylosing spondylitis

Treatment of adults with severe active ankylosing spondylitis who have had an inadequate response to conventional therapy.

Plaque psoriasis

Treatment of adults with moderate to severe plaque psoriasis who failed to respond to, or who have a contraindication to, or are intolerant to other systemic therapy, including ciclosporin, methotrexate or psoralen and ultraviolet-A light (PUVA) (see section 5.1).

Paediatric plaque psoriasis

Treatment of chronic severe plaque psoriasis in children and adolescents from the age of 6 years who are inadequately controlled by, or are intolerant to, other systemic therapies or phototherapies.